

wwPDB X-ray Structure Validation Summary Report (i)

Nov 20, 2023 – 07:41 PM JST

PDB ID : 7DAH

Title : Adenosine triphosphate phosphoribosyltransferase from Vibrio cholerae in

complex with ATP and PRPP

Authors: Pal, R.K.; Gourinath, S.; Biswal, B.K.

Deposited on : 2020-10-16

Resolution : 2.92 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(\mathring{A}))$
R_{free}	130704	2307 (2.94-2.90)
Clashscore	141614	2531 (2.94-2.90)
Ramachandran outliers	138981	2462 (2.94-2.90)
Sidechain outliers	138945	2464 (2.94-2.90)
RSRZ outliers	127900	2248 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	310	72%	19%	• 7%
1	В	310	69%	22%	• 6%
1	С	310	69%	20%	• 7%
1	D	310	73%	18%	• 7%
1	E	310	75%	17%	• 5%
1	F	310	71%	17%	• 7%



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PEG	D	401	-	-	X	-



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 13159 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ATP phosphoribosyltransferase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	288	Total	С	N	О	S	0	0	0
1	A	200	2149	1362	372	399	16	0	0	0
1	В	292	Total	С	N	О	S	0	0	0
1	Б	292	2131	1361	358	398	14		U	
1	С	288	Total C N O S 0	0	0	0				
1		200	2134	1350	370	398	16	U	U	
1	D	287	Total	С	N	О	S	0	0	0
1	D	201	2142	1353	375	398	16	0	0	
1	Е	293	Total	С	N	О	S	0	0	0
1	12	290	2123	1350	365	394	14	U	0	0
1	F	287	Total	С	N	О	S	0	0	0
1	I.	201	2105	1335	363	393	14	U		U

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP C3LU29
A	-4	HIS	-	expression tag	UNP C3LU29
A	-3	HIS	-	expression tag	UNP C3LU29
A	-2	HIS	-	expression tag	UNP C3LU29
A	-1	HIS	-	expression tag	UNP C3LU29
A	0	HIS	-	expression tag	UNP C3LU29
В	-5	HIS	-	expression tag	UNP C3LU29
В	-4	HIS	-	expression tag	UNP C3LU29
В	-3	HIS	-	expression tag	UNP C3LU29
В	-2	HIS	-	expression tag	UNP C3LU29
В	-1	HIS	-	expression tag	UNP C3LU29
В	0	HIS	-	expression tag	UNP C3LU29
С	-5	HIS	-	expression tag	UNP C3LU29
С	-4	HIS	=	expression tag	UNP C3LU29
С	-3	HIS	-	expression tag	UNP C3LU29
С	-2	HIS	-	expression tag	UNP C3LU29
С	-1	HIS	-	expression tag	UNP C3LU29


Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	0	HIS	-	expression tag	UNP C3LU29
D	-5	HIS	-	expression tag	UNP C3LU29
D	-4	HIS	-	expression tag	UNP C3LU29
D	-3	HIS	-	expression tag	UNP C3LU29
D	-2	HIS	-	expression tag	UNP C3LU29
D	-1	HIS	-	expression tag	UNP C3LU29
D	0	HIS	-	expression tag	UNP C3LU29
E	-5	HIS	-	expression tag	UNP C3LU29
E	-4	HIS	-	expression tag	UNP C3LU29
Е	-3	HIS	-	expression tag	UNP C3LU29
Е	-2	HIS	-	expression tag	UNP C3LU29
Е	-1	HIS	-	expression tag	UNP C3LU29
E	0	HIS	-	expression tag	UNP C3LU29
F	-5	HIS	-	expression tag	UNP C3LU29
F	-4	HIS	-	expression tag	UNP C3LU29
F	-3	HIS	-	expression tag	UNP C3LU29
F	-2	HIS	-	expression tag	UNP C3LU29
F	-1	HIS	-	expression tag	UNP C3LU29
F	0	HIS	-	expression tag	UNP C3LU29

• Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 7 4 3	0	0


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total C O 7 4 3	0	0


• Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	С	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	Е	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0


• Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O_4P) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	D	1	Total O P 5 4 1	0	0

• Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0

Continued on next page...




Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total C O 6 3 3	0	0

• Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	С	1	Total Mg 1 1	0	0
6	F	1	Total Mg 1 1	0	0


• Molecule 7 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
7	С	1	Total	С	N	О	Р	0	0
'		1	31	10	5	13	3	U	
7	Г	1	Total	С	N	О	Р	0	0
1	1'	1	31	10	5	13	3	U	

• Molecule 8 is 1-O-pyrophosphono-5-O-phosphono-alpha-D-ribofuranose (three-letter code: PRP) (formula: $C_5H_{13}O_{14}P_3$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
8	С	1	Total	С	О	Р	0	0	
0		1	22	5	14	3	U		
Q	E	1	Total	С	О	Р	0	0	
0	I'	1	22	5	14	3	0	0	

• Molecule 9 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	35	Total O 35 35	0	0
9	В	27	Total O 27 27	0	0
9	С	47	Total O 47 47	0	0
9	D	42	Total O 42 42	0	0
9	E	22	Total O 22 22	0	0
9	F	26	Total O 26 26	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: ATP phosphoribosyltransferase



• Molecule 1: ATP phosphoribosyltransferase




• Molecule 1: ATP phosphoribosyltransferase









 \bullet Molecule 1: ATP phosphoribosyltransferase



• Molecule 1: ATP phosphoribosyltransferase



• Molecule 1: ATP phosphoribosyltransferase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 32	Depositor
Cell constants	136.60Å 136.60Å 121.93Å	D
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.32 - 2.92	Depositor
Resolution (A)	28.30 - 2.92	EDS
% Data completeness	99.6 (28.32-2.92)	Depositor
(in resolution range)	99.7 (28.30-2.92)	EDS
R_{merge}	(Not available)	Depositor
R_{sum}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.03 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
рρ.	0.216 , 0.268	Depositor
R, R_{free}	0.215 , 0.264	DCC
R_{free} test set	2553 reflections (4.63%)	wwPDB-VP
Wilson B-factor (Å ²)	61.3	Xtriage
Anisotropy	0.153	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.28, 23.7	EDS
L-test for twinning ²	$< L > = 0.52, < L^2> = 0.35$	Xtriage
	0.009 for -h,-k,l	
Estimated twinning fraction	0.470 for h,-h-k,-l	Xtriage
	0.010 for -k,-h,-l	
F_o, F_c correlation	0.94	EDS
Total number of atoms	13159	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ATP, PEG, EDO, GOL, PO4, PRP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bo	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z >5
1	A	0.80	0/2170	0.95	0/2924
1	В	0.80	1/2155~(0.0%)	0.97	0/2916
1	С	0.78	$1/2157 \ (0.0\%)$	0.94	0/2911
1	D	0.82	0/2164	0.97	0/2917
1	Е	0.80	0/2150	0.95	1/2912 (0.0%)
1	F	0.79	1/2129 (0.0%)	0.94	0/2876
All	All	0.80	$3/12925 \ (0.0\%)$	0.95	1/17456 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintenain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	С	0	1
1	D	0	1
All	All	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$Ideal(\AA)$
1	F	28	GLU	CD-OE1	5.36	1.31	1.25
1	В	276	GLU	CD-OE1	5.30	1.31	1.25
1	С	28	GLU	CD-OE1	5.12	1.31	1.25

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
1	Е	146	ARG	CB-CA-C	5.08	120.57	110.40



There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	С	257	THR	Peptide
1	D	259	LEU	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2149	0	2130	50	0
1	В	2131	0	2087	65	0
1	С	2134	0	2101	64	0
1	D	2142	0	2124	43	0
1	Ε	2123	0	2056	44	0
1	F	2105	0	2057	58	0
2	A	7	0	10	0	0
2	D	7	0	10	4	0
3	A	8	0	12	0	0
3	С	4	0	6	0	0
3	D	12	0	18	1	0
3	Ε	4	0	6	0	0
3	F	4	0	6	0	0
4	A	5	0	0	0	0
4	D	5	0	0	0	0
5	A	6	0	8	0	0
5	D	6	0	8	1	0
6	С	1	0	0	0	0
6	F	1	0	0	0	0
7	С	31	0	12	2	0
7	F	31	0	12	2	0
8	С	22	0	8	2	0
8	F	22	0	8	2	0
9	A	35	0	0	2	0
9	В	27	0	0	4	0
9	С	47	0	0	6	0
9	D	42	0	0	1	0
9	Ε	22	0	0	3	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
9	F	26	0	0	0	0
All	All	13159	0	12679	301	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 301 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:D:17:GLN:NE2	1:D:20:GLY:HA3	1.77	0.99
1:A:17:GLN:NE2	1:A:20:GLY:HA3	1.75	0.99
1:B:17:GLN:NE2	1:B:20:GLY:HA3	1.78	0.99
1:E:17:GLN:NE2	1:E:20:GLY:HA3	1.79	0.97
1:C:17:GLN:NE2	1:C:20:GLY:HA3	1.78	0.97

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	276/310 (89%)	260 (94%)	14 (5%)	2 (1%)	22	53
1	В	286/310 (92%)	271 (95%)	14 (5%)	1 (0%)	41	70
1	С	282/310 (91%)	269 (95%)	12 (4%)	1 (0%)	34	65
1	D	279/310 (90%)	262 (94%)	15 (5%)	2 (1%)	22	53
1	E	289/310 (93%)	275 (95%)	14 (5%)	0	100	100
1	F	283/310 (91%)	269 (95%)	14 (5%)	0	100	100
All	All	1695/1860 (91%)	1606 (95%)	83 (5%)	6 (0%)	34	65

5 of 6 Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	A	242	LYS
1	В	208	ALA
1	С	242	LYS
1	D	260	PRO
1	D	259	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percen	ntiles
1	A	$213/263\ (81\%)$	196 (92%)	17 (8%)	12	32
1	В	201/263~(76%)	174 (87%)	27 (13%)	4	11
1	С	207/263~(79%)	182 (88%)	25 (12%)	5	14
1	D	212/263 (81%)	198 (93%)	14 (7%)	16	42
1	E	197/263~(75%)	177 (90%)	20 (10%)	7	21
1	F	201/263~(76%)	177 (88%)	24 (12%)	5	14
All	All	1231/1578 (78%)	1104 (90%)	127 (10%)	7	21

5 of 127 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	159	SER
1	F	99	THR
1	D	43	ARG
1	F	97	GLU
1	F	192	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	96	ASN
1	F	17	GLN
1	F	226	GLN
1	F	27	GLN

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	Ε	93	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 20 ligands modelled in this entry, 2 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Т	Clasica	Res	Link	Во	ond leng	ths	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PEG	D	401	-	6,6,6	0.23	0	5,5,5	0.18	0
3	EDO	D	402	-	3,3,3	0.14	0	2,2,2	0.03	0
2	PEG	A	401	-	6,6,6	0.19	0	5,5,5	0.23	0
5	GOL	A	405	-	5,5,5	0.18	0	5,5,5	0.34	0
3	EDO	С	404	-	3,3,3	0.11	0	2,2,2	0.13	0
4	PO4	A	404	-	4,4,4	1.47	1 (25%)	6,6,6	0.36	0
8	PRP	С	403	-	19,22,22	0.67	0	33,35,35	1.35	4 (12%)
8	PRP	F	403	-	19,22,22	0.71	0	33,35,35	1.26	2 (6%)
3	EDO	D	405	-	3,3,3	0.41	0	2,2,2	0.21	0
4	PO4	D	406	-	4,4,4	0.71	0	6,6,6	0.58	0
7	ATP	С	402	6	26,33,33	0.61	0	31,52,52	1.11	2 (6%)
3	EDO	F	402	-	3,3,3	0.37	0	2,2,2	0.42	0



Mal	Mol Type Chain		Res	Link	Bo	nd leng	$ ag{ths}$	Bond angles		
MIOI	Туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	402	-	3,3,3	0.27	0	2,2,2	0.27	0
5	GOL	D	403	-	5,5,5	0.17	0	5,5,5	0.36	0
3	EDO	Е	401	_	3,3,3	0.20	0	2,2,2	0.26	0
7	ATP	F	404	6	26,33,33	0.62	0	31,52,52	1.05	1 (3%)
3	EDO	D	404	-	3,3,3	0.23	0	2,2,2	0.26	0
3	EDO	A	403	-	3,3,3	0.27	0	2,2,2	0.13	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PEG	D	401	-	-	3/4/4/4	-
3	EDO	D	402	-	-	0/1/1/1	-
2	PEG	A	401	-	-	0/4/4/4	-
5	GOL	A	405	-	-	1/4/4/4	-
3	EDO	С	404	-	-	1/1/1/1	-
8	PRP	С	403	-	-	3/16/33/33	0/1/1/1
8	PRP	F	403	-	-	4/16/33/33	0/1/1/1
3	EDO	D	405	-	-	1/1/1/1	-
7	ATP	С	402	6	-	7/18/38/38	0/3/3/3
3	EDO	F	402	-	-	0/1/1/1	-
3	EDO	A	402	-	-	0/1/1/1	-
5	GOL	D	403	-	-	3/4/4/4	-
3	EDO	Е	401	-	-	1/1/1/1	-
7	ATP	F	404	6	-	6/18/38/38	0/3/3/3
3	EDO	D	404	-	-	0/1/1/1	-
3	EDO	A	403	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$[Ideal(\AA)]$
4	Α	404	PO4	P-O1	2.92	1.57	1.50

The worst 5 of 9 bond angle outliers are listed below:

	Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
	8	С	403	PRP	O5-P-O1P	-4.10	94.98	106.47
Ī	8	F	403	PRP	O2P-P-O5	-3.27	98.04	106.73

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
8	С	403	PRP	O1-C1-C2	2.83	111.52	106.72
7	F	404	ATP	C5-C6-N6	2.63	124.35	120.35
7	С	402	ATP	C5-C6-N6	2.63	124.34	120.35

There are no chirality outliers.

5 of 31 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	403	GOL	C1-C2-C3-O3
5	D	403	GOL	O2-C2-C3-O3
7	С	402	ATP	PB-O3B-PG-O3G
7	С	402	ATP	C5'-O5'-PA-O3A
7	С	402	ATP	O4'-C4'-C5'-O5'


There are no ring outliers.


7 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	401	PEG	4	0
3	D	402	EDO	1	0
8	С	403	PRP	2	0
8	F	403	PRP	2	0
7	С	402	ATP	2	0
5	D	403	GOL	1	0
7	F	404	ATP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ} {>} 2$	$\mathrm{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	A	288/310~(92%)	-0.35	1 (0%) 94 94	39, 62, 110, 136	1 (0%)
1	В	292/310~(94%)	-0.23	0 100 100	45, 78, 111, 128	1 (0%)
1	С	$288/310 \ (92\%)$	-0.33	1 (0%) 94 94	43, 67, 109, 137	0
1	D	287/310 (92%)	-0.35	0 100 100	38, 61, 108, 130	0
1	Е	293/310 (94%)	-0.26	1 (0%) 94 94	47, 78, 112, 125	1 (0%)
1	F	287/310 (92%)	-0.35	1 (0%) 94 94	42, 68, 108, 138	1 (0%)
All	All	1735/1860 (93%)	-0.31	4 (0%) 95 95	38, 69, 111, 138	4 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	170	ASP	2.8
1	A	237	HIS	2.1
1	Ε	261	LEU	2.1
1	С	93	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
4	PO4	D	406	5/5	0.82	0.12	106,116,123,123	0
8	PRP	F	403	22/22	0.82	0.21	106,168,191,198	0
3	EDO	E	401	4/4	0.84	0.24	89,89,89,90	0
8	PRP	С	403	22/22	0.85	0.19	104,159,200,202	0
3	EDO	D	405	4/4	0.85	0.14	71,76,81,87	0
7	ATP	F	404	31/31	0.86	0.16	78,97,116,118	12
3	EDO	F	402	4/4	0.87	0.25	77,82,85,85	0
3	EDO	D	402	4/4	0.88	0.31	91,96,96,97	0
2	PEG	A	401	7/7	0.89	0.15	84,87,89,90	0
2	PEG	D	401	7/7	0.89	0.17	80,89,102,102	0
7	ATP	С	402	31/31	0.90	0.16	81,111,120,124	1
5	GOL	D	403	6/6	0.91	0.17	67,68,70,71	0
4	PO4	A	404	5/5	0.92	0.09	97,104,113,114	0
3	EDO	D	404	4/4	0.92	0.26	69,75,78,79	0
3	EDO	С	404	4/4	0.92	0.17	91,91,92,92	0
6	MG	F	401	1/1	0.92	0.09	76,76,76,76	0
3	EDO	A	402	4/4	0.94	0.15	74,75,77,77	0
5	GOL	A	405	6/6	0.95	0.21	69,72,77,80	0
6	MG	С	401	1/1	0.98	0.06	71,71,71,71	0
3	EDO	A	403	4/4	0.98	0.17	74,75,76,77	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around PO4 D 406: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





Electron density around PRP F 403: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

Electron density around EDO E 401:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around PRP C 403:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO D 405:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around ATP F 404:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO F 402:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO D 402: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)


Electron density around PEG A 401: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around PEG D 401:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around ATP C 402:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
















Electron density around PO4 A 404:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO D 404:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around EDO C 404:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around EDO A 402:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around GOL A 405:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)
















6.5 Other polymers (i)

There are no such residues in this entry.

