



# Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2020 – 11:58 pm BST

PDB ID : 6DFD  
Title : Crystal structure of CNNM3 cyclic nucleotide-binding homology domain  
Authors : Kozlov, G.; Gehring, K.  
Deposited on : 2018-05-14  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	
1	B	260	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4240 atoms, of which 2097 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metal transporter CNNM3.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S	Se			
1	A	137	2251	721	1135	190	200	1	4	0	0	0
1	B	125	1919	621	962	163	169	1	3	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	448	GLY	-	expression tag	UNP Q8NE01
A	449	PRO	-	expression tag	UNP Q8NE01
A	450	LEU	-	expression tag	UNP Q8NE01
A	451	GLY	-	expression tag	UNP Q8NE01
A	452	SER	-	expression tag	UNP Q8NE01
A	516	MSE	ILE	engineered mutation	UNP Q8NE01
A	591	MSE	THR	engineered mutation	UNP Q8NE01
A	623	MSE	ALA	engineered mutation	UNP Q8NE01
A	651	MSE	LEU	engineered mutation	UNP Q8NE01
A	669	MSE	VAL	engineered mutation	UNP Q8NE01
A	670	MSE	ILE	engineered mutation	UNP Q8NE01
B	448	GLY	-	expression tag	UNP Q8NE01
B	449	PRO	-	expression tag	UNP Q8NE01
B	450	LEU	-	expression tag	UNP Q8NE01
B	451	GLY	-	expression tag	UNP Q8NE01
B	452	SER	-	expression tag	UNP Q8NE01
B	516	MSE	ILE	engineered mutation	UNP Q8NE01
B	591	MSE	THR	engineered mutation	UNP Q8NE01
B	623	MSE	ALA	engineered mutation	UNP Q8NE01
B	651	MSE	LEU	engineered mutation	UNP Q8NE01
B	669	MSE	VAL	engineered mutation	UNP Q8NE01
B	670	MSE	ILE	engineered mutation	UNP Q8NE01

- Molecule 2 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
2	A	64	Total O 64 64	0	0
2	B	6	Total O 6 6	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	101.27Å 101.27Å 77.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	32.47 – 1.90 32.47 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (32.47-1.90) 99.6 (32.47-1.90)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.49 (at 1.91Å)	Xtrriage
Refinement program	PHENIX (1.11.1_2575: ???)	Depositor
R, $R_{free}$	0.219 , 0.239 0.220 , 0.239	Depositor DCC
$R_{free}$ test set	1587 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.0	Xtrriage
Anisotropy	0.304	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 59.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4240	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.32	0/1135	0.53	0/1530
1	B	0.26	0/971	0.46	0/1314
All	All	0.29	0/2106	0.50	0/2844

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1116	1135	1133	1	0
1	B	957	962	927	2	0
2	A	64	0	0	0	1
2	B	6	0	0	0	0
All	All	2143	2097	2060	3	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:565:ARG:NH1	1:B:578:GLU:OE2	2.25	0.68

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:569:GLU:HA	1:A:575:LEU:O	2.17	0.44
1:B:650:LEU:O	1:B:652:ALA:N	2.54	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:854:HOH:O	2:A:854:HOH:O[7_555]	1.57	0.63

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	133/260 (51%)	128 (96%)	5 (4%)	0	100	100
1	B	121/260 (46%)	119 (98%)	2 (2%)	0	100	100
All	All	254/520 (49%)	247 (97%)	7 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/225 (55%)	121 (98%)	2 (2%)	62	60

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	95/225 (42%)	95 (100%)	0	100	100
All	All	218/450 (48%)	216 (99%)	2 (1%)	78	79

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	562	LEU
1	A	569	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	133/260 (51%)	0.69	19 (14%) <b>2</b>   <b>2</b>	30, 44, 76, 95	0
1	B	122/260 (46%)	1.22	27 (22%) <b>0</b>   <b>0</b>	45, 74, 110, 130	0
All	All	255/520 (49%)	0.95	46 (18%) <b>1</b>   <b>1</b>	30, 58, 102, 130	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	653	THR	6.5
1	B	522	LEU	6.4
1	A	488	TYR	5.3
1	B	544	THR	5.0
1	B	542	LEU	4.5
1	B	496	LEU	4.0
1	B	543	ALA	3.9
1	B	521	LEU	3.8
1	B	552	SER	3.6
1	B	536	PHE	3.5
1	B	540	ASN	3.4
1	B	499	ALA	3.4
1	B	539	SER	3.3
1	A	552	SER	3.2
1	B	498	LEU	3.1
1	B	650	LEU	3.0
1	A	577	PHE	3.0
1	A	652	ALA	2.9
1	B	625	CYS	2.9
1	B	652	ALA	2.9
1	A	582	PHE	2.9
1	A	562	LEU	2.8
1	A	553	GLN	2.8
1	B	541	ARG	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	624	TYR	2.5
1	B	637	GLN	2.5
1	A	654	ARG	2.4
1	B	527	HIS	2.4
1	B	545	HIS	2.4
1	A	496	LEU	2.4
1	A	560	LEU	2.4
1	B	497	LEU	2.4
1	B	519	LYS	2.3
1	B	530	VAL	2.2
1	A	625	CYS	2.2
1	A	585	TYR	2.2
1	B	553	GLN	2.2
1	B	515	ARG	2.2
1	A	639	ILE	2.2
1	A	579	ASN	2.2
1	A	655	ALA	2.1
1	B	537	ASP	2.1
1	A	568	VAL	2.1
1	B	506	ARG	2.1
1	B	555	VAL	2.0
1	A	581	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.