



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 6, 2018 – 11:10 AM EDT

PDB ID : 6DI4  
Title : Rational Modification of Vanillin Derivatives to Stereospecifically Destabilize Sickle Hemoglobin Polymer Formation  
Authors : Pagare, P.P.; Musayev, F.N.  
Deposited on : 2018-05-22  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtrriage (Phenix) : 1.13  
EDS : rb-20031633  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031633


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	5502 (1.90-1.90)
Clashscore	122126	6115 (1.90-1.90)
Ramachandran outliers	120053	6048 (1.90-1.90)
Sidechain outliers	120020	6048 (1.90-1.90)
RSRZ outliers	108989	5379 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	141	 89% 10%
1	C	141	 89% 11%
2	B	146	 82% 18%
2	D	146	 77% 21%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Hemoglobin subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	141	1068	685	187	193	3	0	0	0
1	C	141	1068	685	187	193	3	0	0	0


- Molecule 2 is a protein called Hemoglobin subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	146	1122	724	195	200	3	0	0	0
2	D	146	1122	724	195	200	3	0	0	0

- Molecule 3 is CARBON MONOXIDE (three-letter code: CMO) (formula: CO).







Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			19	15	1	3		
5	C	1	Total	C	N	O	0	0
			19	15	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	158	Total	O	0	0
			158	158		
6	B	226	Total	O	0	0
			226	226		
6	C	196	Total	O	0	0
			196	196		
6	D	204	Total	O	0	0
			204	204		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemoglobin subunit alpha




- Molecule 1: Hemoglobin subunit alpha



- Molecule 2: Hemoglobin subunit beta



- Molecule 2: Hemoglobin subunit beta



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.70Å 83.40Å 104.93Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.34 – 1.90 29.34 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.2 (29.34-1.90) 97.3 (29.34-1.90)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.18 (at 1.91Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.195 , 0.249 0.187 , 0.240	Depositor DCC
$R_{free}$ test set	2133 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.9	Xtrriage
Anisotropy	0.207	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 37.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5382	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CMO, HEM, GJ1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/1096	0.53	0/1491
1	C	0.30	0/1096	0.52	0/1491
2	B	0.30	0/1152	0.53	0/1566
2	D	0.60	4/1152 (0.3%)	0.79	4/1566 (0.3%)
All	All	0.40	4/4496 (0.1%)	0.60	4/6114 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	LEU	CG-CD2	-9.25	1.17	1.51
2	D	3	LEU	C-O	-6.83	1.10	1.23
2	D	4	THR	C-O	-6.31	1.11	1.23
2	D	3	LEU	CA-CB	-5.33	1.41	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	3	LEU	CB-CG-CD2	-18.39	79.75	111.00
2	D	3	LEU	CA-CB-CG	-6.29	100.84	115.30
2	D	3	LEU	CB-CG-CD1	5.36	120.11	111.00
2	D	1	VAL	O-C-N	-5.13	114.50	122.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1068	0	1071	20	0
1	C	1068	0	1071	13	0
2	B	1122	0	1118	22	0
2	D	1122	0	1118	38	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	43	0	30	0	0
4	B	43	0	30	0	0
4	C	43	0	30	0	0
4	D	43	0	30	0	0
5	A	19	0	0	0	0
5	C	19	0	0	0	0
6	A	158	0	0	2	0
6	B	226	0	0	5	0
6	C	196	0	0	3	0
6	D	204	0	0	9	0
All	All	5382	0	4498	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:141:ARG:HH11	1:A:141:ARG:HG3	0.96	1.13
1:A:141:ARG:NH1	1:A:141:ARG:HG3	1.70	0.98
2:D:2:HIS:O	2:D:3:LEU:CB	2.29	0.77
1:A:141:ARG:HH11	1:A:141:ARG:CG	1.89	0.68
2:B:143:HIS:HA	2:B:146:HIS:CD2	2.28	0.68
2:D:2:HIS:O	6:D:301:HOH:O	2.11	0.68
2:B:18:VAL:HG13	2:B:23:VAL:HG21	1.83	0.61
1:A:75:ASP:HB2	6:A:421:HOH:O	2.00	0.61
1:A:85:ASP:O	1:A:89:HIS:HB2	2.02	0.60
2:D:139:ASN:HB3	6:D:332:HOH:O	2.00	0.60
1:A:141:ARG:NH1	1:A:141:ARG:CG	2.54	0.59
2:D:100:PRO:HD3	2:D:145:TYR:CE1	2.37	0.58
1:A:92:ARG:HH11	2:D:39:GLN:HE21	1.51	0.58

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1:VAL:HG21	2:B:78:LEU:HD22	1.85	0.58
2:D:75:LEU:HD21	2:D:133:VAL:HG11	1.85	0.58
2:D:3:LEU:N	6:D:303:HOH:O	2.25	0.57
2:D:18:VAL:HG13	2:D:23:VAL:HG21	1.86	0.57
1:C:47:ASP:H	1:C:54:GLN:NE2	2.02	0.56
2:D:75:LEU:HD23	2:D:78:LEU:HD21	1.88	0.56
2:D:26:GLU:HG3	6:D:394:HOH:O	2.04	0.56
1:A:52:SER:O	1:A:56:LYS:HG2	2.05	0.56
1:A:119:PRO:HB3	2:B:55:MET:CE	2.37	0.54
1:A:3:SER:HB3	1:C:140:TYR:HA	1.90	0.54
2:D:143:HIS:HA	2:D:146:HIS:CD2	2.43	0.54
1:C:85:ASP:O	1:C:89:HIS:HB2	2.07	0.53
2:D:2:HIS:O	2:D:3:LEU:HB2	2.05	0.53
2:B:144:LYS:HD2	6:B:312:HOH:O	2.08	0.53
2:D:120:LYS:HG2	6:D:357:HOH:O	2.09	0.52
2:D:3:LEU:HB3	6:D:301:HOH:O	2.08	0.52
2:D:11:VAL:HG11	2:D:75:LEU:HD22	1.92	0.52
1:C:21:ALA:HB1	1:C:63:ALA:HB1	1.91	0.51
2:B:90:GLU:HG2	2:B:144:LYS:HE2	1.91	0.51
1:A:85:ASP:HA	1:A:89:HIS:CD2	2.46	0.51
1:A:72:HIS:HB3	1:A:75:ASP:OD2	2.11	0.51
2:B:140:ALA:O	2:B:143:HIS:ND1	2.40	0.51
2:B:37:TRP:HA	1:C:92:ARG:HB3	1.92	0.51
2:D:3:LEU:CB	6:D:301:HOH:O	2.58	0.51
2:B:86:ALA:O	2:B:90:GLU:HG3	2.12	0.50
1:C:141:ARG:HG3	6:C:389:HOH:O	2.12	0.50
1:A:92:ARG:NH1	2:D:39:GLN:HE21	2.10	0.49
1:C:85:ASP:OD1	1:C:141:ARG:HD3	2.11	0.49
2:D:90:GLU:CG	2:D:144:LYS:HE2	2.43	0.49
1:A:119:PRO:HB3	2:B:55:MET:HE2	1.93	0.49
1:A:43:PHE:HB3	1:A:46:PHE:HB2	1.95	0.49
1:A:60:LYS:HE3	1:A:64:ASP:OD2	2.13	0.49
2:B:8:LYS:HD2	6:B:388:HOH:O	2.12	0.48
1:C:112:HIS:C	1:C:113:LEU:HD12	2.34	0.48
2:B:14:LEU:HD11	2:B:118:PHE:CG	2.50	0.46
1:C:114:PRO:HA	2:D:116:HIS:CD2	2.51	0.46
1:C:107:VAL:HG13	2:D:115:ALA:HB2	1.97	0.46
2:D:142:ALA:O	2:D:145:TYR:HB2	2.16	0.46
2:D:90:GLU:HG2	2:D:144:LYS:HE2	1.98	0.46
2:D:3:LEU:CD2	2:D:8:LYS:HG3	2.46	0.46
2:D:3:LEU:HA	6:D:303:HOH:O	2.16	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:LYS:NZ	6:A:302:HOH:O	2.48	0.46
2:D:4:THR:OG1	2:D:7:GLU:HG3	2.16	0.46
1:A:140:TYR:HA	1:C:3:SER:HB3	1.98	0.45
2:B:63:HIS:HD2	6:B:418:HOH:O	1.98	0.45
1:C:30:GLU:HG2	6:C:414:HOH:O	2.16	0.45
2:B:142:ALA:O	2:B:145:TYR:HB2	2.17	0.45
2:D:82:LYS:HE3	2:D:143:HIS:CD2	2.52	0.44
2:D:68:LEU:HD13	2:D:68:LEU:O	2.18	0.44
2:B:124:PRO:HB2	2:B:125:PRO:HD3	2.00	0.43
2:B:143:HIS:NE2	2:B:144:LYS:NZ	2.67	0.43
2:B:20:VAL:HG22	6:B:396:HOH:O	2.18	0.42
1:C:60:LYS:HE2	6:C:392:HOH:O	2.19	0.42
2:D:123:THR:OG1	2:D:125:PRO:HD2	2.19	0.42
2:B:32:LEU:HD22	2:B:39:GLN:HA	2.03	0.41
2:D:3:LEU:CA	6:D:303:HOH:O	2.69	0.41
2:B:144:LYS:HA	2:B:144:LYS:HD3	1.87	0.41
2:B:19:ASN:ND2	6:B:307:HOH:O	2.54	0.41
1:A:119:PRO:HB3	2:B:55:MET:HE3	2.03	0.41
2:D:32:LEU:HD22	2:D:39:GLN:HA	2.03	0.41
2:D:11:VAL:CG1	2:D:75:LEU:HD22	2.51	0.41
2:B:127:GLN:O	2:B:131:GLN:HG2	2.20	0.41
2:D:89:SER:CB	2:D:144:LYS:HD3	2.51	0.40
2:D:89:SER:OG	2:D:144:LYS:HD3	2.21	0.40
2:D:133:VAL:O	2:D:137:VAL:HG23	2.21	0.40
2:D:82:LYS:HE3	2:D:143:HIS:HD2	1.86	0.40
1:A:92:ARG:HH11	2:D:39:GLN:NE2	2.19	0.40
2:D:124:PRO:HB2	2:D:125:PRO:HD3	2.04	0.40
2:D:89:SER:HB3	2:D:144:LYS:HD3	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/141 (99%)	138 (99%)	1 (1%)	0	100	100
1	C	139/141 (99%)	137 (99%)	2 (1%)	0	100	100
2	B	144/146 (99%)	140 (97%)	4 (3%)	0	100	100
2	D	144/146 (99%)	141 (98%)	3 (2%)	0	100	100
All	All	566/574 (99%)	556 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/113 (100%)	112 (99%)	1 (1%)	81	81
1	C	113/113 (100%)	113 (100%)	0	100	100
2	B	118/118 (100%)	118 (100%)	0	100	100
2	D	118/118 (100%)	117 (99%)	1 (1%)	83	83
All	All	462/462 (100%)	460 (100%)	2 (0%)	92	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	ARG
2	D	3	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	ASN
1	C	54	GLN
1	C	89	HIS
2	D	39	GLN
2	D	117	HIS
2	D	146	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CMO	A	201	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	A	202	1,3	27,50,50	1.64	7 (25%)	17,82,82	1.09	1 (5%)
5	GJ1	A	203	1	20,20,20	1.91	8 (40%)	26,26,26	1.61	6 (23%)
3	CMO	B	201	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	B	202	3,2	27,50,50	1.73	6 (22%)	17,82,82	0.86	1 (5%)
3	CMO	C	201	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	C	202	1,3	27,50,50	1.54	5 (18%)	17,82,82	1.01	1 (5%)
5	GJ1	C	203	1	20,20,20	2.04	12 (60%)	26,26,26	1.60	6 (23%)
3	CMO	D	201	4	0,1,1	0.00	-	0,0,0	0.00	-
4	HEM	D	202	3,2	27,50,50	1.59	6 (22%)	17,82,82	1.09	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CMO	A	201	4	-	0/0/0/0	0/0/0/0
4	HEM	A	202	1,3	-	0/6/54/54	0/0/8/8
5	GJ1	A	203	1	-	0/9/9/9	0/2/2/2
3	CMO	B	201	4	-	0/0/0/0	0/0/0/0
4	HEM	B	202	3,2	-	0/6/54/54	0/0/8/8
3	CMO	C	201	4	-	0/0/0/0	0/0/0/0
4	HEM	C	202	1,3	-	0/6/54/54	0/0/8/8
5	GJ1	C	203	1	-	0/9/9/9	0/2/2/2
3	CMO	D	201	4	-	0/0/0/0	0/0/0/0
4	HEM	D	202	3,2	-	0/6/54/54	0/0/8/8

All (44) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	202	HEM	C3C-CAC	-4.19	1.39	1.47
4	C	202	HEM	C3B-CAB	-4.16	1.39	1.47
4	D	202	HEM	C3C-CAC	-3.87	1.39	1.47
4	A	202	HEM	C3C-CAC	-3.86	1.40	1.47
4	D	202	HEM	C3B-CAB	-3.83	1.40	1.47
4	A	202	HEM	C3B-CAB	-3.83	1.40	1.47
4	B	202	HEM	C3B-CAB	-3.57	1.40	1.47
4	B	202	HEM	C3B-C2B	-3.32	1.35	1.40
4	C	202	HEM	C3C-CAC	-3.29	1.41	1.47
4	A	202	HEM	C3B-C2B	-3.05	1.36	1.40
4	D	202	HEM	C3B-C2B	-2.95	1.36	1.40
4	B	202	HEM	C3C-C2C	-2.81	1.36	1.40
4	D	202	HEM	C3C-C2C	-2.58	1.36	1.40
4	C	202	HEM	C3B-C2B	-2.25	1.37	1.40
4	B	202	HEM	CBB-CAB	2.02	1.42	1.29
5	C	203	GJ1	C14-C13	2.06	1.42	1.38
5	C	203	GJ1	O8-C4	2.07	1.42	1.37
5	A	203	GJ1	C6-C7	2.10	1.44	1.39
4	A	202	HEM	CBC-CAC	2.12	1.43	1.29
4	A	202	HEM	C1C-C2C	2.14	1.47	1.42
4	D	202	HEM	CBC-CAC	2.17	1.43	1.29
5	C	203	GJ1	C5-C4	2.17	1.43	1.38
4	A	202	HEM	CBB-CAB	2.18	1.43	1.29
5	C	203	GJ1	C18-C16	2.20	1.54	1.51
4	A	202	HEM	C4A-NA	2.20	1.40	1.36
4	B	202	HEM	CBC-CAC	2.20	1.44	1.29
5	A	203	GJ1	C7-C2	2.23	1.44	1.39
5	A	203	GJ1	C12-N17	2.28	1.38	1.34
5	C	203	GJ1	C12-N17	2.28	1.38	1.34

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	202	HEM	CBB-CAB	2.30	1.44	1.29
5	A	203	GJ1	C3-C4	2.32	1.43	1.38
5	C	203	GJ1	C6-C7	2.32	1.44	1.39
5	C	203	GJ1	C7-C2	2.32	1.45	1.39
4	C	202	HEM	CBB-CAB	2.33	1.44	1.29
5	A	203	GJ1	C16-N17	2.37	1.38	1.34
4	C	202	HEM	CBC-CAC	2.38	1.45	1.29
5	C	203	GJ1	C16-N17	2.42	1.38	1.34
5	A	203	GJ1	O10-C7	2.42	1.42	1.37
5	C	203	GJ1	C3-C2	2.44	1.43	1.39
5	A	203	GJ1	C3-C2	2.45	1.43	1.39
5	C	203	GJ1	C6-C5	2.48	1.43	1.38
5	C	203	GJ1	O10-C7	2.60	1.42	1.37
5	A	203	GJ1	C5-C4	2.72	1.44	1.38
5	C	203	GJ1	C3-C4	3.05	1.44	1.38

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	203	GJ1	C15-C16-N17	-2.38	119.35	122.40
4	C	202	HEM	C4C-C3C-C2C	-2.29	105.30	106.90
4	A	202	HEM	C4C-C3C-C2C	-2.28	105.31	106.90
4	D	202	HEM	CMA-C3A-C4A	-2.24	125.02	128.46
5	A	203	GJ1	C15-C16-N17	-2.24	119.53	122.40
5	C	203	GJ1	C13-C12-N17	-2.18	119.61	122.40
5	A	203	GJ1	C13-C12-N17	-2.15	119.64	122.40
4	B	202	HEM	CAD-C3D-C2D	-2.05	123.16	129.00
5	C	203	GJ1	C11-C12-N17	2.13	120.11	115.74
5	A	203	GJ1	C11-C12-N17	2.60	121.08	115.74
5	A	203	GJ1	C12-N17-C16	3.26	122.14	118.14
5	A	203	GJ1	C11-O10-C7	3.27	124.30	117.74
5	C	203	GJ1	C11-O10-C7	3.40	124.55	117.74
5	C	203	GJ1	C18-C16-N17	3.66	119.15	115.16
5	C	203	GJ1	C12-N17-C16	3.75	122.73	118.14
5	A	203	GJ1	C18-C16-N17	4.02	119.54	115.16

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	141/141 (100%)	-0.02	1 (0%) 87 89	22, 29, 42, 56	0
1	C	141/141 (100%)	-0.04	0 100 100	22, 29, 42, 46	0
2	B	146/146 (100%)	0.31	5 (3%) 45 49	23, 31, 49, 61	0
2	D	146/146 (100%)	0.22	4 (2%) 54 58	21, 31, 45, 71	0
All	All	574/574 (100%)	0.12	10 (1%) 70 73	21, 30, 45, 71	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	1	VAL	11.4
2	B	2	HIS	5.9
2	B	143	HIS	5.4
2	B	1	VAL	4.2
2	D	2	HIS	3.2
2	B	5	PRO	3.1
1	A	50	HIS	3.0
2	B	146	HIS	2.8
2	D	145	TYR	2.1
2	D	143	HIS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GJ1	A	203	19/19	0.62	0.32	41,46,50,51	0
5	GJ1	C	203	19/19	0.72	0.26	35,42,50,53	0
4	HEM	B	202	43/43	0.93	0.12	25,29,41,44	0
4	HEM	D	202	43/43	0.95	0.11	21,27,36,47	0
4	HEM	C	202	43/43	0.96	0.10	21,26,37,50	0
4	HEM	A	202	43/43	0.96	0.10	19,26,38,43	0
3	CMO	B	201	2/2	0.96	0.09	28,28,28,29	0
3	CMO	D	201	2/2	0.98	0.14	28,28,28,32	0
3	CMO	C	201	2/2	0.98	0.10	25,25,25,32	0
3	CMO	A	201	2/2	0.99	0.12	25,25,25,33	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.