



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 06:41 PM EDT

PDB ID : 5DJ1
Title : Structure of the PLP-Dependent L-Arginine Hydroxylase MppP Holoenzyme
Authors : Silvaggi, N.R.; Han, L.
Deposited on : 2015-09-01
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	376	 2% 87% 6% 7%
1	B	376	 2% 89% 5% 6%
1	C	376	 2% 88% 6% 6%
1	D	376	 6% 92% 5% 5%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 22818 atoms, of which 10799 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PLP-Dependent L-Arginine Hydroxylase MppP.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P	S			
1	A	351	5432	1729	2682	492	521	1	7	0	4	0
1	B	353	5419	1726	2674	487	524	1	7	0	0	0
1	C	353	5419	1726	2674	487	524	1	7	0	0	0
1	D	364	5620	1795	2769	504	544	1	7	0	2	0

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	B	1	Total	Cl	0	0
			1	1		
2	C	1	Total	Cl	0	0
			1	1		
2	D	1	Total	Cl	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	242	Total	O	0	0
			242	242		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	235	Total 235	O 235	0	0
4	C	223	Total 223	O 223	0	0
4	D	223	Total 223	O 223	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PLP-Dependent L-Arginine Hydroxylase MppP



- Molecule 1: PLP-Dependent L-Arginine Hydroxylase MppP




- Molecule 1: PLP-Dependent L-Arginine Hydroxylase MppP



- Molecule 1: PLP-Dependent L-Arginine Hydroxylase MppP





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.72Å 108.28Å 195.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.45 – 2.10 41.45 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (41.45-2.10) 99.8 (41.45-2.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.75 (at 2.10Å)	Xtrriage
Refinement program	PHENIX (1.10_2148: ???)	Depositor
R, R_{free}	0.144 , 0.171 0.146 , 0.173	Depositor DCC
R_{free} test set	5257 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	24.0	Xtrriage
Anisotropy	0.030	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 59.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	22818	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL, LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/2793	0.62	1/3801 (0.0%)
1	B	0.52	0/2778	0.63	0/3782
1	C	0.49	0/2778	0.61	0/3782
1	D	0.50	0/2893	0.63	0/3940
All	All	0.50	0/11242	0.62	1/15305 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	285	ARG	NE-CZ-NH1	6.15	123.38	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2750	2682	2688	21	0
1	B	2745	2674	2689	12	0
1	C	2745	2674	2690	15	0
1	D	2851	2769	2779	13	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	B	1	0	0	0	0
4	A	242	0	0	10	2
4	B	235	0	0	6	0
4	C	223	0	0	5	2
4	D	223	0	0	3	0
All	All	12019	10799	10846	59	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:23:LEU:N	4:C:601:HOH:O	2.05	0.89
1:A:345:SER:OG	4:A:601:HOH:O	1.95	0.84
1:A:176:GLU:OE1	4:A:602:HOH:O	2.00	0.80
1:D:147:LEU:O	4:D:601:HOH:O	2.08	0.71
1:A:123[A]:ARG:HD3	4:A:603:HOH:O	1.90	0.70
1:B:54:GLN:HB3	4:B:665:HOH:O	1.92	0.70
1:C:180:GLU:OE2	4:C:603:HOH:O	2.09	0.70
1:A:173:ARG:NH1	4:A:602:HOH:O	2.25	0.70
1:B:318:THR:HG22	1:B:321:GLU:H	1.62	0.65
1:A:54:GLN:NE2	4:A:608:HOH:O	2.30	0.64
1:A:80:GLU:OE1	4:A:604:HOH:O	2.15	0.64
1:D:29[B]:HIS:HE1	1:D:31:ARG:HE	1.44	0.63
1:D:374:GLU:O	4:D:603:HOH:O	2.16	0.62
1:C:123:ARG:NH2	4:C:602:HOH:O	2.05	0.62
1:D:120:ASP:OD1	1:D:123:ARG:NH2	2.34	0.59
1:D:29[B]:HIS:CE1	1:D:31:ARG:HE	2.20	0.59
1:D:164:ARG:NH2	4:D:602:HOH:O	2.13	0.58
1:A:138:HIS:HE1	4:A:648:HOH:O	1.93	0.52
1:C:141:ASP:OD1	1:C:173:ARG:HG2	2.12	0.49
1:D:107:ARG:NH1	1:D:128:ASP:OD2	2.43	0.49
1:B:195:ASP:OD1	1:B:306:ARG:NH2	2.45	0.49
1:A:29[B]:HIS:CE1	1:A:31:ARG:HE	2.31	0.48
1:A:193:GLY:HA3	1:A:222:LEU:HD21	1.94	0.48
1:A:53:GLU:CD	4:A:619:HOH:O	2.52	0.48
1:A:123[A]:ARG:NH1	4:A:603:HOH:O	2.07	0.47
1:A:120:ASP:OD2	4:A:605:HOH:O	2.20	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:332:PHE:HB3	4:B:781:HOH:O	2.15	0.47
1:C:323:TRP:CZ2	1:C:327:GLN:NE2	2.83	0.46
1:B:288:VAL:HG22	1:B:366:VAL:HG11	1.97	0.46
1:C:288:VAL:HG22	1:C:366:VAL:HG11	1.96	0.46
1:D:193:GLY:HA3	1:D:222:LEU:HD21	1.98	0.46
1:D:90:SER:HB2	1:D:218:ASP:HB2	1.97	0.46
1:D:29[B]:HIS:HD2	1:D:227:ASP:OD2	1.99	0.45
1:B:27:ASP:OD2	4:B:601:HOH:O	2.21	0.45
1:A:50:ALA:HB2	1:D:13:GLN:HG3	1.99	0.44
1:C:193:GLY:HA3	1:C:222:LEU:HD21	1.99	0.44
1:A:95:GLU:OE2	1:B:95:GLU:OE2	2.35	0.44
1:A:27:ASP:OD2	1:A:29[B]:HIS:CE1	2.71	0.43
1:A:29[B]:HIS:HE1	1:A:31:ARG:HE	1.66	0.43
1:D:27:ASP:OD2	1:D:29[B]:HIS:CE1	2.71	0.43
1:A:90:SER:HB3	1:A:221:LLP:H5'2	2.01	0.43
1:B:244:GLU:HB3	4:B:743:HOH:O	2.17	0.43
1:C:99:ARG:HG2	1:C:99:ARG:HH11	1.85	0.42
1:C:190:SER:HA	1:C:218:ASP:HB3	2.01	0.42
1:C:323:TRP:CD1	1:C:335:PRO:HD3	2.54	0.42
1:C:138:HIS:CE1	1:C:164:ARG:HG2	2.55	0.42
1:A:157:THR:HA	1:A:158:PRO:C	2.40	0.42
1:C:138:HIS:HE1	4:C:643:HOH:O	2.03	0.42
1:C:99:ARG:HG2	1:C:99:ARG:NH1	2.35	0.42
1:A:217:GLU:HB2	1:A:232:LEU:HB2	2.02	0.42
1:B:181:HIS:HD2	4:B:813:HOH:O	2.03	0.42
1:A:221:LLP:H4'1	1:A:221:LLP:OP4	2.20	0.41
1:B:190:SER:HA	1:B:218:ASP:HB3	2.01	0.41
1:B:360:GLU:HB3	1:B:361:PRO:HD3	2.01	0.41
1:B:318:THR:HG21	4:B:808:HOH:O	2.20	0.41
1:C:338:GLN:HB3	4:C:611:HOH:O	2.21	0.41
1:C:317:ARG:HD2	1:C:322:VAL:HG22	2.02	0.40
1:A:190:SER:HA	1:A:218:ASP:HB3	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:829:HOH:O	4:C:614:HOH:O[3_645]	2.10	0.10
4:A:829:HOH:O	4:C:637:HOH:O[3_645]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	352/376 (94%)	348 (99%)	4 (1%)	0	100	100
1	B	350/376 (93%)	344 (98%)	6 (2%)	0	100	100
1	C	350/376 (93%)	344 (98%)	6 (2%)	0	100	100
1	D	363/376 (96%)	357 (98%)	6 (2%)	0	100	100
All	All	1415/1504 (94%)	1393 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/314 (93%)	292 (100%)	0	100	100
1	B	292/314 (93%)	291 (100%)	1 (0%)	92	95
1	C	292/314 (93%)	292 (100%)	0	100	100
1	D	303/314 (96%)	302 (100%)	1 (0%)	92	95
All	All	1179/1256 (94%)	1177 (100%)	2 (0%)	93	96

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	314	PHE
1	D	164	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	HIS
1	B	340	HIS
1	C	138	HIS
1	C	181	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	LLP	D	221	1	23,24,25	2.13	3 (13%)	25,32,34	1.72	6 (24%)
1	LLP	A	221	1	23,24,25	1.98	3 (13%)	25,32,34	1.62	5 (20%)
1	LLP	B	221	1	23,24,25	2.02	3 (13%)	25,32,34	1.42	4 (16%)
1	LLP	C	221	1	23,24,25	2.18	3 (13%)	25,32,34	1.77	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	D	221	1	-	3/16/17/19	0/1/1/1
1	LLP	A	221	1	-	5/16/17/19	0/1/1/1
1	LLP	B	221	1	-	3/16/17/19	0/1/1/1
1	LLP	C	221	1	-	3/16/17/19	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	221	LLP	C3-C2	6.92	1.47	1.40
1	D	221	LLP	C3-C2	6.79	1.47	1.40
1	A	221	LLP	C3-C2	6.47	1.47	1.40
1	B	221	LLP	C3-C2	6.07	1.47	1.40
1	D	221	LLP	C4-C5	5.17	1.48	1.42
1	C	221	LLP	C4-C3	5.17	1.48	1.40
1	B	221	LLP	C4-C5	5.10	1.48	1.42
1	C	221	LLP	C4-C5	4.88	1.48	1.42
1	A	221	LLP	C4-C5	4.83	1.48	1.42
1	B	221	LLP	C4-C3	4.64	1.47	1.40
1	D	221	LLP	C4-C3	4.57	1.47	1.40
1	A	221	LLP	C4-C3	3.95	1.46	1.40

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	221	LLP	OP4-C5'-C5	4.68	118.27	109.35
1	D	221	LLP	C3-C4-C5	-4.17	115.06	118.26
1	C	221	LLP	C4-C3-C2	-4.16	117.61	120.19
1	A	221	LLP	OP4-C5'-C5	4.05	117.07	109.35
1	D	221	LLP	OP4-C5'-C5	3.71	116.42	109.35
1	A	221	LLP	C3-C4-C5	-3.58	115.51	118.26
1	B	221	LLP	C4-C3-C2	-2.94	118.37	120.19
1	D	221	LLP	CD-CE-NZ	-2.79	104.09	110.93
1	D	221	LLP	OP4-P-OP1	-2.74	98.78	106.47
1	B	221	LLP	OP4-C5'-C5	2.57	114.25	109.35
1	C	221	LLP	OP2-P-OP4	-2.56	99.92	106.73
1	D	221	LLP	C4-C3-C2	-2.51	118.63	120.19
1	D	221	LLP	C6-N1-C2	2.48	123.76	119.17
1	A	221	LLP	CD-CE-NZ	-2.38	105.09	110.93
1	C	221	LLP	CD-CE-NZ	-2.34	105.20	110.93
1	B	221	LLP	C6-N1-C2	2.30	123.42	119.17
1	B	221	LLP	OP4-P-OP1	-2.27	100.12	106.47
1	A	221	LLP	C2'-C2-C3	-2.09	118.31	120.89
1	A	221	LLP	C2'-C2-N1	2.02	121.61	117.67
1	C	221	LLP	C6-N1-C2	2.01	122.88	119.17

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	221	LLP	C4-C5-C5'-OP4
1	A	221	LLP	C6-C5-C5'-OP4
1	B	221	LLP	C4-C5-C5'-OP4
1	B	221	LLP	C6-C5-C5'-OP4
1	C	221	LLP	C4-C5-C5'-OP4
1	C	221	LLP	C6-C5-C5'-OP4
1	D	221	LLP	C4-C5-C5'-OP4
1	D	221	LLP	C6-C5-C5'-OP4
1	B	221	LLP	CG-CD-CE-NZ
1	C	221	LLP	CG-CD-CE-NZ
1	A	221	LLP	CA-CB-CG-CD
1	A	221	LLP	CG-CD-CE-NZ
1	A	221	LLP	C5'-OP4-P-OP2
1	D	221	LLP	CA-CB-CG-CD

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	221	LLP	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	350/376 (93%)	-0.28	9 (2%) 56 61	8, 23, 54, 69	0
1	B	352/376 (93%)	-0.22	7 (1%) 65 69	11, 26, 50, 72	0
1	C	352/376 (93%)	-0.19	9 (2%) 56 61	12, 27, 51, 71	0
1	D	363/376 (96%)	-0.15	21 (5%) 23 28	13, 27, 59, 115	0
All	All	1417/1504 (94%)	-0.21	46 (3%) 47 54	8, 26, 55, 115	0

All (46) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	19	LEU	5.2
1	D	372	VAL	4.8
1	D	328	ARG	4.8
1	D	317	ARG	4.5
1	D	18	ALA	4.1
1	D	323	TRP	3.9
1	A	329	HIS	3.7
1	C	317	ARG	3.6
1	C	296	GLU	3.2
1	D	11	LEU	3.2
1	C	315	ALA	3.1
1	A	328	ARG	3.1
1	D	293	ALA	3.0
1	D	330	HIS	3.0
1	D	329	HIS	2.9
1	D	14	TRP	2.9
1	C	327	GLN	2.8
1	D	21	SER	2.8
1	D	316	GLY	2.7
1	A	364	LYS	2.7
1	A	327	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	294	GLY	2.7
1	D	324	GLU	2.6
1	B	315	ALA	2.6
1	B	290	ARG	2.5
1	D	373	LEU	2.5
1	D	321	GLU	2.5
1	D	296	GLU	2.4
1	C	328	ARG	2.4
1	D	327	GLN	2.4
1	A	372	VAL	2.3
1	D	15	GLU	2.3
1	B	317	ARG	2.3
1	A	330	HIS	2.3
1	C	290	ARG	2.3
1	C	375	THR	2.3
1	D	371	THR	2.3
1	C	316	GLY	2.2
1	B	328	ARG	2.2
1	B	296	GLU	2.2
1	D	17	LEU	2.1
1	A	326	LEU	2.1
1	B	316	GLY	2.1
1	A	208	GLU	2.1
1	B	375	THR	2.1
1	A	368	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	B	221	24/25	0.97	0.22	21,28,37,37	0
1	LLP	A	221	24/25	0.98	0.15	18,23,30,31	0
1	LLP	C	221	24/25	0.98	0.21	21,28,34,35	0
1	LLP	D	221	24/25	0.98	0.13	20,26,31,32	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	B	501	1/1	0.88	0.06	35,35,35,35	0
2	CL	B	500	1/1	0.93	0.27	33,33,33,33	0
2	CL	C	500	1/1	0.96	0.19	33,33,33,33	0
2	CL	D	500	1/1	0.98	0.08	27,27,27,27	0
2	CL	A	500	1/1	0.98	0.10	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.