



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 07:11 pm GMT

PDB ID : 3DKQ  
Title : Crystal structure of Putative Oxygenase (YP\_001051978.1) from SHE-  
WANELLA BALTICA OS155 at 2.26 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2008-06-25  
Resolution : 2.26 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.26 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1178 (2.26-2.26)
Clashscore	122126	1286 (2.26-2.26)
Ramachandran outliers	120053	1253 (2.26-2.26)
Sidechain outliers	120020	1254 (2.26-2.26)
RSRZ outliers	108989	1158 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	
1	B	243	
1	C	243	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	IMD	B	502	-	-	-	X

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 5600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PKHD-type hydroxylase Sbal\_3634.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	230	Total 1867	C 1180	N 332	O 349	Se 6	0	7	0
1	B	207	Total 1640	C 1036	N 288	O 312	Se 4	0	2	0
1	C	221	Total 1778	C 1125	N 309	O 339	Se 5	0	4	0

There are 57 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	leader sequence	UNP A3D8P6
A	-17	GLY	-	leader sequence	UNP A3D8P6
A	-16	SER	-	leader sequence	UNP A3D8P6
A	-15	ASP	-	leader sequence	UNP A3D8P6
A	-14	LYS	-	leader sequence	UNP A3D8P6
A	-13	ILE	-	leader sequence	UNP A3D8P6
A	-12	HIS	-	leader sequence	UNP A3D8P6
A	-11	HIS	-	leader sequence	UNP A3D8P6
A	-10	HIS	-	leader sequence	UNP A3D8P6
A	-9	HIS	-	leader sequence	UNP A3D8P6
A	-8	HIS	-	leader sequence	UNP A3D8P6
A	-7	HIS	-	leader sequence	UNP A3D8P6
A	-6	GLU	-	leader sequence	UNP A3D8P6
A	-5	ASN	-	leader sequence	UNP A3D8P6
A	-4	LEU	-	leader sequence	UNP A3D8P6
A	-3	TYR	-	leader sequence	UNP A3D8P6
A	-2	PHE	-	leader sequence	UNP A3D8P6
A	-1	GLN	-	leader sequence	UNP A3D8P6
A	0	GLY	-	leader sequence	UNP A3D8P6
B	-18	MSE	-	leader sequence	UNP A3D8P6
B	-17	GLY	-	leader sequence	UNP A3D8P6
B	-16	SER	-	leader sequence	UNP A3D8P6
B	-15	ASP	-	leader sequence	UNP A3D8P6

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	-14	LYS	-	leader sequence	UNP A3D8P6
B	-13	ILE	-	leader sequence	UNP A3D8P6
B	-12	HIS	-	leader sequence	UNP A3D8P6
B	-11	HIS	-	leader sequence	UNP A3D8P6
B	-10	HIS	-	leader sequence	UNP A3D8P6
B	-9	HIS	-	leader sequence	UNP A3D8P6
B	-8	HIS	-	leader sequence	UNP A3D8P6
B	-7	HIS	-	leader sequence	UNP A3D8P6
B	-6	GLU	-	leader sequence	UNP A3D8P6
B	-5	ASN	-	leader sequence	UNP A3D8P6
B	-4	LEU	-	leader sequence	UNP A3D8P6
B	-3	TYR	-	leader sequence	UNP A3D8P6
B	-2	PHE	-	leader sequence	UNP A3D8P6
B	-1	GLN	-	leader sequence	UNP A3D8P6
B	0	GLY	-	leader sequence	UNP A3D8P6
C	-18	MSE	-	leader sequence	UNP A3D8P6
C	-17	GLY	-	leader sequence	UNP A3D8P6
C	-16	SER	-	leader sequence	UNP A3D8P6
C	-15	ASP	-	leader sequence	UNP A3D8P6
C	-14	LYS	-	leader sequence	UNP A3D8P6
C	-13	ILE	-	leader sequence	UNP A3D8P6
C	-12	HIS	-	leader sequence	UNP A3D8P6
C	-11	HIS	-	leader sequence	UNP A3D8P6
C	-10	HIS	-	leader sequence	UNP A3D8P6
C	-9	HIS	-	leader sequence	UNP A3D8P6
C	-8	HIS	-	leader sequence	UNP A3D8P6
C	-7	HIS	-	leader sequence	UNP A3D8P6
C	-6	GLU	-	leader sequence	UNP A3D8P6
C	-5	ASN	-	leader sequence	UNP A3D8P6
C	-4	LEU	-	leader sequence	UNP A3D8P6
C	-3	TYR	-	leader sequence	UNP A3D8P6
C	-2	PHE	-	leader sequence	UNP A3D8P6
C	-1	GLN	-	leader sequence	UNP A3D8P6
C	0	GLY	-	leader sequence	UNP A3D8P6

- Molecule 2 is NICKEL (II) ION (three-letter code: NI) (formula: Ni).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Ni 1 1	0	0
2	A	1	Total Ni 1 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total Ni 1 1	0	0

- Molecule 3 is IMIDAZOLE (three-letter code: IMD) (formula:  $C_3H_5N_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N 5 3 2	0	0
3	A	1	Total C N 5 3 2	0	0
3	B	1	Total C N 5 3 2	0	0
3	B	1	Total C N 5 3 2	0	0
3	C	1	Total C N 5 3 2	0	0
3	C	1	Total C N 5 3 2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	119	Total O 119 119	0	0
5	B	52	Total O 54 54	0	2
5	C	73	Total O 73 73	0	0





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	153.45Å 153.45Å 84.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.86 – 2.26 28.86 – 2.26	Depositor EDS
% Data completeness (in resolution range)	99.9 (28.86-2.26) 100.0 (28.86-2.26)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	0.12	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.52 (at 2.26Å)	Xtrriage
Refinement program	PHENIX, REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.196 , 0.232 0.239 , 0.268	Depositor DCC
$R_{free}$ test set	2424 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	36.6	Xtrriage
Anisotropy	0.260	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 51.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	5600	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	55.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IMD, NI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	0/1919	0.96	6/2586 (0.2%)
1	B	0.68	0/1670	0.84	4/2255 (0.2%)
1	C	0.70	0/1821	0.88	5/2458 (0.2%)
All	All	0.70	0/5410	0.89	15/7299 (0.2%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	179	ARG	NE-CZ-NH2	-10.97	114.81	120.30
1	A	184	ARG	NE-CZ-NH1	10.66	125.63	120.30
1	C	184	ARG	NE-CZ-NH1	9.78	125.19	120.30
1	C	184	ARG	NE-CZ-NH2	-9.60	115.50	120.30
1	A	179	ARG	NE-CZ-NH1	9.45	125.03	120.30
1	A	184	ARG	NE-CZ-NH2	-8.50	116.05	120.30
1	B	184	ARG	NE-CZ-NH2	-7.83	116.39	120.30
1	B	184	ARG	NE-CZ-NH1	6.85	123.73	120.30
1	B	219	ARG	NE-CZ-NH1	6.15	123.37	120.30
1	B	220	ARG	NE-CZ-NH2	-5.81	117.39	120.30
1	A	219[A]	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	A	219[B]	ARG	NE-CZ-NH1	5.62	123.11	120.30
1	C	110	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	C	220	ARG	NE-CZ-NH2	-5.55	117.53	120.30
1	C	80	TYR	N-CA-C	-5.06	97.34	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1867	0	1849	10	0
1	B	1640	0	1590	5	0
1	C	1778	0	1740	10	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	10	0	10	0	0
3	B	10	0	10	0	0
3	C	10	0	10	0	0
4	A	30	0	40	0	0
4	C	6	0	8	0	0
5	A	119	0	0	1	0
5	B	54	0	0	0	0
5	C	73	0	0	0	0
All	All	5600	0	5257	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:MSE:HE2	1:A:97:ILE:HB	1.44	0.98
1:C:-4:LEU:HD21	1:C:3:ILE:HD11	1.55	0.86
1:A:36:MSE:HE2	1:A:97:ILE:CB	2.11	0.80
1:A:75:LEU:HB3	1:A:177:MSE:HE3	1.68	0.75
1:A:32:THR:HG21	1:A:35:ALA:HB3	1.77	0.66
1:A:36:MSE:HE2	1:A:97:ILE:CG2	2.34	0.57
1:A:171[B]:PHE:HE2	1:A:173:TRP:CZ2	2.24	0.55
1:A:186:LEU:HD21	1:B:213:VAL:HG13	1.89	0.54
1:C:2:LEU:HD11	1:C:139:GLN:HE22	1.74	0.53
1:A:32:THR:CG2	1:A:35:ALA:HB3	2.38	0.52
1:C:14:VAL:HG11	1:C:145:ALA:HB1	1.91	0.52
1:C:177:MSE:CE	1:C:218:LEU:HD11	2.41	0.50
1:C:-4:LEU:HD12	1:C:69:GLN:CD	2.32	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:-4:LEU:HD21	1:C:3:ILE:CD1	2.37	0.49
1:B:75:LEU:HB3	1:B:177:MSE:HE3	1.94	0.48
1:A:100:ALA:C	1:A:101:ILE:HD12	2.36	0.47
1:B:14:VAL:HG11	1:B:145:ALA:HB1	1.97	0.45
1:C:-4:LEU:HD22	1:C:-2:PHE:CE1	2.51	0.45
1:C:3:ILE:HG23	1:C:150:LEU:HB3	1.98	0.45
1:C:132:ILE:HG23	1:C:155:SER:HB3	2.02	0.42
1:B:186:LEU:HD23	1:B:186:LEU:C	2.40	0.41
1:B:203:ALA:HB3	1:B:206:GLU:HG2	2.03	0.41
1:C:177:MSE:HE2	1:C:218:LEU:HD11	2.03	0.41
1:A:177:MSE:HE2	5:A:588:HOH:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	235/243 (97%)	234 (100%)	1 (0%)	0	100	100
1	B	203/243 (84%)	200 (98%)	3 (2%)	0	100	100
1	C	221/243 (91%)	216 (98%)	5 (2%)	0	100	100
All	All	659/729 (90%)	650 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/204 (99%)	199 (98%)	3 (2%)	67	77
1	B	174/204 (85%)	172 (99%)	2 (1%)	76	84
1	C	193/204 (95%)	187 (97%)	6 (3%)	43	52
All	All	569/612 (93%)	558 (98%)	11 (2%)	62	69

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	93	PHE
1	A	148	LEU
1	B	83	LEU
1	B	179	ARG
1	C	-4	LEU
1	C	-2	PHE
1	C	19[A]	GLU
1	C	19[B]	GLU
1	C	28	ASP
1	C	93	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	ASN
1	A	139	GLN
1	B	69	GLN
1	B	124	ASN
1	B	189	GLN
1	C	124	ASN
1	C	139	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 3 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	IMD	A	501	2	3,5,5	0.39	0	4,5,5	0.86	0
3	IMD	A	502	-	3,5,5	0.42	0	4,5,5	0.78	0
4	GOL	A	503	-	5,5,5	0.47	0	5,5,5	0.76	0
4	GOL	A	504	-	5,5,5	0.33	0	5,5,5	0.29	0
4	GOL	A	505	-	5,5,5	0.39	0	5,5,5	0.65	0
4	GOL	A	506	-	5,5,5	0.49	0	5,5,5	0.73	0
4	GOL	A	507	-	5,5,5	0.42	0	5,5,5	0.60	0
3	IMD	B	501	-	3,5,5	0.44	0	4,5,5	0.71	0
3	IMD	B	502	-	3,5,5	0.38	0	4,5,5	0.77	0
3	IMD	C	501	-	3,5,5	0.27	0	4,5,5	0.73	0
3	IMD	C	502	-	3,5,5	0.25	0	4,5,5	0.81	0
4	GOL	C	503	-	5,5,5	0.47	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IMD	A	501	2	-	0/0/0/0	0/1/1/1
3	IMD	A	502	-	-	0/0/0/0	0/1/1/1
4	GOL	A	503	-	-	0/4/4/4	0/0/0/0
4	GOL	A	504	-	-	0/4/4/4	0/0/0/0
4	GOL	A	505	-	-	0/4/4/4	0/0/0/0
4	GOL	A	506	-	-	0/4/4/4	0/0/0/0
4	GOL	A	507	-	-	0/4/4/4	0/0/0/0
3	IMD	B	501	-	-	0/0/0/0	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IMD	B	502	-	-	0/0/0/0	0/1/1/1
3	IMD	C	501	-	-	0/0/0/0	0/1/1/1
3	IMD	C	502	-	-	0/0/0/0	0/1/1/1
4	GOL	C	503	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	224/243 (92%)	0.76	33 (14%) 2 2	45, 52, 83, 99	0
1	B	203/243 (83%)	0.99	39 (19%) 1 1	42, 52, 74, 109	0
1	C	216/243 (88%)	0.85	31 (14%) 2 2	44, 52, 78, 104	0
All	All	643/729 (88%)	0.87	103 (16%) 1 1	42, 52, 79, 109	0

All (103) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	-4	LEU	8.2
1	B	95	TYR	7.5
1	A	171[A]	PHE	6.7
1	B	136	TYR	6.5
1	C	30	ASN	6.5
1	B	97	ILE	6.1
1	A	38	THR	6.0
1	B	90	GLY	5.8
1	C	135	THR	5.7
1	C	134	ASP	5.5
1	B	135	THR	5.3
1	C	-3	TYR	5.3
1	B	42	ARG	4.9
1	C	-2	PHE	4.8
1	C	-4	LEU	4.8
1	B	96	HIS	4.7
1	B	-5	ASN	4.5
1	B	49	ASP	4.5
1	C	29	GLY	4.2
1	C	123	GLU	4.1
1	B	28	ASP	4.1
1	A	30	ASN	4.0
1	A	116	THR	4.0

*Continued on next page...*



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	27	ILE	3.9
1	C	31	GLN	3.9
1	C	163	LEU	3.9
1	C	165	GLY	3.8
1	C	12	GLN	3.8
1	C	162	VAL	3.7
1	B	47	ASP	3.7
1	B	94	GLY	3.6
1	A	31	GLN	3.6
1	C	40	ARG	3.4
1	A	32	THR	3.4
1	A	201	THR	3.4
1	B	201	THR	3.4
1	B	48	LYS	3.4
1	C	41	LYS	3.3
1	C	-1	GLN	3.3
1	A	33	SER	3.3
1	B	156	LEU	3.3
1	A	115	ALA	3.3
1	C	136	TYR	3.3
1	A	113	LEU	3.2
1	A	37	ALA	3.1
1	C	137	GLY	3.1
1	B	92	THR	3.1
1	B	98	ASP	3.1
1	B	29	GLY	3.1
1	C	90	GLY	3.0
1	A	151	TYR	3.0
1	A	150	LEU	2.9
1	B	116	THR	2.9
1	A	-4	LEU	2.9
1	A	130	LEU	2.9
1	B	23	ALA	2.9
1	B	-3	TYR	2.9
1	B	89	GLY	2.8
1	A	224	LEU	2.8
1	B	150	LEU	2.8
1	C	127	GLY	2.8
1	A	29	GLY	2.8
1	A	-1	GLN	2.8
1	B	44	GLN	2.8
1	B	171	PHE	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	169	ALA	2.7
1	B	115	ALA	2.7
1	B	110	ARG	2.7
1	A	114	SER	2.6
1	B	24	ARG	2.6
1	B	22	ASP	2.5
1	A	148	LEU	2.5
1	C	28	ASP	2.5
1	A	117	LEU	2.5
1	B	16	HIS	2.5
1	B	-2	PHE	2.4
1	A	170	ALA	2.4
1	A	105	PRO	2.3
1	B	46	LEU	2.3
1	B	52	VAL	2.3
1	C	140	SER	2.3
1	A	35	ALA	2.2
1	B	149	VAL	2.2
1	B	155	SER	2.2
1	A	173	TRP	2.2
1	C	89	GLY	2.2
1	A	118	PHE	2.1
1	C	116	THR	2.1
1	B	51	PRO	2.1
1	A	-2	PHE	2.1
1	C	164	SER	2.1
1	C	42	ARG	2.1
1	C	19[A]	GLU	2.1
1	B	199	ALA	2.1
1	A	149	VAL	2.1
1	A	136	TYR	2.1
1	C	138	GLN	2.1
1	C	113	LEU	2.0
1	A	159	VAL	2.0
1	C	151	TYR	2.0
1	A	28	ASP	2.0
1	A	205	GLN	2.0
1	C	10	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	IMD	B	502	5/5	0.72	0.45	72,76,77,78	0
4	GOL	A	503	6/6	0.76	0.24	79,81,84,84	0
4	GOL	A	506	6/6	0.78	0.21	79,83,85,88	0
2	NI	B	500	1/1	0.80	0.18	59,59,59,59	1
4	GOL	A	504	6/6	0.84	0.23	70,75,78,82	0
4	GOL	A	507	6/6	0.85	0.23	76,84,85,87	0
4	GOL	A	505	6/6	0.86	0.25	74,80,82,83	0
4	GOL	C	503	6/6	0.89	0.27	92,96,97,99	0
3	IMD	B	501	5/5	0.92	0.31	87,88,90,90	0
3	IMD	C	502	5/5	0.92	0.23	62,63,65,68	0
3	IMD	C	501	5/5	0.95	0.16	48,49,53,53	0
2	NI	C	500	1/1	0.96	0.15	53,53,53,53	0
3	IMD	A	502	5/5	0.97	0.24	46,48,51,53	0
3	IMD	A	501	5/5	0.97	0.17	36,40,45,46	0
2	NI	A	500	1/1	0.99	0.15	42,42,42,42	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.