



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 28, 2024 – 12:32 AM EST

PDB ID : 1DLL  
Title : The HC fragment of tetanus toxin complexed with lactose  
Authors : Emsley, P.; Fotinou, C.; Black, I.; Fairweather, N.F.; Charles, I.G.; Watts, C.; Hewitt, E.; Isaacs, N.W.  
Deposited on : 1999-12-10  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	
2	B	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	3001[B]	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4302 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TETANUS TOXIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	441	3563	2282	597	675	9	0	0	0

- Molecule 2 is an oligosaccharide called beta-D-galactopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	B	2	23	12	11	0	2	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	1
3	A	1	Total C O 6 3 3	0	1
3	A	1	Total C O 6 3 3	0	1
3	A	1	Total C O 6 3 3	0	1


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	692	Total O 692 692	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TETANUS TOXIN



- Molecule 2: beta-D-galactopyranose-(1-4)-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.08Å 70.88Å 122.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.80 27.88 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.4 (30.00-1.80) 99.2 (27.88-1.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.40 (at 1.80Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.201 , 0.267 0.204 , 0.252	Depositor DCC
$R_{free}$ test set	2757 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.8	Xtrriage
Anisotropy	0.574	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 54.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4302	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, GOL, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.46	0/3644	1.12	15/4945 (0.3%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1123	ARG	NE-CZ-NH1	-7.36	116.62	120.30
1	A	957	ARG	NE-CZ-NH2	-7.09	116.76	120.30
1	A	1167	ARG	NE-CZ-NH1	6.82	123.71	120.30
1	A	957	ARG	NE-CZ-NH1	6.67	123.63	120.30
1	A	1021	ASP	CB-CG-OD1	6.67	124.30	118.30
1	A	1040	ARG	NE-CZ-NH1	6.23	123.41	120.30
1	A	1053	MET	CG-SD-CE	5.63	109.21	100.20
1	A	1229	TYR	CB-CG-CD2	-5.31	117.81	121.00
1	A	1274	GLN	CA-CB-CG	5.28	125.01	113.40
1	A	1124	TYR	CB-CG-CD1	-5.16	117.90	121.00
1	A	1123	ARG	CD-NE-CZ	5.14	130.80	123.60
1	A	1126	THR	N-CA-CB	5.10	120.00	110.30
1	A	1145	ILE	C-N-CA	5.10	134.45	121.70
1	A	1284	LEU	CB-CG-CD2	5.10	119.67	111.00
1	A	1090	ARG	NE-CZ-NH2	-5.08	117.76	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3563	0	3516	85	0
2	B	23	0	17	0	0
3	A	24	0	25	11	0
4	A	692	0	0	13	0
All	All	4302	0	3558	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1003:THR:HB	1:A:1013:GLN:HG2	1.51	0.92
1:A:1141:GLN:HE21	1:A:1142:LEU:H	1.18	0.90
1:A:932:GLU:HG3	1:A:1073:LYS:HB2	1.57	0.86
1:A:1226:ARG:HH12	3:A:3001[B]:GOL:H31	1.43	0.82
1:A:1115:ARG:CG	1:A:1115:ARG:HH11	2.00	0.75
1:A:1223:ARG:HG2	1:A:1223:ARG:HH11	1.53	0.74
1:A:972:ASN:HD21	1:A:1078:ASN:H	1.34	0.73
1:A:1111:ILE:HA	4:A:637:HOH:O	1.89	0.72
1:A:976:ILE:HG22	1:A:1074:LEU:HD23	1.72	0.71
1:A:1216:ASN:H	3:A:3001[B]:GOL:H32	1.61	0.66
1:A:972:ASN:ND2	1:A:1078:ASN:H	1.94	0.65
1:A:918:ILE:H	1:A:918:ILE:HD12	1.62	0.64
1:A:1097:ASN:ND2	1:A:1100:GLU:H	1.96	0.64
1:A:918:ILE:HG23	4:A:678:HOH:O	1.96	0.63
1:A:1115:ARG:CG	1:A:1115:ARG:NH1	2.60	0.62
1:A:880:ILE:HG13	4:A:683:HOH:O	1.99	0.62
1:A:1003:THR:HG21	1:A:1013:GLN:HE21	1.66	0.60
1:A:918:ILE:HD11	1:A:1031:TRP:CE3	2.37	0.59
1:A:918:ILE:HD12	1:A:918:ILE:N	2.17	0.59
1:A:949:ASN:ND2	1:A:1040:ARG:H	2.01	0.59
1:A:929:GLU:HG3	4:A:594:HOH:O	2.04	0.57
1:A:1126:THR:HG21	4:A:66:HOH:O	2.03	0.57
1:A:1141:GLN:NE2	1:A:1142:LEU:H	1.98	0.56
1:A:1183:ASN:HD22	3:A:3003[B]:GOL:H11	1.70	0.56
1:A:1221:LEU:O	1:A:1223:ARG:NH1	2.39	0.55
1:A:1115:ARG:HH11	1:A:1115:ARG:HG2	1.73	0.54
1:A:1184:ASN:HB3	4:A:483:HOH:O	2.08	0.53

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1202:TYR:O	1:A:1203:ASN:HB2	2.09	0.53
1:A:878:ASP:OD1	1:A:1103:LYS:HD3	2.09	0.52
1:A:1115:ARG:NH1	1:A:1115:ARG:HG3	2.24	0.52
1:A:1223:ARG:NH1	1:A:1223:ARG:CG	2.69	0.52
1:A:1220:ASN:O	1:A:1221:LEU:HB2	2.09	0.52
1:A:915:VAL:HB	1:A:916:PRO:HD2	1.92	0.51
1:A:981:LYS:O	1:A:982:LYS:HB2	2.10	0.51
1:A:1170:TYR:CE1	1:A:1308:THR:HA	2.45	0.51
1:A:939:MET:HG2	1:A:942:GLU:OE2	2.11	0.51
1:A:1183:ASN:HB3	3:A:3003[B]:GOL:H12	1.93	0.51
1:A:1003:THR:CG2	1:A:1013:GLN:HE21	2.23	0.51
1:A:1223:ARG:HG2	1:A:1223:ARG:NH1	2.25	0.51
1:A:1130:LEU:HD22	1:A:1303:TRP:HB3	1.92	0.50
1:A:947:PHE:O	1:A:1040:ARG:HG3	2.12	0.50
1:A:1097:ASN:HD21	1:A:1100:GLU:HG3	1.77	0.50
1:A:1178:LYS:HE2	4:A:114:HOH:O	2.10	0.50
1:A:915:VAL:HB	1:A:916:PRO:CD	2.42	0.49
1:A:1226:ARG:NH1	3:A:3001[B]:GOL:H31	2.20	0.49
1:A:876:ASP:OD2	1:A:879:VAL:HG23	2.12	0.49
1:A:1149:MET:HB2	1:A:1284:LEU:HG	1.95	0.49
1:A:936:HIS:HD2	1:A:1069:ASN:OD1	1.95	0.48
1:A:1003:THR:HG23	4:A:371:HOH:O	2.14	0.48
1:A:1216:ASN:N	3:A:3001[B]:GOL:H32	2.28	0.47
1:A:1094:LYS:HB2	3:A:3002[B]:GOL:H31	1.96	0.47
1:A:1231:ALA:HB3	1:A:1234:ILE:HD12	1.97	0.47
1:A:947:PHE:CZ	1:A:1063:ALA:HB2	2.50	0.46
1:A:1216:ASN:H	3:A:3001[B]:GOL:C3	2.28	0.46
1:A:982:LYS:HE3	1:A:1076:ARG:HH21	1.81	0.46
1:A:1183:ASN:OD1	1:A:1183:ASN:N	2.49	0.46
1:A:1223:ARG:HH11	1:A:1223:ARG:CG	2.13	0.46
1:A:1093:CYS:HB2	4:A:139:HOH:O	2.15	0.46
1:A:976:ILE:HG22	1:A:1074:LEU:CD2	2.43	0.45
1:A:1097:ASN:HD22	1:A:1097:ASN:C	2.21	0.45
1:A:932:GLU:OE2	1:A:934:ILE:HD11	2.17	0.44
1:A:1099:LYS:HE2	4:A:80:HOH:O	2.16	0.44
1:A:918:ILE:HG12	1:A:955:TRP:CD2	2.53	0.44
1:A:1194:ASP:HB3	3:A:3003[B]:GOL:H31	1.99	0.44
1:A:1094:LYS:HB2	3:A:3002[B]:GOL:C3	2.48	0.44
1:A:1219:ASN:HB3	4:A:644:HOH:O	2.17	0.44
1:A:1226:ARG:HH12	3:A:3001[B]:GOL:C3	2.22	0.44
1:A:932:GLU:HG3	1:A:1073:LYS:CB	2.37	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1115:ARG:HD3	4:A:108:HOH:O	2.18	0.43
1:A:974:TYR:CE2	1:A:1074:LEU:HD22	2.54	0.43
1:A:1181:THR:HG22	4:A:234:HOH:O	2.19	0.43
1:A:902:PHE:O	1:A:903:ASN:HB2	2.18	0.42
1:A:985:LEU:HD12	1:A:985:LEU:O	2.19	0.42
1:A:1104:LEU:HD22	1:A:1108:TYR:HE2	1.83	0.42
1:A:1045:ASN:HB3	1:A:1052:LEU:CD1	2.50	0.42
1:A:891:ILE:HG12	1:A:896:ILE:HG12	2.01	0.42
1:A:1213:LYS:O	1:A:1214:ASP:HB2	2.20	0.42
1:A:1097:ASN:HD22	1:A:1100:GLU:H	1.67	0.41
1:A:1234:ILE:HA	1:A:1235:PRO:HD3	1.93	0.41
1:A:1097:ASN:HB2	1:A:1098:PRO:HD2	2.03	0.41
1:A:981:LYS:N	1:A:1067:ASP:HB2	2.36	0.41
1:A:1042:SER:OG	1:A:1043:SER:N	2.54	0.41
1:A:1111:ILE:H	1:A:1111:ILE:HG12	1.63	0.40
1:A:918:ILE:H	1:A:918:ILE:CD1	2.32	0.40
1:A:1067:ASP:OD1	1:A:1068:ASN:N	2.55	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	439/441 (100%)	414 (94%)	23 (5%)	2 (0%)	29 15

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	982	LYS
1	A	1219	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	397/397 (100%)	373 (94%)	24 (6%)	19 7

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	929	GLU
1	A	939	MET
1	A	987	ILE
1	A	1003	THR
1	A	1035	THR
1	A	1097	ASN
1	A	1099	LYS
1	A	1103	LYS
1	A	1104	LEU
1	A	1111	ILE
1	A	1115	ARG
1	A	1122	LEU
1	A	1126	THR
1	A	1167	ARG
1	A	1183	ASN
1	A	1184	ASN
1	A	1192	SER
1	A	1204	ASN
1	A	1216	ASN
1	A	1238	LYS
1	A	1245	LEU
1	A	1261	LYS
1	A	1284	LEU
1	A	1297	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	924	HIS

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	A	936	HIS
1	A	944	ASN
1	A	949	ASN
1	A	972	ASN
1	A	998	ASN
1	A	1013	GLN
1	A	1068	ASN
1	A	1097	ASN
1	A	1141	GLN
1	A	1171	ASN
1	A	1184	ASN
1	A	1216	ASN
1	A	1277	ASN
1	A	1293	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BGC	B	1[A]	2	12,12,12	0.54	0	17,17,17	0.86	0
2	GAL	B	2[A]	2	11,11,12	0.39	0	15,15,17	1.06	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	1[A]	2	-	2/2/22/22	0/1/1/1
2	GAL	B	2[A]	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2[A]	GAL	O2-C2-C1	2.24	113.74	109.15
2	B	2[A]	GAL	C2-C3-C4	-2.11	107.24	110.89

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1[A]	BGC	O5-C5-C6-O6
2	B	1[A]	BGC	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	GOL	A	3002[B]	-	5,5,5	0.59	0	5,5,5	0.55	0
3	GOL	A	3001[B]	-	5,5,5	0.22	0	5,5,5	0.16	0
3	GOL	A	3000[B]	-	5,5,5	0.23	0	5,5,5	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	3002[B]	-	-	2/4/4/4	-
3	GOL	A	3001[B]	-	-	4/4/4/4	-
3	GOL	A	3000[B]	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	3000[B]	GOL	C1-C2-C3-O3
3	A	3001[B]	GOL	O1-C1-C2-C3
3	A	3001[B]	GOL	C1-C2-C3-O3
3	A	3000[B]	GOL	O2-C2-C3-O3
3	A	3001[B]	GOL	O1-C1-C2-O2
3	A	3001[B]	GOL	O2-C2-C3-O3
3	A	3002[B]	GOL	O2-C2-C3-O3
3	A	3002[B]	GOL	C1-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	3002[B]	GOL	2	0
3	A	3001[B]	GOL	6	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	441/441 (100%)	0.40	40 (9%) <b>9</b> <b>7</b>	20, 30, 55, 82	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	983	HIS	17.0
1	A	985	LEU	10.2
1	A	984	SER	7.5
1	A	1183	ASN	6.3
1	A	875	GLU	6.1
1	A	982	LYS	6.0
1	A	939	MET	5.8
1	A	986	SER	5.4
1	A	1112	THR	5.4
1	A	1184	ASN	4.9
1	A	1065	ARG	4.7
1	A	942	GLU	4.7
1	A	943	TYR	4.4
1	A	1111	ILE	3.6
1	A	1185	GLU	3.5
1	A	902	PHE	3.5
1	A	1221	LEU	3.4
1	A	987	ILE	3.3
1	A	940	ASP	3.3
1	A	1182	PRO	3.1
1	A	1113	PHE	3.0
1	A	1233	GLY	2.9
1	A	938	ALA	2.9
1	A	1234	ILE	2.7
1	A	941	ILE	2.6
1	A	1181	THR	2.6
1	A	883	LYS	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	936	HIS	2.5
1	A	1036	ILE	2.5
1	A	954	PHE	2.5
1	A	1099	LYS	2.4
1	A	952	VAL	2.4
1	A	879	VAL	2.3
1	A	1255	LEU	2.1
1	A	981	LYS	2.1
1	A	1089	PHE	2.1
1	A	1232	PRO	2.0
1	A	1034	ILE	2.0
1	A	1204	ASN	2.0
1	A	1110	SER	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	BGC	B	1[A]	12/12	0.71	0.20	34,36,39,40	12
2	GAL	B	2[A]	11/12	0.78	0.18	33,34,35,35	11

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	A	3002[B]	6/6	0.60	0.33	68,69,70,70	0
3	GOL	A	3001[B]	6/6	0.67	0.23	66,67,68,68	0
3	GOL	A	3003[B]	6/6	0.73	0.19	42,45,46,47	0
3	GOL	A	3000[B]	6/6	0.87	0.18	86,86,86,86	6

## 6.5 Other polymers [i](#)

There are no such residues in this entry.