



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2020 – 05:21 AM BST

PDB ID : 3DNI
Title : CRYSTALLOGRAPHIC REFINEMENT AND STRUCTURE OF DNASE I
AT 2 ANGSTROMS RESOLUTION
Authors : Oefner, C.; Suck, D.
Deposited on : 1992-08-20
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	260	
2	B	7	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	B	2	X	-	-	-
2	MAN	B	6	X	-	-	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 2494 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DEOXYRIBONUCLEASE I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	258	2034	1290	339	397	8	0	0	0

- Molecule 2 is an oligosaccharide called alpha-D-galactopyranose-(1-6)-beta-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	7	83	46	2	35	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	375	Total	O	0	0
			375	375		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: DEOXYRIBONUCLEASE I



- Molecule 2: alpha-D-galactopyranose-(1-6)-beta-D-mannopyranose-(1-3)-[beta-D-mannopyranose-(1-6)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	131.60Å 54.90Å 38.40Å 90.00° 91.40° 90.00°	Depositor
Resolution (Å)	6.00 – 2.00	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.177 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2494	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GLA, CA, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.30	5/2079 (0.2%)	2.09	70/2830 (2.5%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	143	GLU	CD-OE2	-7.83	1.17	1.25
1	A	178	SER	CB-OG	7.02	1.51	1.42
1	A	17	SER	CB-OG	5.24	1.49	1.42
1	A	209	CYS	CB-SG	-5.22	1.73	1.81
1	A	143	GLU	CG-CD	5.01	1.59	1.51

All (70) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	143	GLU	OE1-CD-OE2	20.41	147.80	123.30
1	A	9	ARG	NE-CZ-NH1	14.85	127.72	120.30
1	A	73	ARG	NE-CZ-NH1	-14.38	113.11	120.30
1	A	79	ARG	NE-CZ-NH1	14.17	127.38	120.30
1	A	121	HIS	CA-CB-CG	12.17	134.29	113.60
1	A	127	GLU	CA-CB-CG	11.73	139.20	113.40
1	A	187	ARG	NE-CZ-NH1	11.19	125.89	120.30
1	A	111	ARG	NE-CZ-NH1	11.06	125.83	120.30
1	A	9	ARG	CD-NE-CZ	10.78	138.69	123.60
1	A	33	ASP	CB-CG-OD2	10.71	127.94	118.30
1	A	27	ARG	CD-NE-CZ	-10.41	109.03	123.60
1	A	143	GLU	CG-CD-OE1	-10.32	97.65	118.30
1	A	162	ASP	CB-CG-OD1	-10.23	109.10	118.30
1	A	166	MET	CA-CB-CG	9.90	130.12	113.30
1	A	33	ASP	CB-CG-OD1	-9.50	109.75	118.30
1	A	27	ARG	NE-CZ-NH2	9.42	125.01	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	166	MET	N-CA-CB	-8.97	94.46	110.60
1	A	111	ARG	NE-CZ-NH2	-8.95	115.83	120.30
1	A	63	TYR	CB-CG-CD1	8.80	126.28	121.00
1	A	187	ARG	NE-CZ-NH2	-8.76	115.92	120.30
1	A	13	GLU	OE1-CD-OE2	8.66	133.70	123.30
1	A	30	ARG	NE-CZ-NH1	8.61	124.61	120.30
1	A	41	ARG	NE-CZ-NH2	-8.55	116.02	120.30
1	A	41	ARG	NE-CZ-NH1	8.18	124.39	120.30
1	A	234	ASP	CB-CG-OD1	8.16	125.64	118.30
1	A	251	ASP	CB-CG-OD2	-7.93	111.16	118.30
1	A	169	PHE	CB-CG-CD2	-7.91	115.26	120.80
1	A	164	MET	CG-SD-CE	7.24	111.79	100.20
1	A	185	ARG	NE-CZ-NH1	7.10	123.85	120.30
1	A	212	ASP	CB-CG-OD2	-7.03	111.97	118.30
1	A	166	MET	CG-SD-CE	6.97	111.35	100.20
1	A	9	ARG	NE-CZ-NH2	-6.96	116.82	120.30
1	A	77	LYS	CB-CA-C	-6.88	96.63	110.40
1	A	73	ARG	NH1-CZ-NH2	6.77	126.85	119.40
1	A	89	VAL	N-CA-CB	-6.75	96.65	111.50
1	A	99	ASP	CB-CG-OD1	6.74	124.37	118.30
1	A	256	GLU	O-C-N	6.59	133.25	122.70
1	A	63	TYR	CB-CG-CD2	-6.56	117.06	121.00
1	A	167	GLY	O-C-N	6.51	133.12	122.70
1	A	153	ASP	CB-CG-OD2	-6.42	112.52	118.30
1	A	32	TYR	CB-CG-CD2	6.35	124.81	121.00
1	A	99	ASP	CB-CG-OD2	-6.32	112.61	118.30
1	A	251	ASP	CB-CG-OD1	6.29	123.96	118.30
1	A	178	SER	CA-CB-OG	-6.26	94.29	111.20
1	A	125	VAL	CG1-CB-CG2	6.19	120.80	110.90
1	A	169	PHE	CB-CG-CD1	6.00	125.00	120.80
1	A	74	ASN	CB-CA-C	5.98	122.35	110.40
1	A	148	TYR	CB-CG-CD1	-5.94	117.44	121.00
1	A	32	TYR	CB-CG-CD1	-5.84	117.50	121.00
1	A	77	LYS	CA-CB-CG	5.79	126.14	113.40
1	A	125	VAL	CB-CA-C	-5.74	100.49	111.40
1	A	85	ARG	NE-CZ-NH2	5.71	123.16	120.30
1	A	191	THR	CA-CB-CG2	5.59	120.23	112.40
1	A	244	GLU	CA-CB-CG	5.53	125.56	113.40
1	A	17	SER	CA-CB-OG	-5.43	96.54	111.20
1	A	73	ARG	N-CA-CB	-5.33	101.01	110.60
1	A	178	SER	O-C-N	5.28	131.15	122.70
1	A	1	LEU	CB-CA-C	5.27	120.22	110.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	5	ALA	N-CA-CB	5.26	117.47	110.10
1	A	76	TYR	N-CA-CB	-5.16	101.31	110.60
1	A	198	ASP	CB-CG-OD2	-5.16	113.65	118.30
1	A	57	GLN	CA-CB-CG	5.13	124.68	113.40
1	A	93	ASP	O-C-N	5.11	130.87	122.70
1	A	41	ARG	CD-NE-CZ	-5.08	116.48	123.60
1	A	177	THR	CA-CB-CG2	5.07	119.50	112.40
1	A	138	SER	CB-CA-C	-5.05	100.51	110.10
1	A	30	ARG	NE-CZ-NH2	-5.05	117.78	120.30
1	A	56	ASN	O-C-N	5.04	130.77	122.70
1	A	74	ASN	CA-CB-CG	-5.03	102.34	113.40
1	A	139	ASP	CA-CB-CG	-5.00	102.39	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2034	0	1969	17	0
2	B	83	0	64	5	0
3	A	2	0	0	0	0
4	A	375	0	0	7	2
All	All	2494	0	2033	22	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:2:NAG:H4	2:B:3:BMA:O2	1.34	1.08
2:B:2:NAG:C4	2:B:3:BMA:O2	2.15	0.94
4:A:618:HOH:O	2:B:7:BMA:H2	1.72	0.89
1:A:5:ALA:HB2	1:A:166:MET:HG2	1.69	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:3:BMA:O4	2:B:4:BMA:C1	2.38	0.72
2:B:4:BMA:H2	2:B:7:BMA:H62	1.71	0.70
1:A:30:ARG:HD2	4:A:303:HOH:O	1.99	0.62
1:A:9:ARG:NH2	4:A:609:HOH:O	2.33	0.57
1:A:174:SER:HB2	4:A:434:HOH:O	2.08	0.53
1:A:109:PHE:CD1	1:A:143:GLU:HG3	2.46	0.51
1:A:10:THR:HG23	1:A:41:ARG:HH11	1.76	0.51
1:A:92:LEU:HB2	1:A:117:LYS:HD2	1.95	0.48
1:A:243:ASN:O	1:A:247:LEU:HB2	2.14	0.47
1:A:42:ASP:OD2	1:A:45:LEU:HA	2.14	0.47
1:A:155:GLN:NE2	4:A:444:HOH:O	2.48	0.46
1:A:126:LYS:NZ	4:A:606:HOH:O	2.49	0.45
1:A:112:GLU:HA	1:A:113:PRO:HD3	1.84	0.43
1:A:22:ALA:O	1:A:26:VAL:HG23	2.18	0.43
1:A:200:ALA:O	1:A:210:ALA:HB2	2.20	0.41
1:A:50:LYS:NZ	4:A:578:HOH:O	2.53	0.41
1:A:52:LEU:HD11	1:A:65:TYR:CE1	2.56	0.41
1:A:27:ARG:HD3	1:A:27:ARG:HH11	1.48	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:618:HOH:O	4:A:618:HOH:O 2_556]	1.82	0.38
4:A:368:HOH:O	4:A:368:HOH:O 2_456]	2.10	0.10

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	254/260 (98%)	247 (97%)	6 (2%)	1 (0%)	34 30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	74	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	227/229 (99%)	210 (92%)	17 (8%)	13 9

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	THR
1	A	21	LEU
1	A	23	SER
1	A	27	ARG
1	A	45	LEU
1	A	101	CYS
1	A	107	ASP
1	A	125	VAL
1	A	138	SER
1	A	139	ASP
1	A	143	GLU
1	A	219	SER
1	A	220	LEU
1	A	221	LEU
1	A	242	SER
1	A	244	GLU
1	A	247	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	155	GLN
1	A	159	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	161	ASN
1	A	236	GLN
1	A	243	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

7 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	1.66	2 (14%)	17,19,21	5.29	12 (70%)
2	NAG	B	2	2	14,14,15	2.35	6 (42%)	17,19,21	5.39	9 (52%)
2	BMA	B	3	2	11,11,12	1.26	1 (9%)	15,15,17	3.10	6 (40%)
2	BMA	B	4	2	11,11,12	1.36	2 (18%)	15,15,17	3.78	6 (40%)
2	GLA	B	5	2	11,11,12	2.46	2 (18%)	15,15,17	4.27	7 (46%)
2	MAN	B	6	2	11,11,12	0.84	0	15,15,17	3.44	9 (60%)
2	BMA	B	7	2	11,11,12	2.78	3 (27%)	15,15,17	4.32	9 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	4/6/23/26	0/1/1/1
2	NAG	B	2	2	1/1/5/7	3/6/23/26	0/1/1/1
2	BMA	B	3	2	-	2/2/19/22	0/1/1/1
2	BMA	B	4	2	-	2/2/19/22	0/1/1/1
2	GLA	B	5	2	-	0/2/19/22	0/1/1/1
2	MAN	B	6	2	1/1/4/5	2/2/19/22	0/1/1/1
2	BMA	B	7	2	-	2/2/19/22	0/1/1/1

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	7	BMA	O2-C2	-7.98	1.26	1.43
2	B	5	GLA	C4-C5	-6.00	1.40	1.53
2	B	5	GLA	C2-C3	-4.78	1.45	1.52
2	B	2	NAG	O7-C7	4.39	1.33	1.23
2	B	2	NAG	C1-C2	4.23	1.58	1.52
2	B	1	NAG	O5-C1	-4.07	1.37	1.43
2	B	2	NAG	C8-C7	3.93	1.58	1.50
2	B	3	BMA	O5-C1	-3.23	1.38	1.43
2	B	4	BMA	C2-C3	-3.21	1.47	1.52
2	B	2	NAG	C4-C3	2.91	1.59	1.52
2	B	1	NAG	O3-C3	2.80	1.49	1.43
2	B	4	BMA	O5-C5	-2.55	1.38	1.43
2	B	7	BMA	C2-C3	2.51	1.56	1.52
2	B	7	BMA	O5-C5	-2.30	1.38	1.43
2	B	2	NAG	O3-C3	-2.23	1.37	1.43
2	B	2	NAG	O5-C5	-2.18	1.39	1.43

All (58) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C1-O5-C5	14.02	131.19	112.19
2	B	1	NAG	O7-C7-C8	-12.47	98.90	122.06
2	B	7	BMA	C1-O5-C5	11.77	128.14	112.19
2	B	1	NAG	C1-O5-C5	11.55	127.84	112.19
2	B	2	NAG	O3-C3-C2	10.20	130.58	109.47
2	B	5	GLA	C3-C4-C5	10.05	128.17	110.24
2	B	5	GLA	C1-O5-C5	9.29	124.78	112.19
2	B	3	BMA	C1-O5-C5	8.96	124.33	112.19
2	B	4	BMA	C1-C2-C3	8.84	120.53	109.67
2	B	1	NAG	C2-N2-C7	8.40	134.87	122.90
2	B	4	BMA	C1-O5-C5	7.63	122.53	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C8-C7-N2	-6.85	104.50	116.10
2	B	6	MAN	O3-C3-C2	6.79	122.99	109.99
2	B	2	NAG	O7-C7-C8	6.53	134.18	122.06
2	B	6	MAN	C1-C2-C3	6.36	117.49	109.67
2	B	7	BMA	O4-C4-C3	5.92	124.04	110.35
2	B	4	BMA	O3-C3-C2	5.88	121.25	109.99
2	B	1	NAG	C4-C3-C2	5.62	119.25	111.02
2	B	2	NAG	O5-C1-C2	-5.57	102.50	111.29
2	B	7	BMA	C2-C3-C4	-5.39	101.57	110.89
2	B	5	GLA	O4-C4-C5	5.24	122.31	109.30
2	B	5	GLA	O2-C2-C3	4.92	120.00	110.14
2	B	1	NAG	O5-C5-C6	-4.73	99.79	107.20
2	B	4	BMA	C2-C3-C4	4.51	118.71	110.89
2	B	2	NAG	O5-C5-C6	4.39	114.09	107.20
2	B	6	MAN	O2-C2-C3	4.37	118.88	110.14
2	B	6	MAN	C2-C3-C4	4.25	118.25	110.89
2	B	2	NAG	C4-C3-C2	4.08	117.00	111.02
2	B	7	BMA	O5-C1-C2	4.03	116.99	110.77
2	B	3	BMA	C3-C4-C5	-3.96	103.17	110.24
2	B	7	BMA	O6-C6-C5	3.81	124.35	111.29
2	B	2	NAG	O5-C5-C4	3.71	119.86	110.83
2	B	3	BMA	C1-C2-C3	3.70	114.22	109.67
2	B	7	BMA	O3-C3-C2	-3.68	102.95	109.99
2	B	5	GLA	O3-C3-C2	3.67	117.02	109.99
2	B	2	NAG	C1-C2-N2	-3.58	104.36	110.49
2	B	1	NAG	C8-C7-N2	3.55	122.11	116.10
2	B	6	MAN	O5-C5-C6	3.40	112.53	107.20
2	B	6	MAN	O3-C3-C4	3.30	117.98	110.35
2	B	7	BMA	C3-C4-C5	-3.28	104.39	110.24
2	B	3	BMA	C2-C3-C4	-3.20	105.37	110.89
2	B	1	NAG	C6-C5-C4	3.00	120.03	113.00
2	B	3	BMA	O4-C4-C5	2.94	116.60	109.30
2	B	6	MAN	C1-O5-C5	2.88	116.09	112.19
2	B	4	BMA	O2-C2-C1	2.86	115.01	109.15
2	B	1	NAG	O3-C3-C2	-2.77	103.74	109.47
2	B	1	NAG	O4-C4-C5	2.73	116.08	109.30
2	B	7	BMA	O3-C3-C4	2.55	116.24	110.35
2	B	3	BMA	O5-C1-C2	2.42	114.51	110.77
2	B	1	NAG	O4-C4-C3	2.42	115.94	110.35
2	B	7	BMA	O2-C2-C3	2.35	114.84	110.14
2	B	1	NAG	C3-C4-C5	2.28	114.31	110.24
2	B	6	MAN	O5-C5-C4	2.27	116.36	110.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	BMA	O2-C2-C3	2.24	114.63	110.14
2	B	5	GLA	O4-C4-C3	-2.23	105.19	110.35
2	B	5	GLA	O6-C6-C5	2.09	118.46	111.29
2	B	6	MAN	C6-C5-C4	2.03	117.75	113.00
2	B	1	NAG	O5-C1-C2	2.01	114.46	111.29

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	2	NAG	C3
2	B	6	MAN	C3

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2
2	B	6	MAN	O5-C5-C6-O6
2	B	6	MAN	C4-C5-C6-O6
2	B	4	BMA	C4-C5-C6-O6
2	B	2	NAG	C8-C7-N2-C2
2	B	4	BMA	O5-C5-C6-O6
2	B	7	BMA	O5-C5-C6-O6
2	B	3	BMA	O5-C5-C6-O6
2	B	7	BMA	C4-C5-C6-O6
2	B	3	BMA	C4-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6
2	B	1	NAG	C1-C2-N2-C7
2	B	2	NAG	O7-C7-N2-C2
2	B	1	NAG	C3-C2-N2-C7

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	7	BMA	2	0
2	B	3	BMA	3	0
2	B	2	NAG	2	0
2	B	4	BMA	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.