



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 23, 2023 – 02:11 AM EDT

PDB ID : 3DR0
Title : Structure of reduced cytochrome c6 from *Synechococcus* sp. PCC 7002
Authors : Bialek, W.; Krzywda, S.; Jaskolski, M.; Szczepaniak, A.
Deposited on : 2008-07-10
Resolution : 1.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2024 (1.28-1.20)
Clashscore	141614	1007 (1.26-1.22)
Ramachandran outliers	138981	2053 (1.28-1.20)
Sidechain outliers	138945	2051 (1.28-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	93	92% (green), 8% (yellow), 0% (orange), 0% (red), 0% (grey)
1	B	93	89% (green), 9% (yellow), 0% (orange), 0% (red), 0% (grey)
1	C	93	96% (green), 0% (yellow), 0% (orange), 0% (red), 0% (grey)

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	176	Total	O	0	0
			176	176		
4	B	203	Total	O	0	0
			203	203		
4	C	144	Total	O	0	0
			144	144		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cytochrome c6



- Molecule 1: Cytochrome c6



- Molecule 1: Cytochrome c6



4 Data and refinement statistics

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	82.88Å 82.88Å 28.28Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.58 – 1.23 19.58 – 1.23	Depositor EDS
% Data completeness (in resolution range)	98.8 (19.58-1.23) 98.8 (19.58-1.23)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 1.23Å)	Xtrriage
Refinement program	REFMAC 5.5.0039	Depositor
R, R_{free}	0.107 , 0.138 0.156 , 0.182	Depositor DCC
R_{free} test set	1274 reflections (2.03%)	wwPDB-VP
Wilson B-factor (Å ²)	13.8	Xtrriage
Anisotropy	0.005	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 71.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.026 for -h,-k,l 0.027 for h,-h-k,-l 0.019 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	2745	wwPDB-VP
Average B, all atoms (Å ²)	11.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.77	0/723	0.79	0/976
1	B	0.86	0/749	0.85	0/1010
1	C	0.81	0/722	0.79	0/974
All	All	0.81	0/2194	0.81	0/2960

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	691	0	642	14	0
1	B	701	0	654	12	0
1	C	691	0	646	9	0
2	A	43	0	30	1	0
2	B	43	0	30	3	0
2	C	43	0	30	2	0
3	B	10	0	0	0	0
4	A	176	0	0	12	0
4	B	203	0	0	8	0
4	C	144	0	0	7	0
All	All	2745	0	2032	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26[B]:MET:SD	4:B:283:HOH:O	1.94	1.21
1:A:93[B]:TRP:C	4:A:200:HOH:O	1.85	1.11
1:A:44:LYS:HE3	4:A:268:HOH:O	1.56	1.03
1:B:93[A]:TRP:C	4:B:176:HOH:O	1.98	1.00
1:B:8[A]:GLN:NE2	4:B:286:HOH:O	1.89	0.99
1:B:32[A]:LYS:NZ	4:B:244:HOH:O	1.92	0.97
1:C:93[A]:TRP:C	4:C:213:HOH:O	2.04	0.94
1:A:93[A]:TRP:OXT	4:A:264:HOH:O	1.84	0.93
1:A:26[B]:MET:HG2	4:A:220:HOH:O	1.68	0.92
1:A:93[B]:TRP:OXT	4:A:200:HOH:O	1.87	0.88
1:C:93[B]:TRP:O	4:C:234:HOH:O	1.99	0.80
1:C:28[A]:THR:HG23	4:C:203:HOH:O	1.84	0.77
1:B:93[A]:TRP:O	4:B:176:HOH:O	2.01	0.74
1:C:93[A]:TRP:OXT	4:C:213:HOH:O	2.00	0.74
1:C:28[B]:THR:OG1	4:C:204:HOH:O	2.06	0.72
1:B:74:ASP:OD2	4:B:266:HOH:O	2.13	0.66
1:B:49:SER:OG	1:B:52[B]:GLU:HG3	1.99	0.62
1:A:26[A]:MET:HE3	4:A:220:HOH:O	2.01	0.60
1:A:26[A]:MET:CE	4:A:220:HOH:O	2.49	0.60
1:A:93[A]:TRP:C	4:A:264:HOH:O	2.33	0.59
1:C:93[A]:TRP:O	4:C:213:HOH:O	2.17	0.59
1:C:93[B]:TRP:C	4:C:234:HOH:O	2.39	0.56
1:B:84[B]:ILE:HD11	2:B:94:HEM:HMB3	1.86	0.56
1:B:93[A]:TRP:OXT	4:B:176:HOH:O	2.15	0.55
1:A:93[B]:TRP:O	4:A:200:HOH:O	2.10	0.54
1:B:26[B]:MET:SD	2:B:94:HEM:O2D	2.66	0.53
1:A:65:MET:HB2	2:A:94:HEM:C4D	2.44	0.52
1:C:65:MET:HB2	2:C:94:HEM:C4D	2.48	0.48
1:B:93[B]:TRP:OXT	4:B:177:HOH:O	2.20	0.47
1:B:65:MET:HB2	2:B:94:HEM:C4D	2.50	0.46
1:C:84:ILE:HD11	2:C:94:HEM:HMB3	1.98	0.45
1:A:76:ASP:OD1	4:A:142:HOH:O	2.22	0.43
1:A:51:GLU:HG3	4:A:186:HOH:O	2.19	0.43
1:A:26[A]:MET:HE2	4:A:220:HOH:O	2.17	0.42
1:A:62:GLN:O	1:A:65:MET:HB3	2.19	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/93 (102%)	93 (98%)	2 (2%)	0	100	100
1	B	98/93 (105%)	96 (98%)	2 (2%)	0	100	100
1	C	95/93 (102%)	93 (98%)	2 (2%)	0	100	100
All	All	288/279 (103%)	282 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	65/60 (108%)	65 (100%)	0	100	100
1	B	68/60 (113%)	63 (93%)	5 (7%)	13	0
1	C	65/60 (108%)	63 (97%)	2 (3%)	40	6
All	All	198/180 (110%)	191 (96%)	7 (4%)	52	5

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	26[A]	MET
1	B	26[B]	MET
1	B	57	GLN
1	B	93[A]	TRP
1	B	93[B]	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	93[A]	TRP
1	C	93[B]	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	96	-	4,4,4	0.23	0	6,6,6	0.18	0
2	HEM	B	94	1	41,50,50	1.68	8 (19%)	45,82,82	1.73	13 (28%)
2	HEM	A	94	1	41,50,50	1.57	8 (19%)	45,82,82	1.56	10 (22%)
3	SO4	B	95	-	4,4,4	0.49	0	6,6,6	1.09	0
2	HEM	C	94	1	41,50,50	1.56	6 (14%)	45,82,82	1.64	10 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	94	1	-	4/12/54/54	-
2	HEM	C	94	1	-	4/12/54/54	-
2	HEM	A	94	1	-	5/12/54/54	-

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	94	HEM	C3D-C2D	5.12	1.47	1.36
2	B	94	HEM	C3D-C2D	4.91	1.47	1.36
2	C	94	HEM	C3C-C2C	-4.17	1.34	1.40
2	A	94	HEM	C3D-C2D	4.14	1.45	1.36
2	B	94	HEM	C3C-C2C	-3.88	1.35	1.40
2	A	94	HEM	C3C-CAC	3.67	1.55	1.47
2	A	94	HEM	C3C-C2C	-3.27	1.35	1.40
2	C	94	HEM	C3C-CAC	3.00	1.53	1.47
2	A	94	HEM	CBB-CAB	2.98	1.45	1.30
2	B	94	HEM	CBB-CAB	2.78	1.44	1.30
2	B	94	HEM	C3C-CAC	2.77	1.53	1.47
2	C	94	HEM	C3B-C2B	-2.76	1.31	1.37
2	A	94	HEM	CAB-C3B	2.58	1.54	1.47
2	B	94	HEM	O1A-CGA	2.53	1.30	1.22
2	B	94	HEM	O2D-CGD	-2.44	1.22	1.30
2	A	94	HEM	C1B-NB	-2.38	1.36	1.40
2	A	94	HEM	O2D-CGD	-2.28	1.23	1.30
2	C	94	HEM	CBB-CAB	2.22	1.41	1.30
2	B	94	HEM	CAB-C3B	2.15	1.53	1.47
2	A	94	HEM	O1A-CGA	2.10	1.29	1.22
2	B	94	HEM	C4D-ND	-2.04	1.36	1.40
2	C	94	HEM	CAB-C3B	2.02	1.52	1.47

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	94	HEM	C2C-C3C-C4C	4.89	110.32	106.90
2	C	94	HEM	C4D-ND-C1D	4.30	109.52	105.07
2	C	94	HEM	C3B-C2B-C1B	3.68	109.22	106.49
2	A	94	HEM	CMA-C3A-C4A	-3.57	122.98	128.46
2	C	94	HEM	C4C-CHD-C1D	3.51	127.19	122.56
2	B	94	HEM	CMD-C2D-C1D	3.33	130.12	125.04
2	A	94	HEM	C4D-ND-C1D	3.33	108.52	105.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	94	HEM	CMA-C3A-C4A	-3.21	123.53	128.46
2	A	94	HEM	CMD-C2D-C1D	3.14	129.82	125.04
2	A	94	HEM	C1B-NB-C4B	3.05	108.23	105.07
2	C	94	HEM	CMD-C2D-C1D	3.00	129.61	125.04
2	C	94	HEM	CMA-C3A-C4A	-2.95	123.93	128.46
2	B	94	HEM	CMA-C3A-C2A	2.83	130.28	124.94
2	B	94	HEM	C1B-NB-C4B	2.80	107.97	105.07
2	A	94	HEM	CMA-C3A-C2A	2.71	130.06	124.94
2	A	94	HEM	C4C-CHD-C1D	2.64	126.05	122.56
2	B	94	HEM	C4C-CHD-C1D	2.61	126.00	122.56
2	A	94	HEM	CBB-CAB-C3B	-2.60	114.69	127.62
2	B	94	HEM	CMC-C2C-C3C	2.57	129.48	124.68
2	B	94	HEM	CHD-C1D-ND	2.55	127.20	124.43
2	C	94	HEM	CBB-CAB-C3B	-2.46	115.37	127.62
2	B	94	HEM	CBB-CAB-C3B	-2.45	115.41	127.62
2	A	94	HEM	C4B-CHC-C1C	2.41	125.74	122.56
2	C	94	HEM	CAD-C3D-C4D	2.29	128.66	124.66
2	B	94	HEM	C3C-C4C-NC	-2.22	106.76	110.94
2	B	94	HEM	O2A-CGA-CBA	2.17	121.02	114.03
2	B	94	HEM	C4D-ND-C1D	2.17	107.32	105.07
2	C	94	HEM	CBD-CAD-C3D	-2.17	106.60	112.63
2	A	94	HEM	CAD-CBD-CGD	-2.09	109.10	113.60
2	B	94	HEM	C1D-C2D-C3D	-2.09	104.76	106.96
2	C	94	HEM	C4B-CHC-C1C	2.07	125.29	122.56
2	C	94	HEM	CBA-CAA-C2A	-2.05	109.13	112.62
2	A	94	HEM	CBD-CAD-C3D	-2.00	107.06	112.63

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	94	HEM	C2B-C3B-CAB-CBB
2	B	94	HEM	C4B-C3B-CAB-CBB
2	A	94	HEM	C2B-C3B-CAB-CBB
2	B	94	HEM	C2B-C3B-CAB-CBB
2	A	94	HEM	C4B-C3B-CAB-CBB
2	C	94	HEM	C4B-C3B-CAB-CBB
2	B	94	HEM	CAD-CBD-CGD-O1D
2	B	94	HEM	CAD-CBD-CGD-O2D
2	C	94	HEM	CAD-CBD-CGD-O2D
2	A	94	HEM	CAD-CBD-CGD-O2D
2	C	94	HEM	CAD-CBD-CGD-O1D

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	94	HEM	CAD-CBD-CGD-O1D
2	A	94	HEM	CAA-CBA-CGA-O2A


There are no ring outliers.


3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	94	HEM	3	0
2	A	94	HEM	1	0
2	C	94	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.




6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HEM A 94:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 94:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

Unable to reproduce the depositor's R factor - this section is therefore empty.