

# wwPDB X-ray Structure Validation Summary Report (i)

#### Sep 26, 2024 – 12:06 PM EDT

PDB ID : 9DRF

Title : Crystal structure of ADP-ribose diphosphatase from Klebsiella pneumoniae

(DTP bound, P21 form)

Authors: Seattle Structural Genomics Center for Infectious Disease; Seattle Structural

Genomics Center for Infectious Disease (SSGCID)

Deposited on : 2024-09-25

Resolution : 1.78 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 2022.3.0, CSD as543be (2022)

Xtriage (Phenix) : 1.20.1

EDS: 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.002 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.38.3

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$ 

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\#  ext{Entries},  ext{ resolution range}( ext{Å}))$
$R_{free}$	164625	1191 (1.78-1.78)
Clashscore	180529	1282 (1.78-1.78)
Ramachandran outliers	177936	1270 (1.78-1.78)
Sidechain outliers	177891	1270 (1.78-1.78)
RSRZ outliers	164620	1191 (1.78-1.78)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	218	83%	8%	10%
1	В	218	78%	11%	10%
1	С	218	85%	6%	9%
1	D	218	83%	7%	11%



Continued from previous page...

Mol	Chain	Length	Quality of chain		
1	Б	010	17%		
1	Е	218	85%	7%	7%
1	F	218	17%	20/	110/
1	I.	210	83%	6%	11%
1	G	218	81%	10%	9%
			15%		
1	Н	218	83%	8%	8%



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 13581 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called ADP-ribose pyrophosphatase.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace	
1	A	197	Total C N O S	0	5	0	
1	A	197	1597 1014 283 297 3				
1	В	197	Total C N O S	0	2	0	
1	Б	197	1571 994 275 299 3		2		
1	С	199	Total C N O S	0	2	0	
1		199	1585 1004 277 301 3	U	2		
1	D	195	Total C N O S	0	1	0	
1	D	190	1559 985 275 296 3	0	1	0	
1	E	202	Total C N O S	0	3	0	
1	Ľ	202	1626 1031 290 301 4	0	3		
1	F	194	Total C N O S	0	3	0	
1	I.	194	1545 981 267 294 3		3		
1	G	198	Total C N O S	0	2	0	
1	G	130	1577 1001 276 297 3		2	U	
1	Н	200	Total C N O S	0	4	0	
1	11	200	1596 1012 279 302 3		<del>'1</del>		

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP A0A0H3GVQ7
A	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
A	-5	HIS	ı	expression tag	UNP A0A0H3GVQ7
A	-4	HIS	ı	expression tag	UNP A0A0H3GVQ7
A	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
A	-2	HIS	ı	expression tag	UNP A0A0H3GVQ7
A	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
A	0	HIS	ı	expression tag	UNP A0A0H3GVQ7
В	-7	MET	-	expression tag	UNP A0A0H3GVQ7
В	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
В	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
В	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
В	-3	HIS	-	expression tag	UNP A0A0H3GVQ7



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
В	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
В	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
В	0	HIS	-	expression tag	UNP A0A0H3GVQ7
С	-7	MET	_	expression tag	UNP A0A0H3GVQ7
С	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
С	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
С	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
С	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
С	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
С	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
С	0	HIS	-	expression tag	UNP A0A0H3GVQ7
D	-7	MET	-	expression tag	UNP A0A0H3GVQ7
D	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
D	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
D	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
D	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
D	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
D	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
D	0	HIS	-	expression tag	UNP A0A0H3GVQ7
Е	-7	MET	-	expression tag	UNP A0A0H3GVQ7
Е	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
Е	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
Е	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
Е	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
Е	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
Е	-1	HIS	_	expression tag	UNP A0A0H3GVQ7
Е	0	HIS	-	expression tag	UNP A0A0H3GVQ7
F	-7	MET	-	expression tag	UNP A0A0H3GVQ7
F	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
F	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
F	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
F	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
F	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
F	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
F	0	HIS	-	expression tag	UNP A0A0H3GVQ7
G	-7	MET	-	expression tag	UNP A0A0H3GVQ7
G	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
G	-5	HIS	-	expression tag	UNP A0A0H3GVQ7
G	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
G	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
G	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
G	-1	HIS	_	expression tag	UNP A0A0H3GVQ7



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	0	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	-7	MET	-	expression tag	UNP A0A0H3GVQ7
Н	-6	ALA	-	expression tag	UNP A0A0H3GVQ7
Н	-5	HIS	ı	expression tag	UNP A0A0H3GVQ7
Н	-4	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	-3	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	-2	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	-1	HIS	-	expression tag	UNP A0A0H3GVQ7
Н	0	HIS	-	expression tag	UNP A0A0H3GVQ7

• Molecule 2 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (three-letter code: DTP) (formula:  $C_{10}H_{16}N_5O_{12}P_3$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	A	1	Total	С	N	О	Р	0	0
2	Λ	1	30	10	5	12	3	U	U
2	В	1	Total	С	N	О	Р	0	0
2	Б	1	30	10	5	12	3	U	U
2	С	1	Total	С	N	О	Р	0	0
2		1	30	10	5	12	3	U	
2	C	1	Total	С	N	Ο	Р	0	0
		1	30	10	5	12	3	U	U
2	E	1	Total	С	N	О	Р	0	0
	ت ا	1	30	10	5	12	3	U	U
2	Е	1	Total	С	N	О	Р	0	0
	שנו	1	30	10	5	12	3	U	U



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
9	G	1	Total	С	N	О	Р	0	0
	G	1	30	10	5	12	3	U	0
9	С	1	Total	С	N	О	Р	0	0
	G	1	30	10	5	12	3	U	U

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	E	1	Total Mg 1 1	0	0
3	F	2	Total Mg 2 2	0	0
3	G	1	Total Mg 1 1	0	0
3	Н	1	Total Mg 1 1	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	91	Total O 91 91	0	0
4	В	85	Total O 85 85	0	1
4	С	103	Total O 104 104	0	1
4	D	63	Total O 63 63	0	0
4	E	78	Total O 78 78	0	0
4	F	83	Total O 83 83	0	0
4	G	95	Total O 95 95	0	0
4	Н	79	Total O 79 79	0	0




# 3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ADP-ribose pyrophosphatase












• Molecule 1: ADP-ribose pyrophosphatase



• Molecule 1: ADP-ribose pyrophosphatase



• Molecule 1: ADP-ribose pyrophosphatase





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	93.03Å 80.13Å 116.19Å	Donogitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.13^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	49.23 - 1.78	Depositor
rtesolution (A)	49.23 - 1.78	EDS
% Data completeness	99.8 (49.23-1.78)	Depositor
(in resolution range)	99.9 (49.23-1.78)	EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.13 (at 1.78Å)	Xtriage
Refinement program	PHENIX (dev_5295: ???)	Depositor
Ρ. Р.	0.179 , 0.217	Depositor
$R, R_{free}$	0.186 , $0.224$	DCC
$R_{free}$ test set	8046 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.5	Xtriage
Anisotropy	0.676	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	$0.35 \; ,  45.4$	EDS
L-test for twinning <sup>2</sup>	$< L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.040 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	13581	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 55.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.1293e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: DTP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z >5	RMSZ	# Z  > 5	
1	A	0.51	0/1642	0.69	0/2226	
1	В	0.49	0/1606	0.69	0/2174	
1	С	0.52	0/1621	0.71	0/2194	
1	D	0.49	0/1591	0.70	0/2155	
1	Е	0.48	0/1669	0.69	0/2259	
1	F	0.48	0/1583	0.67	0/2145	
1	G	0.52	0/1613	0.71	0/2184	
1	Н	0.52	0/1638	0.72	0/2220	
All	All	0.50	0/12963	0.70	0/17557	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1597	0	1561	12	0
1	В	1571	0	1535	16	0
1	С	1585	0	1552	8	0
1	D	1559	0	1525	10	0
1	Е	1626	0	1592	12	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	1545	0	1517	9	0
1	G	1577	0	1547	17	0
1	Н	1596	0	1553	14	0
2	A	30	0	12	1	0
2	В	30	0	12	0	0
2	С	60	0	24	1	0
2	Е	60	0	24	2	0
2	G	60	0	24	1	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	Е	1	0	0	0	0
3	F	2	0	0	0	0
3	G	1	0	0	0	0
3	Н	1	0	0	0	0
4	A	91	0	0	0	0
4	В	85	0	0	1	0
4	С	104	0	0	1	0
4	D	63	0	0	0	0
4	Е	78	0	0	0	0
4	F	83	0	0	0	0
4	G	95	0	0	0	0
4	Н	79	0	0	1	0
All	All	13581	0	12478	85	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 3.

The worst 5 of 85 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	$\operatorname{distance}\ ( ext{Å})$	overlap (Å)
1:B:75:VAL:HG13	1:B:166:ILE:HG23	1.50	0.94
1:F:75:VAL:HG13	1:F:166:ILE:HG23	1.60	0.84
1:E:75[B]:VAL:HG23	1:E:166:ILE:HG23	1.62	0.80
1:H:75:VAL:HG13	1:H:166:ILE:HG23	1.63	0.79
1:G:81:ALA:HB2	1:H:53:ILE:HD12	1.65	0.78

There are no symmetry-related clashes.



## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$196/218 \; (90\%)$	186 (95%)	10 (5%)	0	100	100
1	В	195/218 (89%)	186 (95%)	9 (5%)	0	100	100
1	С	197/218 (90%)	191 (97%)	6 (3%)	0	100	100
1	D	192/218 (88%)	182 (95%)	10 (5%)	0	100	100
1	E	199/218 (91%)	194 (98%)	5 (2%)	0	100	100
1	F	193/218 (88%)	185 (96%)	7 (4%)	1 (0%)	25	11
1	G	196/218 (90%)	191 (97%)	5 (3%)	0	100	100
1	Н	200/218 (92%)	190 (95%)	10 (5%)	0	100	100
All	All	1568/1744 (90%)	1505 (96%)	62 (4%)	1 (0%)	48	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	9	ILE

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	168/183~(92%)	164 (98%)	4 (2%)	44 24		
1	В	163/183~(89%)	157 (96%)	6 (4%)	29 10		
1	С	165/183~(90%)	163 (99%)	2 (1%)	67 54		
1	D	163/183 (89%)	162 (99%)	1 (1%)	84 78		



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	E	171/183 (93%)	169 (99%)	2 (1%)	67 54
1	F	162/183 (88%)	162 (100%)	0	100 100
1	G	164/183 (90%)	164 (100%)	0	100 100
1	Н	165/183 (90%)	161 (98%)	4 (2%)	44 24
All	All	1321/1464 (90%)	1302 (99%)	19 (1%)	62 47

5 of 19 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Е	144	LEU
1	Н	98	MET
1	Н	167	ARG
1	Н	34	TYR
1	В	167	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	175	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 15 ligands modelled in this entry, 7 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and



the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Iol Type Chain Res Link			Bo	ond leng	ths	Bond angles			
MIOI	туре	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DTP	В	301	-	28,32,32	0.76	1 (3%)	35,50,50	0.77	1 (2%)
2	DTP	Е	301	-	28,32,32	1.05	2 (7%)	35,50,50	0.86	2 (5%)
2	DTP	G	302	-	28,32,32	1.28	2 (7%)	35,50,50	0.86	1 (2%)
2	DTP	A	301	-	28,32,32	0.93	2 (7%)	35,50,50	0.86	1 (2%)
2	DTP	С	301	-	28,32,32	0.75	1 (3%)	35,50,50	0.78	1 (2%)
2	DTP	Е	302	-	28,32,32	1.20	2 (7%)	35,50,50	0.77	0
2	DTP	С	302	-	28,32,32	0.68	0	35,50,50	0.78	0
2	DTP	G	301	-	28,32,32	1.04	2 (7%)	35,50,50	0.81	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DTP	В	301	-	-	5/18/34/34	0/3/3/3
2	DTP	Е	301	-	-	6/18/34/34	0/3/3/3
2	DTP	G	302	_	-	3/18/34/34	0/3/3/3
2	DTP	A	301	-	-	6/18/34/34	0/3/3/3
2	DTP	С	301	-	-	10/18/34/34	0/3/3/3
2	DTP	Е	302	-	-	8/18/34/34	0/3/3/3
2	DTP	С	302	-	-	7/18/34/34	0/3/3/3
2	DTP	G	301	-	-	3/18/34/34	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$Ideal(\AA)$
2	G	302	DTP	PA-O3A	5.56	1.65	1.59
2	Е	302	DTP	PA-O3A	4.27	1.64	1.59
2	Е	301	DTP	PA-O3A	3.97	1.63	1.59
2	G	301	DTP	PA-O3A	3.84	1.63	1.59
2	Е	302	DTP	PB-O3B	3.58	1.63	1.59



The worst	5	$\circ f$	7	bond	angle	outliers	are	listed	below.
THE WOLDS	$\cdot$	OI		DOM	angic	Outilities	arc	nouca	DCIOW.

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	В	301	DTP	C5-C6-N6	2.32	123.85	120.31
2	Е	301	DTP	C5-C6-N6	2.31	123.83	120.31
2	С	301	DTP	C5-C6-N6	2.31	123.83	120.31
2	G	301	DTP	C5-C6-N6	2.26	123.75	120.31
2	G	302	DTP	C5-C6-N6	2.18	123.62	120.31

There are no chirality outliers.

5 of 48 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	DTP	C4'-C5'-O5'-PA
2	В	301	DTP	PB-O3B-PG-O2G
2	С	301	DTP	C5'-O5'-PA-O1A
2	С	301	DTP	C5'-O5'-PA-O2A
2	С	301	DTP	C5'-O5'-PA-O3A


There are no ring outliers.

4 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	302	DTP	1	0
2	A	301	DTP	1	0
2	Е	302	DTP	2	0
2	С	302	DTP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







































# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$		$\mathrm{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	197/218 (90%)	0.77	23 (11%) 10 1	12	21, 42, 78, 105	5 (2%)
1	В	197/218 (90%)	0.83	29 (14%) 7 7	7	25, 44, 86, 102	2 (1%)
1	С	199/218 (91%)	0.45	10 (5%) 35 41	1	20, 40, 62, 89	2 (1%)
1	D	195/218 (89%)	0.96	23 (11%) 10 1	12	28, 50, 84, 97	1 (0%)
1	E	202/218 (92%)	1.19	37 (18%) 4 4	1	21, 51, 76, 88	3 (1%)
1	F	194/218 (88%)	0.98	38 (19%) 4 4	1	26, 44, 82, 106	3 (1%)
1	G	198/218 (90%)	0.62	18 (9%) 16 20	0	20, 40, 69, 88	2 (1%)
1	Н	200/218 (91%)	0.82	33 (16%) 5 6	3	21, 44, 81, 104	4 (2%)
All	All	1582/1744 (90%)	0.83	211 (13%) 8	9	20, 44, 80, 106	22 (1%)

The worst 5 of 211 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	156	ILE	5.4
1	A	0	HIS	5.3
1	A	10	THR	5.0
1	A	-3	HIS	4.9
1	Н	156	ILE	4.6

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	DTP	В	301	30/30	0.59	0.13	55,92,128,133	0
2	DTP	Е	302	30/30	0.60	0.15	40,86,125,129	0
2	DTP	С	302	30/30	0.64	0.14	40,81,121,122	0
2	DTP	G	301	30/30	0.64	0.13	48,82,120,126	0
2	DTP	С	301	30/30	0.65	0.12	47,83,125,132	0
2	DTP	Е	301	30/30	0.65	0.13	56,83,116,121	0
2	DTP	A	301	30/30	0.66	0.14	33,67,119,130	0
3	MG	G	303	1/1	0.66	0.17	62,62,62,62	0
2	DTP	G	302	30/30	0.70	0.14	30,72,118,129	0
3	MG	Е	303	1/1	0.75	0.15	64,64,64,64	0
3	MG	F	302	1/1	0.77	0.14	61,61,61,61	0
3	MG	В	302	1/1	0.77	0.21	72,72,72,72	0
3	MG	С	303	1/1	0.82	0.12	56,56,56,56	0
3	MG	Н	301	1/1	0.85	0.16	59,59,59,59	0
3	MG	F	301	1/1	0.89	0.18	52,52,52,52	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




#### Electron density around DTP B 301:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









#### Electron density around DTP E 302:












### Electron density around DTP C 302:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$  (at 0.7 rmsd) in gray  ${\rm mF}_o\text{-}{\rm DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









#### Electron density around DTP G 301:












### Electron density around DTP C 301:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)









#### Electron density around DTP E 301:












### Electron density around DTP A 301:











#### Electron density around MG G 303:

 $2 {
m mF}_o {
m -DF}_c$  (at 0.7 rmsd) in gray  ${
m mF}_o {
m -DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)







































# 6.5 Other polymers (i)

There are no such residues in this entry.

