



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 08:18 pm GMT

PDB ID : 3DVG  
Title : Crystal structure of K63-specific fab Apu.3A8 bound to K63-linked di-ubiquitin  
Authors : Hymowitz, S.G.  
Deposited on : 2008-07-18  
Resolution : 2.60 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122126	3110 (2.60-2.60)
Ramachandran outliers	120053	3062 (2.60-2.60)
Sidechain outliers	120020	3062 (2.60-2.60)
RSRZ outliers	108989	2706 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	
2	B	230	
3	X	80	
4	Y	79	

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4491 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Human IgG1 fab fragment light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	210	1613	1010	272	327	4	0	0	0

- Molecule 2 is a protein called Human IgG1 fab fragment heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	221	1658	1053	275	325	5	0	0	0

- Molecule 3 is a protein called Ubiquitin D77.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	X	75	592	374	101	116	1	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	-2	GLY	-	EXPRESSION TAG	UNP P62988
X	-1	SER	-	EXPRESSION TAG	UNP P62988
X	0	HIS	-	EXPRESSION TAG	UNP P62988
X	77	ASP	-	ENGINEERED	UNP P62988

- Molecule 4 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	Y	76	603	378	107	117	1	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Y	-2	GLY	-	EXPRESSION TAG	UNP P62988
Y	-1	SER	-	EXPRESSION TAG	UNP P62988
Y	0	HIS	-	EXPRESSION TAG	UNP P62988
Y	63	ARG	LYS	ENGINEERED	UNP P62988


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	18	Total O 18 18	0	0
5	B	7	Total O 7 7	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Human IgG1 fab fragment light chain




- Molecule 2: Human IgG1 fab fragment heavy chain



- Molecule 3: Ubiquitin D77



- Molecule 4: Ubiquitin



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.81Å 88.12Å 90.23Å 90.00° 108.28° 90.00°	Depositor
Resolution (Å)	48.00 – 2.60 42.84 – 2.60	Depositor EDS
% Data completeness (in resolution range)	97.9 (48.00-2.60) 97.9 (42.84-2.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.219 , 0.261 0.214 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	59.4	Xtrriage
Anisotropy	0.392	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 69.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4491	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.43% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.44	0/1648	0.58	0/2236
2	B	0.40	0/1701	0.58	0/2324
3	X	0.82	3/598 (0.5%)	0.60	0/806
4	Y	0.45	1/609 (0.2%)	0.56	0/819
All	All	0.50	4/4556 (0.1%)	0.58	0/6185

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	X	10	GLY	C-O	10.84	1.41	1.23
3	X	10	GLY	C-N	6.82	1.49	1.34
3	X	73	LEU	C-O	6.56	1.35	1.23
4	Y	75	GLY	C-N	5.86	1.43	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1613	0	1570	18	0
2	B	1658	0	1618	19	0
3	X	592	0	612	8	0
4	Y	603	0	629	4	0
5	A	18	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	7	0	0	0	0
All	All	4491	0	4429	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:29:VAL:HG21	1:A:90:GLN:HB3	1.57	0.84
2:B:104:TYR:O	2:B:105:THR:HB	1.76	0.83
1:A:25:ALA:HB3	1:A:69:THR:O	1.82	0.78
1:A:200:HIS:CD2	1:A:202:GLY:H	2.18	0.61
2:B:97:ALA:HB1	2:B:107:ILE:CG2	2.31	0.60
2:B:35:HIS:HD2	2:B:47:TRP:HE1	1.49	0.59
3:X:1:MET:HG3	3:X:63:LYS:HG2	1.83	0.59
2:B:64:VAL:HG13	2:B:68:PHE:HB2	1.84	0.58
1:A:160:ASN:HD22	1:A:160:ASN:H	1.53	0.57
2:B:154:PRO:O	2:B:207:HIS:HE1	1.89	0.55
2:B:126:PRO:HB3	2:B:152:TYR:HB3	1.91	0.52
4:Y:4:PHE:HE1	4:Y:64:GLU:HG3	1.75	0.52
2:B:16:GLY:O	2:B:86:LEU:HB2	2.10	0.51
3:X:7:THR:HG22	3:X:69:LEU:HD23	1.93	0.51
2:B:35:HIS:CD2	2:B:47:TRP:HE1	2.28	0.50
1:A:162:GLN:HG2	2:B:176:VAL:HG11	1.94	0.49
1:A:142:TYR:CD2	1:A:143:PRO:HA	2.48	0.49
2:B:134:SER:O	2:B:138:THR:HG23	2.13	0.48
2:B:119:SER:HB3	2:B:153:PHE:HZ	1.79	0.47
1:A:142:TYR:CG	1:A:143:PRO:HA	2.49	0.47
2:B:33:LEU:HB2	2:B:99:GLU:HG2	1.97	0.47
3:X:60:ASN:ND2	3:X:62:GLN:HE21	2.12	0.47
1:A:162:GLN:CG	2:B:176:VAL:HG11	2.45	0.46
2:B:35:HIS:HD2	2:B:47:TRP:NE1	2.13	0.46
1:A:29:VAL:CG2	1:A:90:GLN:HB3	2.38	0.46
2:B:161:TRP:CH2	2:B:203:CYS:HB3	2.50	0.46
1:A:37:GLN:HB2	1:A:47:LEU:HD11	1.97	0.46
1:A:143:PRO:O	1:A:200:HIS:HE1	1.99	0.46
2:B:128:VAL:O	2:B:216:LYS:HE3	2.17	0.45
3:X:5:VAL:HB	3:X:13:ILE:HG13	1.99	0.45
2:B:206:ASN:HD21	2:B:208:LYS:HE3	1.83	0.44
1:A:42:LYS:HG2	1:A:43:ALA:H	1.83	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:X:45:PHE:HB2	3:X:67:LEU:HD22	2.00	0.42
4:Y:26:VAL:O	4:Y:30:ILE:HG13	2.20	0.42
3:X:6:LYS:HG2	3:X:12:THR:OG1	2.20	0.41
1:A:118:PHE:CD1	2:B:137:SER:HA	2.55	0.41
1:A:86:TYR:O	1:A:103:GLY:HA2	2.20	0.41
4:Y:31:GLN:HE21	4:Y:37:PRO:HA	1.84	0.41
1:A:162:GLN:HG2	2:B:176:VAL:CG1	2.50	0.41
1:A:151:LYS:HB2	1:A:195:ALA:HB3	2.02	0.41
3:X:55:THR:O	3:X:58:ASP:HB2	2.20	0.41
4:Y:37:PRO:HA	4:Y:38:PRO:HD3	1.89	0.41
1:A:33:VAL:HG21	1:A:71:PHE:CD2	2.56	0.40
1:A:160:ASN:HD22	1:A:160:ASN:N	2.11	0.40
3:X:30:ILE:O	3:X:34:GLU:HB2	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	208/217 (96%)	205 (99%)	3 (1%)	0	100	100
2	B	219/230 (95%)	209 (95%)	9 (4%)	1 (0%)	31	56
3	X	73/80 (91%)	73 (100%)	0	0	100	100
4	Y	74/79 (94%)	69 (93%)	5 (7%)	0	100	100
All	All	574/606 (95%)	556 (97%)	17 (3%)	1 (0%)	49	74

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	105	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/192 (96%)	173 (94%)	12 (6%)	19	38
2	B	182/192 (95%)	164 (90%)	18 (10%)	8	16
3	X	67/71 (94%)	63 (94%)	4 (6%)	21	42
4	Y	68/70 (97%)	61 (90%)	7 (10%)	8	14
All	All	502/525 (96%)	461 (92%)	41 (8%)	12	25

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	THR
1	A	22	THR
1	A	28	SER
1	A	73	LEU
1	A	107	GLU
1	A	110	ARG
1	A	112	VAL
1	A	127	LEU
1	A	149	GLN
1	A	160	ASN
1	A	162	GLN
1	A	189	GLU
2	B	11	LEU
2	B	12	VAL
2	B	18	LEU
2	B	20	LEU
2	B	52	THR
2	B	57	SER
2	B	62	ASP
2	B	85	SER
2	B	86	LEU
2	B	98	ARG
2	B	102	ARG
2	B	105	THR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	117	THR
2	B	123	THR
2	B	166	LEU
2	B	179	SER
2	B	199	GLN
2	B	211	ASN
3	X	2	GLN
3	X	9	THR
3	X	15	LEU
3	X	36	ILE
4	Y	17	VAL
4	Y	33	LYS
4	Y	42	ARG
4	Y	54	ARG
4	Y	60	ASN
4	Y	63	ARG
4	Y	74	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	90	GLN
1	A	126	GLN
1	A	149	GLN
1	A	157	GLN
1	A	160	ASN
1	A	162	GLN
1	A	200	HIS
2	B	35	HIS
2	B	199	GLN
2	B	206	ASN
2	B	207	HIS
3	X	2	GLN
3	X	40	GLN
3	X	49	GLN
3	X	60	ASN
4	Y	31	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	210/217 (96%)	0.49	5 (2%) 59 52	51, 62, 76, 85	0
2	B	221/230 (96%)	0.61	11 (4%) 29 22	45, 61, 72, 85	0
3	X	75/80 (93%)	1.00	12 (16%) 1 1	57, 69, 81, 84	1 (1%)
4	Y	76/79 (96%)	1.52	20 (26%) 0 0	49, 67, 87, 93	0
All	All	582/606 (96%)	0.73	48 (8%) 11 8	45, 63, 79, 93	1 (0%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	Y	4	PHE	7.8
3	X	40	GLN	6.2
3	X	73	LEU	5.9
4	Y	2	GLN	5.6
4	Y	19	PRO	4.7
4	Y	12	THR	4.7
4	Y	63	ARG	4.6
4	Y	3	ILE	4.2
4	Y	15	LEU	4.2
4	Y	66	THR	4.2
2	B	211	ASN	3.9
4	Y	53	GLY	3.8
4	Y	64	GLU	3.7
4	Y	21	ASP	3.7
4	Y	48	LYS	3.5
2	B	1	GLU	3.4
1	A	128	LYS	3.3
3	X	10	GLY	3.2
4	Y	6	LYS	3.2
1	A	5	THR	3.2
1	A	192	LYS	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
4	Y	5	VAL	3.1
4	Y	62	GLN	3.1
2	B	7	SER	3.0
4	Y	22	THR	3.0
2	B	29	VAL	3.0
3	X	37	PRO	2.9
3	X	72	ARG	2.8
2	B	27	PHE	2.8
3	X	70	VAL	2.8
3	X	32	ASP	2.5
4	Y	54	ARG	2.5
3	X	42	ARG	2.5
3	X	39	ASP	2.5
2	B	84	ASN	2.5
2	B	206	ASN	2.5
2	B	114	THR	2.4
4	Y	1	MET	2.4
3	X	23	ILE	2.3
3	X	36	ILE	2.3
1	A	201	GLN	2.2
1	A	207	VAL	2.2
4	Y	9	THR	2.2
4	Y	8	LEU	2.2
2	B	198	THR	2.1
2	B	12	VAL	2.1
2	B	4	LEU	2.1
3	X	24	GLU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.