



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 15, 2023 – 06:18 AM EDT

PDB ID : 8DXT  
Title : Fab arm of antibody GAR12 bound to the receptor binding domain of SARS-CoV-2.  
Authors : Langley, D.B.; Christ, D.; Henry, J.Y.  
Deposited on : 2022-08-03  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	214	 15% 81% 17% ..
2	H	234	 11% 78% 13% 8%
3	R	204	 90% 6% .
4	A	3	 33% 67%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4857 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Light chain of Fab arm of antibody GAR12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	211	1586	993	265	322	6	0	1	0

- Molecule 2 is a protein called Heavy chain of Fab arm of antibody GAR12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	215	1603	1014	267	316	6	0	2	0


- Molecule 3 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	R	196	1540	987	259	286	8	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	529	GLY	-	expression tag	UNP P0DTC2
R	530	SER	-	expression tag	UNP P0DTC2
R	531	HIS	-	expression tag	UNP P0DTC2
R	532	HIS	-	expression tag	UNP P0DTC2
R	533	HIS	-	expression tag	UNP P0DTC2
R	534	HIS	-	expression tag	UNP P0DTC2
R	535	HIS	-	expression tag	UNP P0DTC2
R	536	HIS	-	expression tag	UNP P0DTC2

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	A	3	38	22	2	14	0	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	L	18	Total	O	0	0
			18	18		
5	H	15	Total	O	0	0
			15	15		
5	R	57	Total	O	0	0
			57	57		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Light chain of Fab arm of antibody GAR12



- Molecule 2: Heavy chain of Fab arm of antibody GAR12



- Molecule 3: Spike protein S1



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.61Å 41.95Å 183.03Å 90.00° 97.08° 90.00°	Depositor
Resolution (Å)	48.09 – 2.25 48.08 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.09-2.25) 99.9 (48.08-2.25)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.06 (at 2.24Å)	Xtrriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, $R_{free}$	0.211 , 0.254 0.211 , 0.254	Depositor DCC
$R_{free}$ test set	1688 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	50.8	Xtrriage
Anisotropy	0.133	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 40.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4857	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	L	0.39	0/1620	0.61	1/2205 (0.0%)
2	H	0.42	1/1641 (0.1%)	0.58	0/2238
3	R	0.58	0/1584	0.63	0/2157
All	All	0.47	1/4845 (0.0%)	0.61	1/6600 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	96	CYS	CB-SG	-5.86	1.72	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	153	LEU	CB-CG-CD2	-5.45	101.73	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1586	0	1501	22	0
2	H	1603	0	1546	22	0
3	R	1540	0	1439	7	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	38	0	34	1	0
5	H	15	0	0	0	0
5	L	18	0	0	0	0
5	R	57	0	0	0	0
All	All	4857	0	4520	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:18:ARG:HG2	1:L:76:SER:HA	1.52	0.91
1:L:119:PRO:HD3	1:L:131:VAL:HG12	1.58	0.86
1:L:145:VAL:HG22	1:L:195:VAL:HG22	1.62	0.81
1:L:131:VAL:HG22	1:L:178:LEU:HB3	1.67	0.76
3:R:359:SER:HA	3:R:524:VAL:HG22	1.69	0.74
2:H:205:ILE:HD11	2:H:218:ASP:HB3	1.74	0.70
2:H:216:LYS:NZ	2:H:218:ASP:OD1	2.24	0.67
1:L:112:PRO:HD2	1:L:200:LEU:HD21	1.82	0.60
1:L:166:ASP:OD1	1:L:167:SER:N	2.36	0.58
1:L:78:LEU:HD11	1:L:103:LEU:HD21	1.86	0.56
1:L:178:LEU:HD13	1:L:180:LEU:HD23	1.87	0.56
2:H:180:LEU:HB2	2:H:186:TYR:CE1	2.39	0.56
2:H:207:ASN:HB3	2:H:216:LYS:NZ	2.20	0.56
1:L:178:LEU:CD1	1:L:180:LEU:HD23	2.36	0.56
2:H:207:ASN:HB3	2:H:216:LYS:HZ1	1.71	0.56
2:H:89:GLU:H	2:H:89:GLU:CD	2.08	0.55
1:L:197:HIS:HB3	1:L:200:LEU:HD23	1.89	0.55
2:H:9:GLY:HA2	2:H:18:LEU:HD21	1.87	0.55
2:H:196:SER:O	2:H:198:SER:N	2.38	0.54
2:H:198:SER:HA	2:H:201:THR:OG1	2.09	0.52
1:L:189:LYS:O	1:L:209:ASN:HA	2.10	0.51
2:H:175[A]:THR:HG22	2:H:190[A]:SER:OG	2.10	0.51
2:H:147:ALA:HB2	2:H:193:THR:HG22	1.93	0.51
2:H:12:VAL:HG21	2:H:86:LEU:HD13	1.91	0.50
1:L:123:GLN:O	1:L:126:SER:HB3	2.11	0.49
1:L:19:VAL:HG22	1:L:75:ILE:HB	1.94	0.49
1:L:131:VAL:CG2	1:L:178:LEU:HB3	2.41	0.49
1:L:166:ASP:O	1:L:170:SER:HA	2.13	0.49
2:H:62:ASP:OD1	2:H:65:LYS:NZ	2.44	0.49

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:51:ILE:HD13	2:H:72:ILE:HG12	1.95	0.48
1:L:160:GLU:HA	1:L:175:SER:O	2.14	0.48
3:R:359:SER:HA	3:R:524:VAL:CG2	2.42	0.47
3:R:354:ASN:O	3:R:398:ASP:HA	2.14	0.46
2:H:173:VAL:HG22	2:H:192:VAL:HG23	1.96	0.45
2:H:39:GLN:HE21	2:H:44:GLY:HA2	1.82	0.45
3:R:431:GLY:HA3	3:R:513:LEU:O	2.17	0.45
3:R:342:PHE:HB2	4:A:1:NAG:H82	1.99	0.43
2:H:65:LYS:HE3	2:H:65:LYS:HB2	1.70	0.43
1:L:78:LEU:HD23	1:L:79:GLN:N	2.33	0.43
2:H:48:VAL:HB	2:H:49:LEU:HD12	1.99	0.43
2:H:48:VAL:C	2:H:49:LEU:HD12	2.39	0.43
2:H:43:LYS:CD	2:H:44:GLY:H	2.32	0.43
2:H:104:LEU:O	2:H:104:LEU:HD23	2.19	0.43
3:R:393:THR:HG21	3:R:520:ALA:HB3	2.00	0.42
1:L:188:HIS:O	1:L:210:ARG:HD3	2.19	0.42
1:L:123:GLN:HG2	1:L:128:THR:O	2.19	0.41
3:R:484:GLU:HG2	3:R:490:PHE:HB2	2.02	0.41
1:L:131:VAL:HG23	1:L:147:TRP:CH2	2.56	0.41
1:L:49:TYR:HB2	2:H:109:TYR:CD2	2.56	0.41
1:L:185:TYR:HA	1:L:191:TYR:OH	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	210/214 (98%)	199 (95%)	10 (5%)	1 (0%)	29	29
2	H	213/234 (91%)	204 (96%)	8 (4%)	1 (0%)	29	29
3	R	195/204 (96%)	186 (95%)	9 (5%)	0	100	100
All	All	618/652 (95%)	589 (95%)	27 (4%)	2 (0%)	41	46

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	196	SER
1	L	154	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	174/187 (93%)	168 (97%)	6 (3%)	37	45
2	H	176/195 (90%)	173 (98%)	3 (2%)	60	71
3	R	163/176 (93%)	162 (99%)	1 (1%)	86	91
All	All	513/558 (92%)	503 (98%)	10 (2%)	55	66

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	7	SER
1	L	31	ARG
1	L	107	ARG
1	L	126	SER
1	L	141	ARG
1	L	210	ARG
2	H	43	LYS
2	H	89	GLU
2	H	206	CYS
3	R	373	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	174	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	A	1	3,4	14,14,15	0.47	0	17,19,21	0.61	0
4	NAG	A	2	4	14,14,15	0.44	0	17,19,21	0.50	0
4	FUC	A	3	4	10,10,11	1.04	0	14,14,16	0.99	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1	3,4	-	0/6/23/26	0/1/1/1
4	NAG	A	2	4	-	0/6/23/26	0/1/1/1
4	FUC	A	3	4	-	-	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	3	FUC	C1-O5-C5	2.11	117.56	112.78

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	L	211/214 (98%)	0.59	33 (15%) <b>2</b> <b>1</b>	46, 78, 130, 140	0
2	H	215/234 (91%)	0.45	25 (11%) <b>4</b> <b>4</b>	47, 72, 130, 147	0
3	R	196/204 (96%)	0.18	0 <b>100</b> <b>100</b>	33, 50, 83, 111	0
All	All	622/652 (95%)	0.42	58 (9%) <b>8</b> <b>9</b>	33, 69, 124, 147	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	202	SER	5.9
2	H	147	ALA	5.8
2	H	204	TYR	5.4
1	L	191	TYR	4.9
2	H	199	LEU	4.9
2	H	201	THR	4.7
1	L	183	ALA	4.6
1	L	126	SER	4.2
2	H	132	PHE	4.1
1	L	149	VAL	4.0
1	L	120	SER	4.0
2	H	221	VAL	3.9
1	L	151	ASN	3.7
1	L	203	PRO	3.7
1	L	116	ILE	3.7
1	L	124	LEU	3.6
1	L	200	LEU	3.5
2	H	217	VAL	3.4
1	L	207	SER	3.4
2	H	133	PRO	3.3
2	H	198	SER	3.2
2	H	134	LEU	3.2
2	H	136	PRO	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	L	180	LEU	3.1
1	L	204	VAL	3.0
1	L	125	LYS	3.0
1	L	114	VAL	2.9
1	L	210	ARG	2.9
1	L	187	LYS	2.8
2	H	146	ALA	2.8
2	H	168	ALA	2.8
2	H	200	GLY	2.7
2	H	173	VAL	2.7
2	H	148	LEU	2.7
1	L	184	ASP	2.6
2	H	191	VAL	2.6
2	H	208	VAL	2.6
2	H	223	PRO	2.6
2	H	197	SER	2.5
2	H	151	LEU	2.5
1	L	147	TRP	2.4
2	H	135	ALA	2.4
2	H	206	CYS	2.4
1	L	185	TYR	2.4
1	L	153	LEU	2.4
1	L	209	ASN	2.3
1	L	179	THR	2.3
1	L	188	HIS	2.3
1	L	119	PRO	2.3
1	L	150	ASP	2.2
2	H	164	TRP	2.2
1	L	115	PHE	2.2
1	L	152	ALA	2.2
1	L	145	VAL	2.1
2	H	169	LEU	2.1
1	L	211	GLY	2.1
1	L	194	GLU	2.0
1	L	208	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains


There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	FUC	A	3	10/11	0.73	0.17	119,122,124,126	0
4	NAG	A	2	14/15	0.79	0.20	111,115,120,123	0
4	NAG	A	1	14/15	0.84	0.14	69,85,104,111	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.