



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2023 – 12:24 AM EDT

PDB ID : 8DY5
Title : Crystal Structure of spFv CAT2200 LH in complex with IL-17A
Authors : Luo, J.; Boucher, L.E.
Deposited on : 2022-08-03
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	 5% 93%
1	B	251	 4% 90% 7%
2	C	122	 20% 73% 8% 1%
2	D	122	 24% 73% 11% 1%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5318 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called spFv CAT2200 LH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	244	Total 1781	C 1106	N 306	O 358	S 11	0	0	0
1	B	244	Total 1764	C 1096	N 303	O 354	S 11	0	0	0

- Molecule 2 is a protein called Interleukin-17A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	100	Total 733	C 461	N 132	O 134	S 6	0	0	0
2	D	103	Total 751	C 471	N 136	O 137	S 7	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	11	MET	-	initiating methionine	UNP Q16552
C	70	GLN	LYS	engineered mutation	UNP Q16552
C	106	SER	CYS	engineered mutation	UNP Q16552
C	132	GLN	ALA	engineered mutation	UNP Q16552
D	11	MET	-	initiating methionine	UNP Q16552
D	70	GLN	LYS	engineered mutation	UNP Q16552
D	106	SER	CYS	engineered mutation	UNP Q16552
D	132	GLN	ALA	engineered mutation	UNP Q16552

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Cl 1	0	0

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Ca 1 1	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	D	1	Total Mg 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	117	Total O 117 117	0	0
6	B	110	Total O 110 110	0	0
6	C	34	Total O 34 34	0	0
6	D	25	Total O 25 25	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: spFv CAT2200 LH




- Molecule 1: spFv CAT2200 LH



- Molecule 2: Interleukin-17A



- Molecule 2: Interleukin-17A



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	82.43Å 226.38Å 75.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.25 – 2.20 45.25 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (45.25-2.20) 99.4 (45.25-2.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.90 (at 2.20Å)	Xtrriage
Refinement program	PHENIX 1.18_3855	Depositor
R, R_{free}	0.188 , 0.232 0.188 , 0.232	Depositor DCC
R_{free} test set	1805 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	37.0	Xtrriage
Anisotropy	0.530	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5318	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.64 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.0945e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, CL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/1821	0.61	0/2475
1	B	0.40	0/1804	0.60	0/2456
2	C	0.38	0/751	0.58	0/1034
2	D	0.34	0/771	0.52	0/1063
All	All	0.40	0/5147	0.59	0/7028

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1781	0	1672	4	0
1	B	1764	0	1637	11	0
2	C	733	0	658	10	0
2	D	751	0	652	9	0
3	A	1	0	0	0	0
4	B	1	0	0	0	0
5	D	1	0	0	0	0
6	A	117	0	0	0	2
6	B	110	0	0	4	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	34	0	0	1	0
6	D	25	0	0	3	0
All	All	5318	0	4619	33	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:100:ARG:O	2:C:110:PHE:CB	1.70	1.40
2:C:101:ARG:HG2	2:C:101:ARG:HH11	1.45	0.80
2:D:64:SER:OG	6:D:301:HOH:O	2.00	0.78
2:C:101:ARG:HG2	2:C:101:ARG:NH1	1.99	0.76
2:D:56:ASN:OD1	2:D:61:ARG:NH2	2.21	0.73
1:B:172:GLU:OE1	6:B:401:HOH:O	2.10	0.69
1:A:209:MET:HE2	1:A:212:LEU:HD21	1.79	0.64
2:C:101:ARG:HH11	2:C:101:ARG:CG	2.13	0.62
1:B:1:ASN:ND2	6:B:403:HOH:O	2.35	0.58
2:D:36:ASN:N	6:D:303:HOH:O	2.38	0.56
1:B:33:TYR:OH	2:D:127:ILE:HD11	2.08	0.54
1:B:3:MET:SD	6:B:510:HOH:O	2.59	0.53
1:B:47:ILE:HD11	1:B:50:PHE:HB3	1.90	0.53
2:C:61:ARG:HD2	2:C:63:PRO:O	2.09	0.52
2:C:36:ASN:N	6:C:201:HOH:O	2.43	0.51
1:A:37:TYR:OH	1:A:231:THR:HB	2.12	0.49
2:C:37:PRO:HD2	2:C:53:LEU:HD13	1.94	0.49
1:B:217:THR:HG23	1:B:241:THR:HA	1.94	0.48
2:C:37:PRO:HD3	2:C:67:TRP:CG	2.49	0.47
1:A:4:LEU:HD11	1:A:93:THR:HG22	1.98	0.46
1:A:40:ARG:NH1	1:A:84:GLU:O	2.28	0.46
2:D:74:LEU:O	2:D:88:ASN:HA	2.15	0.46
2:D:78:ASN:HB2	6:D:319:HOH:O	2.15	0.46
2:D:100:ARG:O	2:D:111:ARG:N	2.45	0.46
1:B:4:LEU:HD11	1:B:93:THR:HG22	1.98	0.46
1:B:111:GLY:HA3	6:B:402:HOH:O	2.18	0.44
2:C:63:PRO:HG2	2:C:66:ILE:HD11	2.00	0.44
2:D:37:PRO:HG2	2:D:53:LEU:HD13	2.00	0.44
1:B:99:VAL:HB	1:B:173:TRP:CG	2.54	0.43
2:C:53:LEU:HD23	2:C:53:LEU:HA	1.81	0.42
1:B:223:ALA:HB1	1:B:231:THR:HG23	2.01	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:42:ASP:HA	2:D:44:TYR:CE1	2.54	0.41
1:B:37:TYR:CE1	1:B:47:ILE:HD12	2.55	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:486:HOH:O	6:A:486:HOH:O[3_554]	2.15	0.05
6:A:432:HOH:O	6:B:502:HOH:O[3_454]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	242/251 (96%)	230 (95%)	11 (4%)	1 (0%)	34	37
1	B	242/251 (96%)	230 (95%)	12 (5%)	0	100	100
2	C	96/122 (79%)	89 (93%)	7 (7%)	0	100	100
2	D	100/122 (82%)	95 (95%)	5 (5%)	0	100	100
All	All	680/746 (91%)	644 (95%)	35 (5%)	1 (0%)	51	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	116	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/203 (94%)	189 (100%)	1 (0%)	88	94
1	B	185/203 (91%)	184 (100%)	1 (0%)	88	94
2	C	75/117 (64%)	73 (97%)	2 (3%)	44	57
2	D	75/117 (64%)	75 (100%)	0	100	100
All	All	525/640 (82%)	521 (99%)	4 (1%)	81	90

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	VAL
1	B	110	LEU
2	C	101	ARG
2	C	112	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	244/251 (97%)	0.02	13 (5%) 26 25	26, 35, 61, 89	0
1	B	244/251 (97%)	-0.07	9 (3%) 41 39	27, 38, 62, 83	0
2	C	100/122 (81%)	0.86	24 (24%) 0 0	28, 46, 101, 105	0
2	D	103/122 (84%)	1.11	29 (28%) 0 0	29, 53, 103, 108	0
All	All	691/746 (92%)	0.28	75 (10%) 5 5	26, 37, 89, 108	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	21	THR	9.1
1	A	115	GLY	8.7
1	A	116	SER	8.2
1	B	113	SER	7.6
2	D	105	HIS	6.6
1	A	114	GLY	6.6
1	B	112	GLY	6.5
2	D	21	THR	6.2
2	C	23	MET	6.1
2	D	23	MET	6.1
1	A	117	GLY	6.1
1	B	114	GLY	5.5
2	D	26	LEU	5.4
2	C	104	PRO	5.3
1	B	122	CYS	5.3
1	A	123	GLY	5.3
1	B	115	GLY	5.2
1	A	113	SER	5.1
2	D	110	PHE	4.9
1	B	125	GLY	4.8
2	D	24	VAL	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	125	GLY	4.5
1	A	112	GLY	4.2
1	A	1	ASN	4.0
2	D	107	PRO	4.0
1	B	116	SER	4.0
2	C	110	PHE	3.9
2	C	24	VAL	3.9
2	D	64	SER	3.9
2	C	37	PRO	3.9
2	C	108	ASN	3.8
2	D	112	LEU	3.8
2	D	99	LEU	3.8
1	A	111	GLY	3.7
2	C	59	PRO	3.6
2	D	25	ASN	3.6
2	D	66	ILE	3.5
2	D	59	PRO	3.5
2	D	22	VAL	3.5
2	D	36	ASN	3.5
2	C	98	VAL	3.4
2	C	105	HIS	3.4
2	C	58	ASP	3.4
2	C	36	ASN	3.3
2	D	103	PRO	3.2
2	D	108	ASN	3.2
2	C	109	SER	3.1
1	B	126	GLY	3.1
2	D	101	ARG	3.0
2	D	63	PRO	3.0
2	D	98	VAL	2.9
2	D	102	GLU	2.9
2	C	107	PRO	2.8
2	D	111	ARG	2.8
2	C	102	GLU	2.8
2	C	62	TYR	2.8
2	C	26	LEU	2.7
2	D	62	TYR	2.7
2	C	103	PRO	2.6
2	C	25	ASN	2.5
2	D	97	LEU	2.5
2	D	39	ARG	2.4
2	D	96	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	244	SER	2.3
2	D	104	PRO	2.2
1	A	126	GLY	2.2
2	C	101	ARG	2.2
2	C	111	ARG	2.2
1	A	122	CYS	2.2
2	D	115	ILE	2.2
2	C	39	ARG	2.2
2	D	109	SER	2.1
2	C	22	VAL	2.1
2	C	40	SER	2.1
1	A	61	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CA	B	301	1/1	0.91	0.17	69,69,69,69	0
3	CL	A	301	1/1	0.94	0.06	68,68,68,68	0
5	MG	D	201	1/1	0.97	0.38	47,47,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.