



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 2, 2024 – 12:40 PM EST

PDB ID : 8DYE
Title : Crystal structure of human SDHA-SDHAF4 assembly intermediate
Authors : Sharma, P.; Maklashina, E.; Cecchini, G.; Iverson, T.M.
Deposited on : 2022-08-04
Resolution : 1.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2021 (1.46-1.42)
Clashscore	141614	2086 (1.46-1.42)
Ramachandran outliers	138981	2047 (1.46-1.42)
Sidechain outliers	138945	2047 (1.46-1.42)
RSRZ outliers	127900	1993 (1.46-1.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	611	 4% 91% 9%
2	B	98	 12% 36% 8% 56%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	A	703	-	-	X	-
5	EDO	A	706	-	-	X	-
6	GOL	A	711	-	-	X	-
6	GOL	A	716	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 9601 atoms, of which 3865 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	611	8469	3022	3636	858	919	34	1	41	0

- Molecule 2 is a protein called Succinate dehydrogenase assembly factor 4, mitochondrial.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
2	B	43	564	211	227	61	64	1	0	0	0

There are 23 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	11	MET	-	initiating methionine	UNP Q5VUM1
B	12	ARG	-	expression tag	UNP Q5VUM1
B	13	SER	-	expression tag	UNP Q5VUM1
B	14	HIS	-	expression tag	UNP Q5VUM1
B	15	HIS	-	expression tag	UNP Q5VUM1
B	16	HIS	-	expression tag	UNP Q5VUM1
B	17	HIS	-	expression tag	UNP Q5VUM1
B	18	HIS	-	expression tag	UNP Q5VUM1
B	19	HIS	-	expression tag	UNP Q5VUM1
B	20	GLU	-	expression tag	UNP Q5VUM1
B	21	ASN	-	expression tag	UNP Q5VUM1
B	22	LEU	-	expression tag	UNP Q5VUM1
B	23	TYR	-	expression tag	UNP Q5VUM1
B	24	PHE	-	expression tag	UNP Q5VUM1
B	25	GLN	-	expression tag	UNP Q5VUM1
B	26	GLY	-	expression tag	UNP Q5VUM1
B	27	ILE	-	expression tag	UNP Q5VUM1
B	28	ASP	-	expression tag	UNP Q5VUM1
B	29	PRO	-	expression tag	UNP Q5VUM1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	30	PHE	-	expression tag	UNP Q5VUM1
B	31	THR	-	expression tag	UNP Q5VUM1
B	32	GLY	-	expression tag	UNP Q5VUM1
B	33	SER	-	expression tag	UNP Q5VUM1

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	53	27	9	15	2	0	0

- Molecule 4 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	A	1	9	3	2	4	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	4	2	2	0	0
5	A	1	4	2	2	0	0
5	A	1	4	2	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0

- Molecule 7 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	2	Total Na 2 2	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	392	Total O 392 392	0	0
8	B	30	Total O 30 30	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial



- Molecule 2: Succinate dehydrogenase assembly factor 4, mitochondrial



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	139.05Å 69.30Å 72.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.57 – 1.44 28.57 – 1.44	Depositor EDS
% Data completeness (in resolution range)	99.4 (28.57-1.44) 99.6 (28.57-1.44)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.44Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.155 , 0.173 0.155 , 0.172	Depositor DCC
R_{free} test set	6232 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	18.5	Xtrriage
Anisotropy	0.164	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 53.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.013 for -h,l,k	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9601	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MLI, NA, EDO, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/5109	0.68	1/6914 (0.0%)
2	B	0.42	0/347	0.66	0/468
All	All	0.42	0/5456	0.68	1/7382 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	459	ASP	CB-CG-OD2	-5.10	113.71	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4833	3636	4551	49	0
2	B	337	227	310	7	0
3	A	53	0	30	0	0
4	A	7	2	2	0	0
5	A	64	0	96	23	0
6	A	18	0	24	15	0
7	A	2	0	0	0	0
8	A	392	0	0	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	B	30	0	0	0	0
All	All	5736	3865	5013	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:ARG:HH22	6:A:711:GOL:H31	1.38	0.86
1:A:232:ARG:HD3	1:A:482:PRO:HB2	1.61	0.81
1:A:277[A]:THR:HG23	1:A:287:CYS:SG	2.22	0.79
1:A:440:GLU:OE1	5:A:703:EDO:H11	1.83	0.78
1:A:544:GLY:O	1:A:547:LYS:HG2	1.91	0.70
5:A:713:EDO:H12	8:A:967:HOH:O	1.93	0.69
5:A:706:EDO:H22	6:A:716:GOL:O2	1.93	0.68
1:A:374:GLU:HG2	8:A:1117:HOH:O	1.95	0.67
1:A:508:THR:HG21	1:A:564:GLU:HG3	1.76	0.66
1:A:607:SER:OG	1:A:608:LYS:HE3	1.96	0.65
1:A:242:GLY:CA	5:A:718:EDO:H21	2.27	0.65
1:A:75:ARG:HH22	6:A:711:GOL:C3	2.09	0.62
1:A:526:PHE:CE1	1:A:595:GLU:HG3	2.35	0.61
1:A:75:ARG:HH12	6:A:711:GOL:C3	2.14	0.60
1:A:75:ARG:HH12	6:A:711:GOL:H32	1.66	0.60
2:B:66:LEU:HA	2:B:102:LYS:NZ	2.16	0.60
1:A:632[B]:VAL:HG23	5:A:714:EDO:H11	1.84	0.59
1:A:75:ARG:NH2	6:A:711:GOL:H31	2.14	0.59
5:A:706:EDO:H12	6:A:716:GOL:C3	2.33	0.59
2:B:66:LEU:CB	2:B:70:PRO:HA	2.33	0.59
2:B:66:LEU:O	2:B:70:PRO:HB3	2.04	0.58
1:A:258:GLY:H	5:A:703:EDO:H21	1.70	0.55
2:B:69:GLU:O	2:B:102:LYS:HE2	2.07	0.54
1:A:444:ALA:N	1:A:445:SER:HA	2.22	0.54
5:A:714:EDO:H12	8:A:918:HOH:O	2.08	0.53
1:A:242:GLY:HA2	5:A:718:EDO:H21	1.91	0.53
1:A:314:GLU:OE1	1:A:379:ARG:HD3	2.09	0.52
1:A:490:GLU:O	1:A:494:MET:HG3	2.09	0.52
5:A:706:EDO:H12	6:A:716:GOL:H32	1.91	0.52
1:A:99:HIS:CD2	5:A:703:EDO:H12	2.45	0.52
1:A:551:THR:OG1	1:A:554:ARG:NH1	2.43	0.51
1:A:61:GLU:OE2	1:A:481:VAL:HB	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:706:EDO:H12	6:A:716:GOL:H31	1.92	0.50
1:A:154:GLU:OE2	1:A:162:ARG:NH2	2.44	0.50
1:A:277[A]:THR:HG22	8:A:1008:HOH:O	2.11	0.50
1:A:277[B]:THR:OG1	1:A:574:CYS:SG	2.66	0.49
1:A:475[B]:CYS:HB2	1:A:479:ASP:OD2	2.13	0.49
2:B:66:LEU:HA	2:B:102:LYS:HZ1	1.77	0.49
1:A:260:GLY:HA3	1:A:266[B]:CYS:SG	2.52	0.48
5:A:706:EDO:H22	6:A:716:GOL:C3	2.42	0.48
1:A:472:GLU:OE2	6:A:711:GOL:H12	2.13	0.48
1:A:374:GLU:HA	1:A:374:GLU:OE1	2.13	0.48
1:A:280:ILE:HG13	1:A:413:ILE:HG12	1.95	0.47
1:A:258:GLY:H	5:A:703:EDO:C2	2.27	0.47
1:A:258:GLY:N	5:A:703:EDO:H21	2.31	0.46
1:A:251:ASN:OD1	5:A:721:EDO:H21	2.16	0.45
5:A:709:EDO:H11	8:A:1087:HOH:O	2.17	0.44
1:A:250:LYS:HE3	1:A:475[A]:CYS:SG	2.58	0.44
1:A:496[A]:LEU:HD21	5:A:718:EDO:O2	2.18	0.44
2:B:94:THR:HA	2:B:98:ASP:O	2.18	0.43
5:A:706:EDO:C1	6:A:716:GOL:H32	2.48	0.43
1:A:277[B]:THR:H	1:A:277[B]:THR:HG23	1.47	0.43
5:A:706:EDO:H22	6:A:716:GOL:H32	2.00	0.43
1:A:78:PHE:CZ	1:A:207:ARG:HD3	2.54	0.42
1:A:116[A]:GLU:HG3	8:A:1048:HOH:O	2.19	0.42
1:A:116[B]:GLU:HB3	8:A:822:HOH:O	2.18	0.42
1:A:632[A]:VAL:HG13	5:A:714:EDO:H11	2.03	0.41
2:B:100:GLU:HA	2:B:104:ARG:O	2.20	0.41
1:A:375:GLN:HG3	1:A:379:ARG:HG3	2.03	0.41
1:A:55:TYR:O	1:A:57:VAL:HG23	2.20	0.41
1:A:248:ARG:HD3	1:A:484:ILE:HG12	2.02	0.41
1:A:491:GLU:HA	1:A:494:MET:HE2	2.03	0.41
1:A:250:LYS:HG2	5:A:721:EDO:H22	2.03	0.40
1:A:282:ARG:CD	6:A:716:GOL:H32	2.50	0.40
5:A:706:EDO:H22	6:A:716:GOL:C2	2.51	0.40
1:A:93:LEU:HD21	1:A:271:THR:HG21	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	650/611 (106%)	637 (98%)	13 (2%)	0	100	100
2	B	41/98 (42%)	39 (95%)	2 (5%)	0	100	100
All	All	691/709 (98%)	676 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	531/495 (107%)	528 (99%)	3 (1%)	86	68
2	B	34/87 (39%)	34 (100%)	0	100	100
All	All	565/582 (97%)	562 (100%)	3 (0%)	86	74

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	456	SER
1	A	459	ASP
1	A	547	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 2 are monoatomic - leaving 21 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	A	704	-	3,3,3	0.53	0	2,2,2	0.12	0
5	EDO	A	712	-	3,3,3	0.47	0	2,2,2	0.34	0
6	GOL	A	716	-	5,5,5	1.09	0	5,5,5	1.04	0
5	EDO	A	710	-	3,3,3	0.45	0	2,2,2	0.26	0
5	EDO	A	719	-	3,3,3	0.47	0	2,2,2	0.31	0
5	EDO	A	708	-	3,3,3	0.44	0	2,2,2	0.46	0
5	EDO	A	717	-	3,3,3	0.45	0	2,2,2	0.27	0
5	EDO	A	703	-	3,3,3	0.45	0	2,2,2	0.72	0
5	EDO	A	718	-	3,3,3	0.41	0	2,2,2	0.50	0
6	GOL	A	707	-	5,5,5	1.07	0	5,5,5	1.22	0
5	EDO	A	705	-	3,3,3	0.45	0	2,2,2	0.44	0
6	GOL	A	711	-	5,5,5	0.90	0	5,5,5	1.41	1 (20%)
5	EDO	A	714	-	3,3,3	0.42	0	2,2,2	0.46	0
5	EDO	A	706	-	3,3,3	0.53	0	2,2,2	0.26	0
5	EDO	A	713	-	3,3,3	0.44	0	2,2,2	0.33	0
5	EDO	A	715	-	3,3,3	0.50	0	2,2,2	0.37	0
5	EDO	A	709	-	3,3,3	0.50	0	2,2,2	0.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MLI	A	702	-	6,6,6	1.56	1 (16%)	7,7,7	1.09	0
5	EDO	A	721	-	3,3,3	0.47	0	2,2,2	0.34	0
3	FAD	A	701	7,1	53,58,58	0.61	1 (1%)	68,89,89	0.68	1 (1%)
5	EDO	A	720	-	3,3,3	0.44	0	2,2,2	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	704	-	-	0/1/1/1	-
5	EDO	A	712	-	-	0/1/1/1	-
6	GOL	A	716	-	-	4/4/4/4	-
5	EDO	A	710	-	-	0/1/1/1	-
5	EDO	A	719	-	-	1/1/1/1	-
5	EDO	A	708	-	-	0/1/1/1	-
5	EDO	A	717	-	-	1/1/1/1	-
5	EDO	A	703	-	-	1/1/1/1	-
5	EDO	A	718	-	-	0/1/1/1	-
6	GOL	A	707	-	-	4/4/4/4	-
5	EDO	A	705	-	-	0/1/1/1	-
6	GOL	A	711	-	-	4/4/4/4	-
5	EDO	A	714	-	-	0/1/1/1	-
5	EDO	A	706	-	-	1/1/1/1	-
5	EDO	A	713	-	-	1/1/1/1	-
5	EDO	A	715	-	-	1/1/1/1	-
5	EDO	A	709	-	-	0/1/1/1	-
4	MLI	A	702	-	-	4/4/4/4	-
5	EDO	A	721	-	-	1/1/1/1	-
3	FAD	A	701	7,1	-	6/30/50/50	0/6/6/6
5	EDO	A	720	-	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	702	MLI	C1-C3	2.62	1.55	1.51
3	A	701	FAD	P-O2P	-2.11	1.45	1.55

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	FAD	C5A-C6A-N6A	2.50	124.15	120.35
6	A	711	GOL	C3-C2-C1	-2.48	102.06	111.70

There are no chirality outliers.

All (30) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	701	FAD	N10-C1'-C2'-O2'
3	A	701	FAD	N10-C1'-C2'-C3'
6	A	707	GOL	O1-C1-C2-O2
6	A	707	GOL	O1-C1-C2-C3
6	A	711	GOL	C1-C2-C3-O3
6	A	716	GOL	C1-C2-C3-O3
6	A	716	GOL	O2-C2-C3-O3
5	A	706	EDO	O1-C1-C2-O2
6	A	707	GOL	C1-C2-C3-O3
6	A	711	GOL	O1-C1-C2-C3
6	A	716	GOL	O1-C1-C2-C3
6	A	711	GOL	O2-C2-C3-O3
4	A	702	MLI	C2-C1-C3-O9
6	A	707	GOL	O2-C2-C3-O3
6	A	711	GOL	O1-C1-C2-O2
4	A	702	MLI	C2-C1-C3-O8
3	A	701	FAD	PA-O3P-P-O5'
6	A	716	GOL	O1-C1-C2-O2
4	A	702	MLI	C3-C1-C2-O6
5	A	719	EDO	O1-C1-C2-O2
5	A	717	EDO	O1-C1-C2-O2
4	A	702	MLI	C3-C1-C2-O7
3	A	701	FAD	P-O3P-PA-O1A
3	A	701	FAD	P-O3P-PA-O2A
5	A	713	EDO	O1-C1-C2-O2
5	A	703	EDO	O1-C1-C2-O2
5	A	715	EDO	O1-C1-C2-O2
3	A	701	FAD	O4B-C4B-C5B-O5B
5	A	720	EDO	O1-C1-C2-O2
5	A	721	EDO	O1-C1-C2-O2


There are no ring outliers.

9 monomers are involved in 30 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	716	GOL	9	0
5	A	703	EDO	5	0
5	A	718	EDO	3	0
6	A	711	GOL	6	0
5	A	714	EDO	3	0
5	A	706	EDO	8	0
5	A	713	EDO	1	0
5	A	709	EDO	1	0
5	A	721	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	611/611 (100%)	0.11	25 (4%) 37 38	13, 23, 44, 74	0
2	B	43/98 (43%)	0.99	12 (27%) 0 0	19, 28, 67, 76	0
All	All	654/709 (92%)	0.17	37 (5%) 23 23	13, 23, 46, 76	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	66	LEU	11.2
1	A	664	TYR	8.4
2	B	70	PRO	8.4
2	B	67	GLU	6.7
1	A	54	GLN	5.5
1	A	612	GLY	5.4
1	A	55	TYR	5.2
1	A	277[A]	THR	4.7
1	A	661	ILE	4.6
2	B	69	GLU	4.5
1	A	614	GLN	4.4
2	B	75	PRO	4.3
1	A	396	THR	3.9
1	A	374	GLU	3.8
1	A	397	LYS	3.6
2	B	71	LEU	3.5
1	A	376	LEU	3.5
2	B	81	VAL	3.4
1	A	366[A]	LEU	3.3
1	A	56	PRO	3.0
1	A	371	LEU	2.9
2	B	80	PRO	2.9
2	B	68	LYS	2.8
1	A	663[A]	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	76	ASP	2.6
1	A	373	PRO	2.5
1	A	551	THR	2.5
1	A	552	PHE	2.5
1	A	607	SER	2.4
2	B	74	PHE	2.4
1	A	359	PRO	2.4
1	A	234[A]	VAL	2.4
1	A	333	VAL	2.4
1	A	372	PRO	2.4
2	B	73	LYS	2.4
1	A	337	LEU	2.2
1	A	332	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	A	721	4/4	0.71	0.19	41,44,45,49	0
6	GOL	A	707	6/6	0.74	0.28	35,41,46,47	0
5	EDO	A	706	4/4	0.76	0.22	38,41,43,49	0
5	EDO	A	709	4/4	0.77	0.27	39,41,49,52	0
6	GOL	A	711	6/6	0.78	0.22	30,36,47,49	0
5	EDO	A	719	4/4	0.82	0.13	48,52,54,56	0
5	EDO	A	718	4/4	0.82	0.28	44,45,47,47	0
5	EDO	A	708	4/4	0.86	0.15	28,32,35,47	0
5	EDO	A	704	4/4	0.87	0.12	29,30,32,39	0
5	EDO	A	717	4/4	0.88	0.18	36,38,40,53	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	EDO	A	712	4/4	0.88	0.17	39,40,41,43	0
5	EDO	A	705	4/4	0.89	0.15	32,32,43,48	0
4	MLI	A	702	7/7	0.90	0.15	29,32,36,39	0
6	GOL	A	716	6/6	0.91	0.19	20,31,35,42	6
5	EDO	A	710	4/4	0.92	0.12	32,33,36,39	0
5	EDO	A	720	4/4	0.92	0.20	35,37,53,55	0
5	EDO	A	703	4/4	0.94	0.22	19,22,25,38	0
5	EDO	A	713	4/4	0.94	0.12	46,50,51,54	0
5	EDO	A	715	4/4	0.96	0.07	32,33,39,44	0
5	EDO	A	714	4/4	0.97	0.06	31,35,36,44	0
3	FAD	A	701	53/53	0.98	0.09	11,14,17,21	0
7	NA	A	723	1/1	0.99	0.08	19,19,19,19	0
7	NA	A	722	1/1	1.00	0.10	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.