

wwPDB X-ray Structure Validation Summary Report (i)

Oct 15, 2023 – 09:24 AM EDT

PDB ID : 8DYQ

Title : Crystal structure of Neisseria gonorrhoeae carbonic anhydrase with Acetazo-

lamide

Authors: Marapaka, A.K.; Das, C.; Flaherty, D.P.; Yadav, R.

Deposited on : 2022-08-04

Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Medit	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain		
1	A	243	75%	16%	9%
1	С	243	70%	18%	• 9%
1	E	243	69%	20%	• 9%
1	G	243	72%	19%	• 9%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Carbonic anhydrase.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	Λ	222	Total	С	N	О	S	0	1	0
1	A	222	1746	1105	306	329	6	0	1	U
1	С	222	Total	С	N	О	S	0	0	0
1		222	1733	1097	306	324	6	U	U	
1	E	222	Total	С	N	О	S	0	1	0
1	12	222	1741	1102	306	327	6	0	1	
1	С	222	Total	С	N	О	S	0	1	0
1	G	$G \mid 222 \mid$	1756	1110	309	331	6		1	

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-16	HIS	-	expression tag	UNP Q50940
A	-15	HIS	-	expression tag	UNP Q50940
A	-14	HIS	-	expression tag	UNP Q50940
A	-13	HIS	-	expression tag	UNP Q50940
A	-12	HIS	-	expression tag	UNP Q50940
A	-11	HIS	-	expression tag	UNP Q50940
A	-10	ASP	-	expression tag	UNP Q50940
A	-9	SER	-	expression tag	UNP Q50940
A	-8	GLY	-	expression tag	UNP Q50940
A	-7	LEU	-	expression tag	UNP Q50940
A	-6	VAL	-	expression tag	UNP Q50940
A	-5	PRO	-	expression tag	UNP Q50940
A	-4	ARG	-	expression tag	UNP Q50940
A	-3	GLY	-	expression tag	UNP Q50940
A	-2	SER	-	expression tag	UNP Q50940
A	-1	HIS	-	expression tag	UNP Q50940
A	0	MET	-	expression tag	UNP Q50940
С	-16	HIS	-	expression tag	UNP Q50940
С	-15	HIS		expression tag	UNP Q50940
С	-14	HIS	-	expression tag	UNP Q50940
С	-13	HIS	-	expression tag	UNP Q50940



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled	Actual	Comment	Reference
С	-12	HIS	-	expression tag	UNP Q50940
С	-11	HIS	-	expression tag	UNP Q50940
С	-10	ASP	-	expression tag	UNP Q50940
С	-9	SER	-	expression tag	UNP Q50940
С	-8	GLY	-	expression tag	UNP Q50940
С	-7	LEU	_	expression tag	UNP Q50940
С	-6	VAL	-	expression tag	UNP Q50940
С	-5	PRO	-	expression tag	UNP Q50940
С	-4	ARG	-	expression tag	UNP Q50940
С	-3	GLY	-	expression tag	UNP Q50940
С	-2	SER	-	expression tag	UNP Q50940
С	-1	HIS	-	expression tag	UNP Q50940
С	0	MET	-	expression tag	UNP Q50940
Е	-16	HIS	-	expression tag	UNP Q50940
Е	-15	HIS	-	expression tag	UNP Q50940
Е	-14	HIS	-	expression tag	UNP Q50940
Е	-13	HIS	-	expression tag	UNP Q50940
Е	-12	HIS	-	expression tag	UNP Q50940
Е	-11	HIS	-	expression tag	UNP Q50940
Е	-10	ASP	-	expression tag	UNP Q50940
Е	-9	SER	-	expression tag	UNP Q50940
Е	-8	GLY	_	expression tag	UNP Q50940
Е	-7	LEU	-	expression tag	UNP Q50940
Е	-6	VAL	-	expression tag	UNP Q50940
Е	-5	PRO	_	expression tag	UNP Q50940
Е	-4	ARG	-	expression tag	UNP Q50940
Е	-3	GLY	-	expression tag	UNP Q50940
Е	-2	SER	-	expression tag	UNP Q50940
Е	-1	HIS	-	expression tag	UNP Q50940
Е	0	MET	-	expression tag	UNP Q50940
G	-16	HIS	-	expression tag	UNP Q50940
G	-15	HIS	-	expression tag	UNP Q50940
G	-14	HIS	-	expression tag	UNP Q50940
G	-13	HIS	-	expression tag	UNP Q50940
G	-12	HIS	-	expression tag	UNP Q50940
G	-11	HIS	-	expression tag	UNP Q50940
G	-10	ASP	-	expression tag	UNP Q50940
G	-9	SER	-	expression tag	UNP Q50940
G	-8	GLY	-	expression tag	UNP Q50940
G	-7	LEU	-	expression tag	UNP Q50940
G	-6	VAL	-	expression tag	UNP Q50940
G	-5	PRO	-	expression tag	UNP Q50940



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	-4	ARG	-	expression tag	UNP Q50940
G	-3	GLY	-	expression tag	UNP Q50940
G	-2	SER	-	expression tag	UNP Q50940
G	-1	HIS	-	expression tag	UNP Q50940
G	0	MET	-	expression tag	UNP Q50940

• Molecule 2 is 5-ACETAMIDO-1,3,4-THIADIAZOLE-2-SULFONAMIDE (three-letter code: AZM) (formula: $C_4H_6N_4O_3S_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
2	Δ	1	Total	С	N	О	S	0	0
	Λ	1	13	4	4	3	2		0
2	С	1	Total	С	N	О	S	0	0
	C	1	13	4	4	3	2		
2	E	1	Total	С	N	О	S	0	0
	ינו	1	13	4	4	3	2	0	U
9	С	1	Total	С	N	О	S	0	0
	G		13	4	4	3	2	0	U

• Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	0
3	С	1	Total Zn 1 1	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total Zn 1 1	0	0
3	G	1	Total Zn 1 1	0	0

 \bullet Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	С	1	Total O S 5 4 1	0	0
4	С	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0
4	G	1	Total O S 5 4 1	0	0

• Molecule 5 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	10	Total O 10 10	0	0
5	С	11	Total O 11 11	0	0
5	Е	9	Total O 9 9	0	0
5	G	7	Total O 7 7	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Carbonic anhydrase









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	71.97Å 122.59Å 78.35Å	Depositor
a, b, c, α , β , γ	90.00° 117.21° 90.00°	Depositor
Resolution (Å)	46.02 - 2.15	Depositor
resolution (A)	46.02 - 2.15	EDS
% Data completeness	91.6 (46.02-2.15)	Depositor
(in resolution range)	91.1 (46.02-2.15)	EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.18 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.8.0352	Depositor
R, R_{free}	0.259 , 0.314	Depositor
it, it free	0.263 , 0.312	DCC
R_{free} test set	2850 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.098	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.36, 10.2	EDS
L-test for twinning ²	$< L >=0.39, < L^2>=0.22$	Xtriage
Estimated twinning fraction	0.377 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7109	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: AZM, SO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles		
IVIOI	Chain	RMSZ # Z > 5		RMSZ	# Z > 5	
1	A	0.46	0/1795	0.80	0/2445	
1	С	0.42	0/1779	0.82	$1/2423 \ (0.0\%)$	
1	Е	0.45	0/1790	0.85	0/2439	
1	G	0.42	0/1805	0.81	$1/2457 \ (0.0\%)$	
All	All	0.43	0/7169	0.82	2/9764 (0.0%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	С	0	2
1	Е	0	3
1	G	0	1
All	All	0	7

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mo	ol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1		G	41	GLU	CB-CA-C	-5.78	98.84	110.40
1		С	156	PRO	N-CD-CG	-5.64	94.75	103.20

There are no chirality outliers.

5 of 7 planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	84	ARG	Sidechain
1	С	26	ARG	Sidechain
1	С	84	ARG	Sidechain
1	Е	166	ARG	Sidechain
1	Е	84	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1746	0	1701	25	0
1	С	1733	0	1683	32	0
1	Е	1741	0	1691	33	0
1	G	1756	0	1716	22	0
2	A	13	0	5	0	0
2	С	13	0	6	2	0
2	Е	13	0	6	2	0
2	G	13	0	6	0	0
3	A	1	0	0	0	0
3	С	1	0	0	0	0
3	Е	1	0	0	0	0
3	G	1	0	0	0	0
4	A	5	0	0	0	0
4	С	10	0	0	1	0
4	Е	20	0	0	1	0
4	G	5	0	0	0	0
5	A	10	0	0	2	0
5	С	11	0	0	0	0
5	Е	9	0	0	0	0
5	G	7	0	0	0	0
All	All	7109	0	6814	109	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 109 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:50:ILE:HG23	1:A:79:LEU:HD21	1.60	0.84
1:C:34:GLN:HE22	1:C:181:CYS:HB3	1.51	0.76
1:C:34:GLN:HE21	1:C:214:ARG:HH12	1.35	0.72
1:C:132:LYS:NZ	4:C:303:SO4:O4	2.24	0.70
1:A:54:TYR:CE2	1:A:163:LEU:HD11	2.27	0.70

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	221/243 (91%)	212 (96%)	8 (4%)	1 (0%)	29	22
1	\mathbf{C}	220/243 (90%)	209 (95%)	9 (4%)	2 (1%)	17	11
1	E	221/243 (91%)	208 (94%)	12 (5%)	1 (0%)	29	22
1	G	221/243 (91%)	207 (94%)	12 (5%)	2 (1%)	17	11
All	All	883/972 (91%)	836 (95%)	41 (5%)	6 (1%)	22	15

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	156	PRO
1	A	75	GLY
1	С	155	GLN
1	G	119	LYS
1	G	12	HIS

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Chain Analysed Rotameric Outliers		Percentiles		
1	A	193/213 (91%)	187 (97%)	6 (3%)	40 39	
1	С	189/213 (89%)	178 (94%)	11 (6%)	20 15	
1	E	191/213 (90%)	179 (94%)	12 (6%)	18 13	
1	G	195/213 (92%)	184 (94%)	11 (6%)	21 16	
All	All	768/852 (90%)	728 (95%)	40 (5%)	23 19	

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Е	166	ARG
1	G	141	TRP
1	G	6	HIS
1	G	82	ASN
1	G	160	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	33	ASN
1	Е	66	HIS
1	G	82	ASN
1	G	66	HIS
1	G	69	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Chain Res		Dag	Link	В	ond leng	gths Bond angles			gles	
MIOI	Type	Chain	Res	Lilik	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	AZM	Е	301	3	8,13,13	3.05	4 (50%)	9,19,19	5.56	6 (66%)
2	AZM	G	301	3	8,13,13	3.63	5 (62%)	9,19,19	5.13	7 (77%)
2	AZM	С	301	3	8,13,13	3.74	4 (50%)	9,19,19	7.84	6 (66%)
4	SO4	С	303	-	4,4,4	0.26	0	6,6,6	0.10	0
4	SO4	С	304	-	4,4,4	0.40	0	6,6,6	0.16	0
4	SO4	Е	303	-	4,4,4	0.28	0	6,6,6	0.07	0
4	SO4	Е	304	-	4,4,4	0.35	0	6,6,6	0.21	0
4	SO4	Е	305	-	4,4,4	0.32	0	6,6,6	0.25	0
2	AZM	A	301	3	8,13,13	3.51	6 (75%)	9,19,19	5.06	4 (44%)
4	SO4	Е	306	-	4,4,4	0.37	0	6,6,6	0.17	0
4	SO4	G	303	-	4,4,4	0.42	0	6,6,6	0.28	0
4	SO4	A	303	-	4,4,4	0.44	0	6,6,6	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AZM	Е	301	3	-	0/2/10/10	0/1/1/1
2	AZM	A	301	3	-	0/2/10/10	0/1/1/1
2	AZM	G	301	3	-	0/2/10/10	0/1/1/1
2	AZM	С	301	3	-	0/2/10/10	0/1/1/1

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\text{\AA})$
2	G	301	AZM	S1-N1	6.16	1.72	1.60
2	С	301	AZM	C3-N4	5.99	1.47	1.36
2	G	301	AZM	C3-N4	5.85	1.46	1.36



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
2	С	301	AZM	S1-N1	5.78	1.71	1.60
2	A	301	AZM	C3-N4	5.76	1.46	1.36

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	С	301	AZM	O2-S1-O1	-15.07	93.99	118.76
2	С	301	AZM	O1-S1-N1	13.06	126.72	107.36
2	Е	301	AZM	O2-S1-O1	-13.01	97.38	118.76
2	A	301	AZM	O2-S1-O1	-12.98	97.43	118.76
2	С	301	AZM	O2-S1-C1	-11.50	95.08	108.59

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	Е	301	AZM	2	0
2	С	301	AZM	2	0
4	С	303	SO4	1	0
4	Е	305	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around AZM C 301:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around AZM E 301:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

