



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 20, 2023 – 11:38 PM JST

PDB ID : 7E0T
Title : Crystal Structure of Human Indoleamine 2,3-dioxygenase 1 (hIDO1) Complexed with (1R,2S)-2-(((5-bromo-1H-indazol-4-yl)amino)methyl)Cyclohexan-1-ol (36)
Authors : Li, G.-B.; Ning, X.-L.
Deposited on : 2021-01-28
Resolution : 2.14 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.14 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	 77% 15% • 5% 7%
1	B	392	 78% 15% • 6% 6%



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	Br	C	N	O		
3	A	1	Total	Br	C	N	O	0	0
			19	1	14	3	1		
3	B	1	Total	Br	C	N	O	0	0
			19	1	14	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	21	Total	O	0	0
			21	21		
4	B	17	Total	O	0	0
			17	17		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Indoleamine 2,3-dioxygenase 1



- Molecule 1: Indoleamine 2,3-dioxygenase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.90Å 97.11Å 131.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.73 – 2.14 19.73 – 2.14	Depositor EDS
% Data completeness (in resolution range)	90.3 (19.73-2.14) 90.3 (19.73-2.14)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 2.13Å)	Xtrriage
Refinement program	PHENIX 1.10.1_2155	Depositor
R, R_{free}	0.213 , 0.272 0.217 , 0.272	Depositor DCC
R_{free} test set	1998 reflections (3.59%)	wwPDB-VP
Wilson B-factor (Å ²)	55.1	Xtrriage
Anisotropy	0.329	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6027	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HU9, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.69	2/3004 (0.1%)	0.74	1/4065 (0.0%)
1	B	0.70	1/2997 (0.0%)	0.76	3/4055 (0.1%)
All	All	0.70	3/6001 (0.0%)	0.75	4/8120 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	269	VAL	CB-CG1	-8.74	1.34	1.52
1	B	308	CYS	CB-SG	-8.17	1.68	1.82
1	A	146	GLU	CG-CD	5.46	1.60	1.51

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	88	MET	CG-SD-CE	-6.54	89.74	100.20
1	B	100	ARG	NE-CZ-NH1	-6.08	117.26	120.30
1	A	385	MET	CG-SD-CE	5.62	109.19	100.20
1	B	308	CYS	CA-CB-SG	5.47	123.84	114.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2936	0	2941	34	0
1	B	2929	0	2934	43	0
2	A	43	0	30	5	0
2	B	43	0	30	4	0
3	A	19	0	0	1	0
3	B	19	0	0	1	0
4	A	21	0	0	1	0
4	B	17	0	0	0	0
All	All	6027	0	5935	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:26:PRO:O	1:A:74:LYS:NZ	2.12	0.83
1:A:62:LEU:O	1:A:106:ASN:ND2	2.12	0.82
1:B:64:MET:N	1:B:106:ASN:OD1	2.18	0.77
1:B:342:LEU:O	1:B:343:ARG:HB2	1.84	0.76
1:B:109:VAL:O	1:B:113:GLN:HG3	1.89	0.72
1:A:136:LYS:NZ	1:A:141:LYS:O	2.22	0.72
1:A:27:ASN:O	1:A:74:LYS:NZ	2.22	0.71
1:B:263:SER:HG	2:B:501:HEM:CGA	2.07	0.66
1:B:136:LYS:HE2	1:B:141:LYS:O	1.97	0.64
1:B:269:VAL:HG23	1:B:270:PHE:HD1	1.60	0.64
1:A:171:GLU:OE1	1:A:269:VAL:CG1	2.46	0.64
2:A:501:HEM:HHA	2:A:501:HEM:HBA2	1.83	0.61
1:A:171:GLU:OE1	1:A:269:VAL:HG13	2.01	0.61
1:A:12:SER:HA	1:A:13:LYS:CB	2.31	0.60
1:B:214:PHE:CE1	1:B:349:ILE:HD12	2.38	0.59
1:B:269:VAL:HG23	1:B:270:PHE:CD1	2.38	0.58
1:A:27:ASN:HB2	1:A:74:LYS:HZ1	1.69	0.57
1:A:144:THR:O	1:A:148:MET:HG3	2.05	0.57
1:B:34:ASP:O	1:B:37:ASN:OD1	2.25	0.55
1:B:99:VAL:HG11	1:B:243:LEU:HD11	1.89	0.54
1:A:56:ARG:O	1:A:60:GLU:OE2	2.26	0.52
1:A:264:ALA:HB3	2:A:501:HEM:C4D	2.46	0.51
1:B:347:LEU:CD2	1:B:385:MET:HE2	2.41	0.51
1:B:39:TRP:NE1	1:B:65:LEU:HD12	2.26	0.50
1:B:356:ILE:HB	1:B:357:PRO:HD3	1.94	0.50
1:A:189:GLN:HG2	1:A:190:MET:HE3	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:347:LEU:HD23	1:B:385:MET:HE2	1.95	0.49
1:B:342:LEU:O	1:B:343:ARG:CB	2.50	0.49
2:A:501:HEM:HBA2	2:A:501:HEM:CHA	2.44	0.48
1:A:189:GLN:HG2	1:A:190:MET:CE	2.44	0.48
1:A:285:GLY:N	4:A:602:HOH:O	2.33	0.48
1:B:172:ILE:O	1:B:175:ALA:HB3	2.13	0.48
1:B:14:GLU:HA	1:B:14:GLU:OE1	2.13	0.48
1:A:171:GLU:CD	1:A:269:VAL:CG1	2.82	0.48
1:B:347:LEU:HD23	1:B:385:MET:CE	2.44	0.48
1:A:80:ARG:NH2	1:A:128:ASP:OD2	2.46	0.47
1:A:171:GLU:CD	1:A:269:VAL:HG13	2.35	0.47
1:B:55:LEU:HD22	1:B:89:ALA:HB1	1.96	0.47
1:A:176:SER:HB3	1:A:206:CYS:SG	2.55	0.46
1:B:270:PHE:CE2	1:B:342:LEU:HD13	2.51	0.46
1:A:46:LEU:N	1:A:47:PRO:CD	2.79	0.45
1:B:214:PHE:CE1	1:B:349:ILE:CD1	3.00	0.45
1:B:240:ASN:O	1:B:244:SER:N	2.50	0.45
1:B:263:SER:OG	2:B:501:HEM:O2A	2.32	0.45
1:B:13:LYS:CB	1:B:14:GLU:CB	2.95	0.45
1:B:141:LYS:HB3	1:B:142:PRO:CD	2.48	0.44
1:B:214:PHE:HZ	1:B:346:HIS:HD2	1.65	0.44
1:A:59:VAL:HA	1:A:62:LEU:HD12	2.00	0.44
1:B:345:TYR:CE2	1:B:349:ILE:HD11	2.52	0.44
1:A:88:MET:HE1	1:A:123:ILE:HG13	1.99	0.44
1:A:42:ILE:HD12	1:A:62:LEU:HD11	2.00	0.44
1:B:321:LEU:HD21	1:B:400:LEU:HD22	1.99	0.44
1:A:245:ASP:O	1:A:256:PRO:HB2	2.18	0.43
1:B:46:LEU:O	1:B:50:ILE:HG13	2.19	0.43
1:B:72:ASP:O	1:B:76:GLN:NE2	2.52	0.43
1:A:175:ALA:HA	1:A:178:ILE:HD12	2.00	0.43
1:A:99:VAL:HG11	1:A:243:LEU:HD11	2.01	0.43
1:B:13:LYS:HA	1:B:14:GLU:HB2	2.00	0.43
1:B:336:VAL:O	1:B:340:VAL:HG23	2.19	0.43
1:B:40:MET:HA	1:B:40:MET:HE3	2.01	0.43
1:B:64:MET:HB2	1:B:106:ASN:OD1	2.18	0.43
1:A:50:ILE:HG21	1:A:155:ARG:HD3	2.01	0.42
1:A:12:SER:CA	1:A:13:LYS:CB	2.97	0.42
1:B:175:ALA:HA	1:B:178:ILE:HD12	2.00	0.42
1:B:264:ALA:HB3	2:B:501:HEM:C1D	2.55	0.42
1:A:275:VAL:HG13	1:A:311:GLU:HG3	2.00	0.42
1:B:91:VAL:O	1:B:99:VAL:HG13	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:LEU:HD12	1:A:26:PRO:HD2	2.01	0.42
2:A:501:HEM:NA	3:A:502:HU9:C17	2.83	0.42
1:B:270:PHE:HD1	1:B:270:PHE:N	2.17	0.42
1:A:87:THR:HG23	1:A:103:LEU:HD13	2.01	0.41
1:B:270:PHE:CD1	1:B:270:PHE:N	2.88	0.41
1:A:43:ALA:HB2	1:A:82:VAL:HG13	2.01	0.41
1:B:270:PHE:CZ	1:B:342:LEU:HD13	2.56	0.41
1:B:13:LYS:CB	1:B:14:GLU:HB3	2.50	0.41
1:A:138:ASP:OD2	1:A:141:LYS:HD2	2.20	0.41
1:B:19:GLU:O	1:B:19:GLU:HG3	2.21	0.41
2:B:501:HEM:NC	3:B:502:HU9:N19	2.69	0.41
1:B:356:ILE:CB	1:B:357:PRO:HD3	2.50	0.41
1:A:12:SER:O	1:A:17:ILE:O	2.40	0.41
1:A:346:HIS:HE1	2:A:501:HEM:NA	2.19	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	367/392 (94%)	353 (96%)	13 (4%)	1 (0%)	41	36
1	B	366/392 (93%)	344 (94%)	20 (6%)	2 (0%)	29	22
All	All	733/784 (94%)	697 (95%)	33 (4%)	3 (0%)	34	29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	13	LYS
1	B	244	SER
1	B	343	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	320/339 (94%)	295 (92%)	25 (8%)	12	7
1	B	319/339 (94%)	303 (95%)	16 (5%)	24	20
All	All	639/678 (94%)	598 (94%)	41 (6%)	17	12

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	SER
1	A	19	GLU
1	A	34	ASP
1	A	40	MET
1	A	44	LYS
1	A	48	ASP
1	A	60	GLU
1	A	61	LYS
1	A	62	LEU
1	A	66	SER
1	A	71	THR
1	A	80	ARG
1	A	141	LYS
1	A	214	PHE
1	A	217	ILE
1	A	222	ASN
1	A	237	TRP
1	A	238	LYS
1	A	244	SER
1	A	245	ASP
1	A	282	THR
1	A	309	SER
1	A	318	GLU
1	A	379	THR
1	A	393	SER
1	B	12	SER
1	B	14	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	58	ARG
1	B	80	ARG
1	B	105	ARG
1	B	107	ILE
1	B	116	LYS
1	B	135	LYS
1	B	212	GLN
1	B	214	PHE
1	B	222	ASN
1	B	237	TRP
1	B	272	CYS
1	B	318	GLU
1	B	355	LEU
1	B	382	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	501	3,1	41,50,50	1.64	3 (7%)	45,82,82	1.55	9 (20%)
2	HEM	B	501	3,1	41,50,50	1.44	2 (4%)	45,82,82	1.44	8 (17%)
3	HU9	B	502	2	20,21,21	0.98	2 (10%)	22,29,29	1.26	2 (9%)
3	HU9	A	502	2	20,21,21	1.15	3 (15%)	22,29,29	1.73	4 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	501	3,1	-	4/12/54/54	-
2	HEM	B	501	3,1	-	2/12/54/54	-
3	HU9	B	502	2	-	1/5/16/16	0/3/3/3
3	HU9	A	502	2	-	1/5/16/16	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	HEM	C3C-C2C	-6.23	1.31	1.40
2	B	501	HEM	C3C-C2C	-5.78	1.32	1.40
2	A	501	HEM	CAB-C3B	3.46	1.56	1.47
2	A	501	HEM	C3C-CAC	3.42	1.54	1.47
3	A	502	HU9	BR7-C05	2.90	1.96	1.89
3	A	502	HU9	C04-N08	2.89	1.46	1.37
3	B	502	HU9	C04-N08	2.67	1.45	1.37
3	B	502	HU9	C03-C02	-2.12	1.37	1.42
3	A	502	HU9	C03-C02	-2.03	1.37	1.42
2	B	501	HEM	CAB-C3B	2.01	1.52	1.47

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	HU9	BR7-C05-C04	4.44	124.92	119.73
2	B	501	HEM	C2C-C3C-C4C	3.75	109.52	106.90
2	B	501	HEM	CMB-C2B-C1B	-3.36	119.92	125.04
2	A	501	HEM	C4C-CHD-C1D	3.35	126.98	122.56
3	A	502	HU9	C15-C10-C09	-3.20	103.48	110.30
3	B	502	HU9	C15-C10-C09	-3.09	103.71	110.30
2	A	501	HEM	C1D-C2D-C3D	2.94	110.04	106.96
2	A	501	HEM	CAA-CBA-CGA	-2.89	105.65	113.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	HEM	CMB-C2B-C1B	-2.65	121.00	125.04
2	A	501	HEM	C3B-C2B-C1B	2.61	108.42	106.49
2	A	501	HEM	O1A-CGA-CBA	-2.61	114.71	123.08
3	A	502	HU9	C15-C10-C11	2.53	115.27	108.91
2	A	501	HEM	CHD-C1D-ND	2.47	127.12	124.43
3	B	502	HU9	BR7-C05-C04	2.42	122.55	119.73
2	B	501	HEM	CMD-C2D-C1D	-2.40	121.38	125.04
2	A	501	HEM	C2D-C1D-ND	-2.34	107.08	109.88
2	B	501	HEM	O1A-CGA-CBA	-2.32	115.62	123.08
2	B	501	HEM	CAA-CBA-CGA	-2.23	107.50	113.76
2	B	501	HEM	CAB-C3B-C2B	2.12	135.59	128.60
2	A	501	HEM	O2A-CGA-O1A	2.05	128.40	123.30
3	A	502	HU9	C10-C09-N08	-2.03	108.80	112.09
2	B	501	HEM	CMB-C2B-C3B	2.01	133.22	128.30
2	B	501	HEM	C4A-C3A-C2A	2.01	108.39	107.00

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	HU9	C10-C09-N08-C04
2	A	501	HEM	C1A-C2A-CAA-CBA
2	A	501	HEM	C3A-C2A-CAA-CBA
3	B	502	HU9	C10-C09-N08-C04
2	B	501	HEM	CAD-CBD-CGD-O2D
2	A	501	HEM	CAD-CBD-CGD-O2D
2	B	501	HEM	CAD-CBD-CGD-O1D
2	A	501	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.


4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	HEM	5	0
2	B	501	HEM	4	0
3	B	502	HU9	1	0
3	A	502	HU9	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	371/392 (94%)	0.43	29 (7%) 13 16	46, 63, 82, 94	0
1	B	370/392 (94%)	0.45	24 (6%) 18 23	48, 63, 80, 120	0
All	All	741/784 (94%)	0.44	53 (7%) 15 19	46, 63, 81, 120	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	12	SER	8.2
1	B	13	LYS	4.8
1	B	14	GLU	3.2
1	B	381	GLY	3.1
1	A	107	ILE	3.1
1	A	71	THR	3.1
1	B	200	LEU	3.0
1	A	350	VAL	2.9
1	A	254	GLU	2.9
1	A	328	LEU	2.9
1	A	379	THR	2.9
1	B	103	LEU	2.9
1	B	380	GLY	2.9
1	B	83	LEU	2.8
1	B	284	GLY	2.8
1	B	142	PRO	2.7
1	B	87	THR	2.7
1	A	13	LYS	2.7
1	A	335	CYS	2.7
1	B	107	ILE	2.7
1	A	200	LEU	2.6
1	A	380	GLY	2.6
1	B	108	ALA	2.6
1	A	105	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	328	LEU	2.6
1	B	230	LEU	2.5
1	A	41	PHE	2.5
1	A	83	LEU	2.5
1	A	103	LEU	2.5
1	A	287	HIS	2.5
1	A	139	PRO	2.5
1	A	86	ILE	2.5
1	B	196	LEU	2.5
1	A	255	ASP	2.5
1	B	184	VAL	2.4
1	A	203	ILE	2.4
1	A	87	THR	2.4
1	B	244	SER	2.4
1	B	287	HIS	2.4
1	A	284	GLY	2.3
1	A	34	ASP	2.3
1	A	332	TYR	2.3
1	A	165	LEU	2.3
1	A	140	ASN	2.2
1	A	277	LEU	2.1
1	B	168	LEU	2.1
1	B	249	TYR	2.1
1	A	141	LYS	2.1
1	A	15	TYR	2.1
1	A	61	LYS	2.1
1	B	34	ASP	2.0
1	B	70	LEU	2.0
1	B	203	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HEM	A	501	43/43	0.97	0.15	40,51,64,68	0
2	HEM	B	501	43/43	0.97	0.16	41,50,66,74	0
3	HU9	B	502	19/19	0.98	0.15	46,55,67,68	0
3	HU9	A	502	19/19	0.99	0.12	46,56,68,72	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around HEM B 501:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HU9 B 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.