



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 18, 2024 – 12:55 AM EDT

PDB ID : 3E15  
Title : 6-phosphogluconolactonase from Plasmodium vivax  
Authors : Arakaki, T.L.; Merritt, E.A.  
Deposited on : 2008-08-01  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	
1	B	312	
1	C	312	
1	D	312	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	AZI	A	307	-	X	-	-
4	AZI	C	306	-	X	-	-

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 10510 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucose-6-phosphate 1-dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	291	2406	1553	385	456	4	8	0	7	0
1	B	295	2441	1569	389	469	5	9	0	8	0
1	C	295	2442	1572	392	464	5	9	0	7	0
1	D	295	2451	1576	392	468	5	10	0	8	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	ALA	-	expression tag	UNP A5K3M1
A	-5	HIS	-	expression tag	UNP A5K3M1
A	-4	HIS	-	expression tag	UNP A5K3M1
A	-3	HIS	-	expression tag	UNP A5K3M1
A	-2	HIS	-	expression tag	UNP A5K3M1
A	-1	HIS	-	expression tag	UNP A5K3M1
A	0	HIS	-	expression tag	UNP A5K3M1
A	1	MSE	-	expression tag	UNP A5K3M1
B	-6	ALA	-	expression tag	UNP A5K3M1
B	-5	HIS	-	expression tag	UNP A5K3M1
B	-4	HIS	-	expression tag	UNP A5K3M1
B	-3	HIS	-	expression tag	UNP A5K3M1
B	-2	HIS	-	expression tag	UNP A5K3M1
B	-1	HIS	-	expression tag	UNP A5K3M1
B	0	HIS	-	expression tag	UNP A5K3M1
B	1	MSE	-	expression tag	UNP A5K3M1
C	-6	ALA	-	expression tag	UNP A5K3M1
C	-5	HIS	-	expression tag	UNP A5K3M1
C	-4	HIS	-	expression tag	UNP A5K3M1
C	-3	HIS	-	expression tag	UNP A5K3M1
C	-2	HIS	-	expression tag	UNP A5K3M1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	HIS	-	expression tag	UNP A5K3M1
C	0	HIS	-	expression tag	UNP A5K3M1
C	1	MSE	-	expression tag	UNP A5K3M1
D	-6	ALA	-	expression tag	UNP A5K3M1
D	-5	HIS	-	expression tag	UNP A5K3M1
D	-4	HIS	-	expression tag	UNP A5K3M1
D	-3	HIS	-	expression tag	UNP A5K3M1
D	-2	HIS	-	expression tag	UNP A5K3M1
D	-1	HIS	-	expression tag	UNP A5K3M1
D	0	HIS	-	expression tag	UNP A5K3M1
D	1	MSE	-	expression tag	UNP A5K3M1


- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Cl 1 1	0	0
2	D	1	Total Cl 1 1	0	0

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total K 1 1	0	0
3	B	1	Total K 1 1	0	0
3	C	1	Total K 1 1	0	0
3	D	1	Total K 1 1	0	0

- Molecule 4 is AZIDE ION (three-letter code: AZI) (formula: N<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total N 3 3	0	0
4	B	1	Total N 3 3	0	0
4	C	1	Total N 3 3	0	0
4	D	1	Total N 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	176	Total O 176 176	0	0
5	B	201	Total O 201 201	0	0
5	C	209	Total O 209 209	0	0
5	D	166	Total O 166 166	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glucose-6-phosphate 1-dehydrogenase




- Molecule 1: Glucose-6-phosphate 1-dehydrogenase



- Molecule 1: Glucose-6-phosphate 1-dehydrogenase



- Molecule 1: Glucose-6-phosphate 1-dehydrogenase







## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.52Å 80.85Å 89.77Å 63.98° 88.10° 80.01°	Depositor
Resolution (Å)	40.00 – 2.00 39.81 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.8 (40.00-2.00) 97.8 (39.81-2.00)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	8.46 (at 2.00Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.150 , 0.202 0.165 , 0.211	Depositor DCC
$R_{free}$ test set	3838 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.5	Xtrriage
Anisotropy	0.073	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 51.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for -h,-k,-k+1	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10510	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 75.83 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1708e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, AZI, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.64	0/2465	0.56	0/3317
1	B	0.67	0/2502	0.56	0/3364
1	C	0.68	0/2501	0.56	0/3364
1	D	0.64	0/2513	0.55	0/3377
All	All	0.66	0/9981	0.56	0/13422

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2406	0	2408	24	0
1	B	2441	0	2437	12	0
1	C	2442	0	2447	21	0
1	D	2451	0	2446	21	0
2	A	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	1	0	0	0	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
4	C	3	0	0	0	0
4	D	3	0	0	0	0
5	A	176	0	0	1	0
5	B	201	0	0	2	0
5	C	209	0	0	3	0
5	D	166	0	0	1	0
All	All	10510	0	9738	78	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (78) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:17:HIS:O	1:D:18[A]:ASN:ND2	1.79	1.15
1:C:18[B]:ASN:HD21	1:C:277[B]:MSE:SE	1.78	1.14
1:D:17:HIS:C	1:D:18[A]:ASN:HD22	1.69	0.96
1:C:18[B]:ASN:ND2	1:C:277[B]:MSE:SE	2.54	0.90
1:A:160:SER:HA	1:A:251:LEU:HD21	1.64	0.80
1:A:194:ALA:HB3	1:A:199:THR:OG1	1.90	0.71
1:B:190:SER:O	1:B:191:ILE:HD13	1.92	0.69
1:D:289:TYR:O	1:D:292[B]:MSE:HB2	1.96	0.65
1:A:63:THR:HB	1:A:64:PRO:HD3	1.79	0.63
1:C:277[A]:MSE:HE2	5:C:421:HOH:O	1.99	0.62
1:A:25:LYS:HZ2	1:A:33:LYS:HZ2	1.46	0.62
1:D:63:THR:HB	1:D:64:PRO:HD3	1.80	0.62
1:D:194:ALA:HB3	1:D:199:THR:OG1	2.00	0.61
1:D:150:LYS:HE2	1:D:237:SER:OG	2.00	0.61
1:C:142[B]:LYS:HG3	1:C:211:TYR:CD2	2.37	0.60
1:C:142[B]:LYS:HG2	1:C:211:TYR:CE2	2.37	0.59
1:D:277[A]:MSE:HE3	5:D:392:HOH:O	2.03	0.59
1:C:294:GLU:O	1:C:298:VAL:HG23	2.03	0.59
1:A:25:LYS:HZ2	1:A:33:LYS:NZ	2.00	0.58
1:B:160:SER:HA	1:B:251:LEU:HD11	1.86	0.58
1:A:25:LYS:NZ	1:A:33:LYS:NZ	2.51	0.58
1:C:1:MSE:HG2	1:C:6:LEU:HD11	1.88	0.56
1:D:295:ASP:OD2	1:D:295:ASP:N	2.39	0.55
1:A:224:LYS:NZ	5:A:438:HOH:O	2.33	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:142:LYS:HG2	1:D:211:TYR:CE2	2.42	0.54
1:C:29:ASP:O	1:C:33:LYS:HB2	2.10	0.52
1:B:51:GLY:O	1:B:83:LYS:HD3	2.10	0.52
1:B:277[B]:MSE:HE2	5:B:428:HOH:O	2.10	0.51
1:A:65:ILE:HD11	1:A:105:ASN:HB3	1.92	0.51
1:D:116:ASN:ND2	1:D:119[B]:GLU:CD	2.65	0.50
1:A:25:LYS:NZ	1:A:33:LYS:HZ2	2.11	0.49
1:C:277[A]:MSE:CE	5:C:421:HOH:O	2.59	0.49
1:C:140:LYS:NZ	5:C:504:HOH:O	2.31	0.49
1:D:190:SER:O	1:D:191:ILE:HD13	2.12	0.49
1:C:205:LEU:HD21	1:C:230:LEU:HD23	1.94	0.49
1:D:65:ILE:HD11	1:D:105:ASN:HB3	1.94	0.49
1:A:13:MSE:HE3	1:A:20:LYS:HB3	1.95	0.48
1:C:142[B]:LYS:CG	1:C:211:TYR:CD2	2.95	0.48
1:A:142:LYS:HG2	1:A:211:TYR:CE2	2.49	0.48
1:B:63:THR:HB	1:B:64:PRO:HD3	1.95	0.48
1:B:139[A]:GLU:OE1	1:B:142:LYS:HE2	2.13	0.48
1:A:15:HIS:CD2	1:A:20:LYS:HG2	2.49	0.47
1:B:50[A]:GLU:H	1:B:50[A]:GLU:CD	2.18	0.47
1:B:140:LYS:NZ	5:B:476:HOH:O	2.37	0.46
1:D:13:MSE:HE2	1:D:20:LYS:HB3	1.96	0.46
1:C:160:SER:HA	1:C:251:LEU:HD11	1.97	0.46
1:A:166:SER:HB2	1:A:227:THR:HB	1.97	0.46
1:C:17:HIS:O	1:C:18[B]:ASN:CB	2.65	0.45
1:A:17:HIS:O	1:A:18:ASN:HB2	2.16	0.45
1:A:160:SER:HA	1:A:251:LEU:CD2	2.43	0.45
1:B:295:ASP:HA	1:B:298:VAL:HG22	1.99	0.44
1:C:63:THR:HB	1:C:64:PRO:HD3	1.99	0.44
1:D:13:MSE:HE1	1:D:280:THR:HG23	2.00	0.44
1:A:41:GLN:HA	1:A:44[B]:GLU:HG2	1.99	0.44
1:C:276:SER:O	1:C:277[A]:MSE:HB2	2.18	0.44
1:B:29:ASP:OD2	1:B:33:LYS:NZ	2.43	0.44
1:A:25:LYS:NZ	1:A:33:LYS:HZ1	2.16	0.43
1:A:25:LYS:HZ1	1:A:33:LYS:HZ1	1.67	0.43
1:B:176:TYR:CE1	1:B:193:VAL:HG21	2.54	0.43
1:A:13:MSE:HG3	1:A:21:TYR:O	2.19	0.43
1:A:203:ASP:O	1:A:206:ASP:HB2	2.19	0.43
1:C:228:VAL:HB	1:C:232:LEU:HD23	2.01	0.43
1:C:37:TYR:CE2	1:C:280:THR:HG21	2.55	0.42
1:D:289:TYR:HB3	1:D:290:PRO:HD3	2.02	0.42
1:A:53:HIS:HA	1:A:83:LYS:O	2.20	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:180:TYR:CE1	1:D:191:ILE:HD11	2.55	0.42
1:A:142:LYS:CG	1:A:211:TYR:CE2	3.03	0.41
1:C:134:VAL:HG21	1:C:214:PHE:HB2	2.01	0.41
1:C:18[B]:ASN:CG	1:C:277[B]:MSE:SE	3.09	0.41
1:D:134:VAL:HG21	1:D:214:PHE:HB2	2.02	0.41
1:C:30:PHE:CZ	1:C:244:ASN:HB3	2.54	0.41
1:B:160:SER:HA	1:B:251:LEU:CD1	2.50	0.41
1:D:77:ILE:O	1:D:77:ILE:HG23	2.20	0.41
1:A:289:TYR:N	1:A:290:PRO:CD	2.84	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	294/312 (94%)	285 (97%)	8 (3%)	1 (0%)	41 37
1	B	299/312 (96%)	290 (97%)	8 (3%)	1 (0%)	41 37
1	C	298/312 (96%)	288 (97%)	9 (3%)	1 (0%)	41 37
1	D	299/312 (96%)	292 (98%)	6 (2%)	1 (0%)	41 37
All	All	1190/1248 (95%)	1155 (97%)	31 (3%)	4 (0%)	41 37

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	222	VAL
1	B	222	VAL
1	C	222	VAL
1	D	222	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/283 (97%)	272 (99%)	2 (1%)	84	88
1	B	281/283 (99%)	281 (100%)	0	100	100
1	C	280/283 (99%)	279 (100%)	1 (0%)	91	93
1	D	281/283 (99%)	274 (98%)	7 (2%)	47	49
All	All	1116/1132 (99%)	1106 (99%)	10 (1%)	84	83

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	75	LYS
1	A	296	ILE
1	C	150	LYS
1	D	18[A]	ASN
1	D	18[B]	ASN
1	D	40[A]	HIS
1	D	40[B]	HIS
1	D	75	LYS
1	D	191	ILE
1	D	295	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	AZI	B	306	-	2,2,2	1.89	1 (50%)	0,1,1	-	-
4	AZI	A	307	-	2,2,2	2.23	2 (100%)	0,1,1	-	-
4	AZI	C	306	-	2,2,2	2.14	2 (100%)	0,1,1	-	-
4	AZI	D	307	-	2,2,2	1.96	1 (50%)	0,1,1	-	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	307	AZI	N3-N2	-2.31	1.18	1.23
4	A	307	AZI	N1-N2	-2.16	1.18	1.23
4	C	306	AZI	N1-N2	-2.15	1.18	1.23
4	C	306	AZI	N3-N2	-2.14	1.18	1.23
4	D	307	AZI	N3-N2	-2.11	1.18	1.23
4	B	306	AZI	N3-N2	-2.07	1.18	1.23

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	284/312 (91%)	-0.24	14 (4%) 29 28	7, 16, 46, 72	1 (0%)
1	B	287/312 (91%)	-0.51	5 (1%) 70 68	6, 13, 35, 48	0
1	C	287/312 (91%)	-0.54	5 (1%) 70 68	6, 13, 34, 54	0
1	D	287/312 (91%)	-0.24	16 (5%) 24 23	7, 17, 46, 57	1 (0%)
All	All	1145/1248 (91%)	-0.38	40 (3%) 44 43	6, 15, 41, 72	2 (0%)

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	8	LYS	5.0
1	A	6	LEU	4.5
1	D	3	CYS	4.2
1	D	5	ALA	4.1
1	D	2	ASP	4.1
1	C	298	VAL	4.0
1	D	190	SER	4.0
1	A	295	ASP	3.8
1	B	72	ALA	3.8
1	B	199	THR	3.7
1	A	296	ILE	3.6
1	A	73	LEU	3.5
1	A	5	ALA	3.2
1	A	72	ALA	3.1
1	C	199	THR	3.1
1	D	193	VAL	3.0
1	D	291[A]	GLN	2.9
1	D	73	LEU	2.9
1	D	297	TYR	2.8
1	B	298	VAL	2.8
1	D	72	ALA	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	296	ILE	2.7
1	D	11	GLU	2.7
1	A	11	GLU	2.7
1	D	6	LEU	2.7
1	A	9	SER	2.5
1	D	295	ASP	2.5
1	A	10	LEU	2.5
1	A	75	LYS	2.5
1	A	27	LEU	2.4
1	C	4	GLN	2.3
1	B	73	LEU	2.3
1	D	8	LYS	2.3
1	C	195	ASN	2.2
1	B	3	CYS	2.2
1	D	191	ILE	2.1
1	D	4	GLN	2.1
1	C	191	ILE	2.0
1	A	297	TYR	2.0
1	A	7	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	AZI	B	306	3/3	0.90	0.14	14,14,22,32	0
4	AZI	C	306	3/3	0.91	0.11	17,17,18,30	0
4	AZI	A	307	3/3	0.92	0.14	16,16,22,30	0
4	AZI	D	307	3/3	0.93	0.11	16,16,21,32	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	K	B	305	1/1	0.99	0.05	20,20,20,20	0
3	K	D	306	1/1	1.00	0.03	18,18,18,18	0
2	CL	D	305	1/1	1.00	0.05	9,9,9,9	0
3	K	A	306	1/1	1.00	0.03	19,19,19,19	0
2	CL	A	305	1/1	1.00	0.07	9,9,9,9	0
3	K	C	305	1/1	1.00	0.03	19,19,19,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.