



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2023 – 12:52 AM JST

PDB ID : 7E36  
Title : A [6+4]-cycloaddition adduct is the biosynthetic intermediate in streptoseomycin biosynthesis  
Authors : Zhang, B.; Ge, H.M.  
Deposited on : 2021-02-08  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	324	 5% 90% 8%
2	B	421	 6% 82% 11% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CL	A	406	-	-	X	-

## 2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5989 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alkanesulfonate monooxygenase SsuD/methylene tetrahydro methanopterin reductase-like flavin-dependent oxidoreductase (Luciferase family).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	319	2384	1477	456	450	1	0	0	0

- Molecule 2 is a protein called FMN-dependent oxidoreductase (Nitrilotriacetate monooxygenase family).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	395	3075	1942	559	567	7	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	A	1	5	4	1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	7	Total	Cl	0	0
			7	7		
4	B	1	Total	Cl	0	0
			1	1		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		

- Molecule 6 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
6	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 7 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C<sub>4</sub>H<sub>12</sub>NO<sub>3</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
7	B	1	8	4	1	3	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
8	A	231	231	231	0	0
8	B	260	260	260	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alkanesulfonate monooxygenase SsuD/methylene tetrahydromethanopterin reductase-like flavin-dependent oxidoreductase (Luciferase family)



- Molecule 2: FMN-dependent oxidoreductase (Nitrilotriacetate monooxygenase family)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.28Å 104.99Å 75.95Å 90.00° 95.95° 90.00°	Depositor
Resolution (Å)	51.00 – 2.00 51.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (51.00-2.00) 99.6 (51.00-2.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.69 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, $R_{free}$	0.165 , 0.205 0.165 , 0.204	Depositor DCC
$R_{free}$ test set	1997 reflections (3.70%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.6	Xtrriage
Anisotropy	0.465	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 56.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5989	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SO4, TRS, MES, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.27	0/2425	0.49	0/3299
2	B	0.29	0/3152	0.50	0/4292
All	All	0.28	0/5577	0.49	0/7591

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2384	0	2361	15	0
2	B	3075	0	3000	34	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	7	0	0	4	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
6	B	12	0	12	0	0
7	B	8	0	12	1	0
8	A	231	0	0	2	1
8	B	260	0	0	3	1
All	All	5989	0	5385	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:134:SER:HB3	2:B:154:ARG:HD3	1.47	0.94
2:B:405:THR:HG22	2:B:408:ASP:H	1.33	0.92
4:A:407:CL:CL	8:A:721:HOH:O	2.34	0.81
1:A:148:SER:HB2	4:A:406:CL:CL	2.22	0.76
2:B:327:ALA:O	2:B:329:ARG:N	2.19	0.76
1:A:185:LEU:N	4:A:406:CL:CL	2.60	0.71
2:B:328:ARG:O	2:B:330:GLU:N	2.23	0.71
1:A:282:VAL:HG23	1:A:284:ASP:H	1.59	0.68
4:A:403:CL:CL	8:A:710:HOH:O	2.52	0.63
2:B:61:VAL:HG11	2:B:139:GLU:OE2	1.98	0.62
1:A:100:SER:O	1:A:103:ARG:HD3	2.01	0.60
2:B:329:ARG:O	2:B:331:ASN:N	2.34	0.59
1:A:55:ASP:OD1	2:B:173:ARG:NH1	2.32	0.58
2:B:273:LEU:HD21	2:B:342:ALA:HB2	1.87	0.56
1:A:297:ARG:HD2	1:A:302:PRO:HD3	1.90	0.54
2:B:105:THR:HG21	2:B:134:SER:HB2	1.88	0.54
2:B:151:HIS:HA	2:B:154:ARG:HE	1.74	0.52
2:B:151:HIS:HA	2:B:154:ARG:NE	2.25	0.52
2:B:349:THR:CG2	2:B:352:GLN:H	2.25	0.50
1:A:217:VAL:HG21	1:A:231:VAL:HG11	1.94	0.49
2:B:86:LEU:HD22	2:B:98:VAL:HB	1.95	0.49
2:B:108:ASP:OD2	8:B:601:HOH:O	2.20	0.49
2:B:405:THR:HG21	8:B:824:HOH:O	2.12	0.48
1:A:279:ASP:OD1	1:A:280:ARG:N	2.46	0.48
2:B:214:ALA:HB1	2:B:221:ILE:HD12	1.95	0.47
1:A:43:LEU:HD12	1:A:63:ALA:HB2	1.97	0.46
2:B:405:THR:HG22	2:B:408:ASP:N	2.16	0.46
1:A:180:ASP:HB2	2:B:144:GLY:HA2	1.98	0.45
2:B:273:LEU:HD21	2:B:342:ALA:CB	2.46	0.45
1:A:280:ARG:O	1:A:282:VAL:HG22	2.17	0.45
2:B:349:THR:OG1	2:B:350:PRO:HD2	2.16	0.45
1:A:53:ALA:N	1:A:54:PRO:HD3	2.32	0.45
2:B:307:VAL:HG22	2:B:334:LEU:HD23	1.99	0.45
1:A:278:ALA:O	1:A:281:SER:OG	2.35	0.44
2:B:203:GLN:HG2	2:B:214:ALA:HA	1.99	0.44
1:A:77:PRO:HD2	1:A:105:GLY:O	2.18	0.43
2:B:65:GLY:H	2:B:68:LEU:HG	1.84	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:41:ARG:HH22	7:B:503:TRS:H21	1.83	0.43
2:B:235:TYR:CZ	2:B:239:LYS:HE3	2.53	0.43
2:B:373:PHE:HA	2:B:374:PRO:HD3	1.92	0.42
2:B:293:LEU:HD22	2:B:326:LEU:HD12	2.01	0.42
2:B:8:LEU:HD12	2:B:368:GLY:HA2	2.02	0.42
2:B:327:ALA:N	8:B:620:HOH:O	2.52	0.42
2:B:338:SER:HA	2:B:341:THR:HG22	2.03	0.41
2:B:64:TRP:CE3	2:B:68:LEU:HD21	2.56	0.41
1:A:164:PHE:HZ	2:B:68:LEU:HD22	1.85	0.40
2:B:329:ARG:HE	2:B:329:ARG:HB3	1.68	0.40
2:B:402:LEU:O	2:B:404:GLY:N	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:730:HOH:O	8:B:766:HOH:O[1_554]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	317/324 (98%)	307 (97%)	8 (2%)	2 (1%)	25 19
2	B	391/421 (93%)	370 (95%)	17 (4%)	4 (1%)	15 9
All	All	708/745 (95%)	677 (96%)	25 (4%)	6 (1%)	19 13

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	146	PRO
2	B	328	ARG

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	329	ARG
2	B	330	GLU
1	A	16	ALA
1	A	282	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/234 (99%)	227 (98%)	4 (2%)	60	65
2	B	317/338 (94%)	308 (97%)	9 (3%)	43	44
All	All	548/572 (96%)	535 (98%)	13 (2%)	49	51

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	LEU
1	A	165	ARG
1	A	169	LEU
1	A	283	ASP
2	B	5	THR
2	B	42	ARG
2	B	203	GLN
2	B	287	ARG
2	B	328	ARG
2	B	329	ARG
2	B	339	LEU
2	B	400	SER
2	B	405	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 9 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	A	401	-	4,4,4	0.12	0	6,6,6	0.09	0
3	SO4	B	502	-	4,4,4	0.14	0	6,6,6	0.09	0
7	TRS	B	503	-	7,7,7	0.34	0	9,9,9	0.35	0
6	MES	B	501	-	12,12,12	2.11	1 (8%)	14,16,16	2.47	7 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	TRS	B	503	-	-	0/9/9/9	-
6	MES	B	501	-	-	1/6/14/14	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	501	MES	C8-S	-7.00	1.67	1.77

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	501	MES	C5-N4-C3	4.93	119.93	108.83
6	B	501	MES	C7-N4-C5	3.44	120.04	111.23
6	B	501	MES	C7-N4-C3	3.28	119.61	111.23
6	B	501	MES	C2-C3-N4	-3.25	105.18	110.10
6	B	501	MES	C6-C5-N4	-2.74	105.94	110.10
6	B	501	MES	O1S-S-C8	2.61	110.06	106.92
6	B	501	MES	O2S-S-C8	2.07	109.41	106.92

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	501	MES	C8-C7-N4-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	503	TRS	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	319/324 (98%)	-0.11	15 (4%) 31 30	16, 27, 61, 105	0
2	B	395/421 (93%)	0.18	26 (6%) 18 17	16, 27, 71, 142	0
All	All	714/745 (95%)	0.05	41 (5%) 23 23	16, 27, 69, 142	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	135	ALA	23.7
2	B	148	HIS	7.8
2	B	146	PRO	7.2
2	B	342	ALA	7.1
1	A	281	SER	7.0
1	A	53	ALA	6.9
2	B	310	LEU	6.2
2	B	147	GLU	5.7
1	A	51	PRO	5.6
2	B	414	ASP	5.6
2	B	295	ALA	5.4
1	A	52	GLY	4.6
1	A	2	SER	4.2
1	A	16	ALA	4.1
2	B	331	ASN	4.0
1	A	282	VAL	4.0
2	B	149	MET	3.8
1	A	279	ASP	3.8
2	B	226	THR	3.7
1	A	283	ASP	3.6
2	B	136	ALA	3.6
2	B	138	TRP	3.6
2	B	309	GLU	3.4
2	B	296	VAL	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	341	THR	3.1
1	A	124	SER	3.0
2	B	151	HIS	3.0
2	B	179	HIS	2.9
2	B	137	PRO	2.9
2	B	308	PRO	2.9
1	A	17	GLU	2.8
2	B	330	GLU	2.8
2	B	5	THR	2.7
1	A	125	ALA	2.7
2	B	60	VAL	2.6
1	A	280	ARG	2.5
1	A	304	GLN	2.3
1	A	50	ARG	2.3
2	B	329	ARG	2.1
2	B	145	PHE	2.0
2	B	326	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	CL	A	406	1/1	0.55	0.25	74,74,74,74	0
7	TRS	B	503	8/8	0.79	0.26	50,55,60,70	0
5	MG	A	409	1/1	0.90	0.09	35,35,35,35	0
4	CL	A	405	1/1	0.93	0.05	55,55,55,55	0
4	CL	A	407	1/1	0.94	0.08	60,60,60,60	0
4	CL	B	504	1/1	0.96	0.12	59,59,59,59	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	CL	A	404	1/1	0.96	0.13	58,58,58,58	0
4	CL	A	408	1/1	0.96	0.28	30,30,30,30	0
6	MES	B	501	12/12	0.97	0.11	33,40,45,55	0
3	SO4	B	502	5/5	0.98	0.12	46,49,52,57	0
4	CL	A	402	1/1	0.98	0.07	42,42,42,42	0
4	CL	A	403	1/1	0.98	0.28	56,56,56,56	0
3	SO4	A	401	5/5	0.98	0.08	51,57,60,73	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.