



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 11:48 am BST

PDB ID : 5E5C  
Title : Crystal structure of dihydropyrimidinase from *Pseudomonas aeruginosa* PAO1  
Authors : Huang, C.C.; Huang, Y.H.; Hsieh, Y.C.; Tzeng, C.T.; Chen, C.J.; Huang, C.Y.  
Deposited on : 2015-10-08  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	485	 90% 7% ..
1	C	485	 3% 83% 15% ..

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 7698 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called D-hydantoinase/dihydropyrimidinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	478	3674	2312	660	686	16	0	0	0
1	C	479	3682	2317	661	687	17	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	480	HIS	-	expression tag	UNP Q9I676
A	481	HIS	-	expression tag	UNP Q9I676
A	482	HIS	-	expression tag	UNP Q9I676
A	483	HIS	-	expression tag	UNP Q9I676
A	484	HIS	-	expression tag	UNP Q9I676
A	485	HIS	-	expression tag	UNP Q9I676
C	480	HIS	-	expression tag	UNP Q9I676
C	481	HIS	-	expression tag	UNP Q9I676
C	482	HIS	-	expression tag	UNP Q9I676
C	483	HIS	-	expression tag	UNP Q9I676
C	484	HIS	-	expression tag	UNP Q9I676
C	485	HIS	-	expression tag	UNP Q9I676

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	227	Total 227	O 227	0	0
3	C	111	Total 111	O 111	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: D-hydantoinase/dihydropyrimidinase



- Molecule 1: D-hydantoinase/dihydropyrimidinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.13Å 112.13Å 161.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	97.11 – 2.10 27.93 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.6 (97.11-2.10) 99.6 (27.93-2.10)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.93 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.170 , 0.215 0.180 , 0.223	Depositor DCC
$R_{free}$ test set	3471 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.3	Xtrriage
Anisotropy	0.013	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 49.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7698	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.87% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.06	3/3753 (0.1%)	1.01	13/5104 (0.3%)
1	C	0.93	3/3761 (0.1%)	0.95	7/5114 (0.1%)
All	All	1.00	6/7514 (0.1%)	0.98	20/10218 (0.2%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	369	SER	CA-CB	6.73	1.63	1.52
1	A	460	SER	CB-OG	-6.41	1.33	1.42
1	A	101	SER	CB-OG	5.46	1.49	1.42
1	C	363	GLU	CD-OE2	-5.28	1.19	1.25
1	C	236	TYR	CG-CD1	-5.17	1.32	1.39
1	A	467	ARG	CZ-NH1	5.03	1.39	1.33

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	74	ASP	CB-CG-OD1	7.82	125.34	118.30
1	A	172	ARG	NE-CZ-NH2	-7.33	116.64	120.30
1	A	412	ARG	NE-CZ-NH1	7.14	123.87	120.30
1	A	468	ARG	NE-CZ-NH2	-6.84	116.88	120.30
1	A	467	ARG	NE-CZ-NH1	6.80	123.70	120.30
1	A	467	ARG	NE-CZ-NH2	-6.77	116.92	120.30
1	A	460	SER	N-CA-CB	-6.65	100.53	110.50
1	C	223	ARG	NE-CZ-NH1	-6.54	117.03	120.30
1	A	412	ARG	NE-CZ-NH2	-6.38	117.11	120.30
1	A	74	ASP	CB-CG-OD2	-6.01	112.89	118.30
1	A	272	ASP	CB-CG-OD1	5.73	123.46	118.30
1	A	182	VAL	CG1-CB-CG2	-5.61	101.92	110.90
1	A	45	ASP	CB-CG-OD1	5.53	123.28	118.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	253	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	C	135	ARG	NE-CZ-NH1	5.43	123.02	120.30
1	C	442	ASP	CB-CG-OD1	5.21	122.99	118.30
1	C	226	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	C	345	ARG	NE-CZ-NH1	5.10	122.85	120.30
1	A	386	ARG	NE-CZ-NH1	5.01	122.80	120.30
1	C	226	ARG	NE-CZ-NH1	5.00	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3674	0	3558	19	0
1	C	3682	0	3570	45	0
2	A	2	0	0	0	0
2	C	2	0	0	0	0
3	A	227	0	0	3	0
3	C	111	0	0	3	0
All	All	7698	0	7128	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:467:ARG:NH2	3:A:601:HOH:O	2.17	0.75
1:C:188:GLU:OE1	3:C:601:HOH:O	2.03	0.75
1:A:64:LEU:HD13	1:A:66:PHE:HB2	1.69	0.73
1:C:233:THR:HG23	1:C:234:PRO:O	1.93	0.69
1:C:75:PHE:O	1:C:79:THR:HG22	1.92	0.68
1:C:317:HIS:HD2	1:C:341:GLY:H	1.42	0.68
1:A:321:CYS:SG	3:A:824:HOH:O	2.43	0.67

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:112:TRP:O	1:C:115:LYS:HE3	1.96	0.65
1:A:257:GLN:HG3	3:A:805:HOH:O	2.00	0.62
1:C:183:HIS:HE1	1:C:185:GLU:OE1	1.82	0.62
1:C:24:ALA:HB3	1:C:29:GLN:HG3	1.81	0.61
1:A:239:HIS:CD2	1:A:239:HIS:O	2.54	0.60
1:C:132:GLU:HA	1:C:132:GLU:OE1	2.02	0.59
1:C:317:HIS:CD2	1:C:341:GLY:H	2.22	0.58
1:A:280:TRP:CZ2	1:A:329:ARG:HA	2.40	0.56
1:A:432:SER:HB3	1:A:437:LEU:HD12	1.87	0.56
1:A:239:HIS:O	1:A:239:HIS:HD2	1.88	0.55
1:C:243:ARG:NH2	1:C:296:GLU:HG3	2.23	0.54
1:A:64:LEU:HD13	1:A:66:PHE:CB	2.38	0.54
1:C:112:TRP:O	1:C:115:LYS:CE	2.57	0.53
1:C:467:ARG:NH2	3:C:608:HOH:O	2.41	0.53
1:C:406:ALA:HA	1:C:409:HIS:CE1	2.45	0.52
1:C:79:THR:CG2	1:C:117:ALA:H	2.23	0.52
1:C:59:HIS:CD2	1:C:316:ASP:HA	2.46	0.51
1:A:72:SER:O	1:A:412:ARG:HD2	2.11	0.50
1:C:75:PHE:O	1:C:79:THR:CG2	2.59	0.50
1:C:432:SER:HB3	1:C:437:LEU:HD12	1.93	0.50
1:C:324:GLN:HA	1:C:327:MET:HE2	1.94	0.48
1:C:228:ALA:HA	1:C:233:THR:HG22	1.96	0.48
1:C:76:PHE:HB2	1:C:115:LYS:HG3	1.97	0.47
1:C:64:LEU:HG	1:C:66:PHE:HB2	1.95	0.47
1:C:24:ALA:HB3	1:C:29:GLN:CG	2.45	0.47
1:C:402:ARG:HB2	1:C:426:ILE:HG21	1.96	0.46
1:A:271:ASP:HA	1:A:293:ARG:O	2.16	0.46
1:C:24:ALA:O	1:C:29:GLN:NE2	2.39	0.46
1:C:1:MET:HB3	1:C:2:SER:HB2	1.98	0.45
1:C:370:THR:O	1:C:374:LYS:HG3	2.17	0.45
1:C:280:TRP:CZ2	1:C:329:ARG:HA	2.52	0.45
1:C:229:GLU:OE1	3:C:602:HOH:O	2.21	0.44
1:C:79:THR:HG21	1:C:117:ALA:H	1.83	0.44
1:C:208:HIS:N	1:C:209:PRO:HD2	2.33	0.44
1:C:429:HIS:CD2	1:C:439:ALA:HA	2.54	0.43
1:A:479:ARG:NH2	1:C:277:HIS:O	2.50	0.43
1:A:213:PRO:O	1:A:216:VAL:HB	2.18	0.43
1:A:64:LEU:HD22	1:A:65:PRO:O	2.19	0.43
1:C:417:ILE:HG13	1:C:418:PHE:CD2	2.54	0.43
1:A:234:PRO:HD3	1:A:465:LEU:CD1	2.49	0.42
1:C:114:GLN:O	1:C:448:GLY:HA2	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:65:PRO:HA	1:C:69:THR:O	2.19	0.42
1:C:153:MET:SD	1:C:182:VAL:HG11	2.59	0.42
1:C:465:LEU:HD23	1:C:465:LEU:HA	1.87	0.42
1:A:238:VAL:HG23	1:A:239:HIS:CD2	2.55	0.41
1:C:325:LYS:HA	1:C:336:PRO:HD2	2.02	0.41
1:C:317:HIS:HD2	1:C:341:GLY:N	2.15	0.41
1:A:208:HIS:N	1:A:209:PRO:HD2	2.36	0.41
1:C:141:VAL:HA	1:C:146:VAL:O	2.21	0.41
1:C:397:ASP:C	1:C:397:ASP:OD1	2.58	0.41
1:C:289:SER:HA	1:C:290:PRO:C	2.41	0.40
1:A:183:HIS:CD2	1:A:183:HIS:C	2.94	0.40
1:C:317:HIS:CD2	1:C:342:ILE:H	2.39	0.40
1:A:479:ARG:NH1	1:C:206:GLU:OE1	2.51	0.40
1:C:238:VAL:HG23	1:C:239:HIS:CD2	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	475/485 (98%)	456 (96%)	16 (3%)	3 (1%)	25	21
1	C	476/485 (98%)	456 (96%)	18 (4%)	2 (0%)	34	32
All	All	951/970 (98%)	912 (96%)	34 (4%)	5 (0%)	29	26

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	239	HIS
1	C	239	HIS
1	C	341	GLY
1	A	341	GLY

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	63	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	375/382 (98%)	366 (98%)	9 (2%)	49	53
1	C	376/382 (98%)	370 (98%)	6 (2%)	62	69
All	All	751/764 (98%)	736 (98%)	15 (2%)	55	60

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	ARG
1	A	64	LEU
1	A	109	TRP
1	A	329	ARG
1	A	337	ASN
1	A	399	GLN
1	A	409	HIS
1	A	478	GLU
1	A	479	ARG
1	C	109	TRP
1	C	114	GLN
1	C	181	THR
1	C	233	THR
1	C	401	SER
1	C	409	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	200	GLN
1	A	239	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	306	GLN
1	C	183	HIS
1	C	200	GLN
1	C	239	HIS
1	C	317	HIS
1	C	337	ASN
1	C	411	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	KCX	A	150	1,2	7,11,12	1.36	0	4,12,14	0.60	0
1	KCX	C	150	1,2	7,11,12	1.46	1 (14%)	4,12,14	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	150	1,2	-	0/7/10/12	-
1	KCX	C	150	1,2	-	0/7/10/12	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	150	KCX	CB-CA	-3.23	1.49	1.53

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	477/485 (98%)	-0.41	2 (0%) 92 93	12, 22, 42, 69	0
1	C	478/485 (98%)	0.04	14 (2%) 51 57	16, 33, 52, 96	0
All	All	955/970 (98%)	-0.19	16 (1%) 70 74	12, 27, 49, 96	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	MET	9.4
1	C	473	ARG	4.3
1	C	33	GLU	3.6
1	C	407	ALA	3.5
1	C	99	ARG	3.4
1	C	40	GLY	2.8
1	C	69	THR	2.6
1	C	68	GLY	2.4
1	C	323	GLU	2.4
1	C	330	ASP	2.4
1	C	475	VAL	2.3
1	C	478	GLU	2.3
1	A	39	SER	2.2
1	C	157	ASN	2.1
1	C	342	ILE	2.1
1	A	473	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	KCX	A	150	12/13	0.93	0.15	14,17,33,38	0
1	KCX	C	150	12/13	0.94	0.16	20,25,40,43	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ZN	A	501	1/1	0.97	0.04	47,47,47,47	0
2	ZN	A	502	1/1	0.99	0.05	32,32,32,32	0
2	ZN	C	502	1/1	0.99	0.03	40,40,40,40	0
2	ZN	C	501	1/1	0.99	0.03	49,49,49,49	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.