



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 20, 2023 – 11:20 PM JST

PDB ID : 7E58
Title : interferon-inducible anti-viral protein 2
Authors : Cui, W.; Wang, W.; Chen, C.; Slater, B.; Xiong, Y.; Ji, X.Y.; Yang, H.T.
Deposited on : 2021-02-18
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	591	
1	B	591	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 17718 atoms, of which 8843 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanylate-binding protein 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	549	8818	2787	4416	743	849	23	0	0	0
1	B	552	8863	2806	4427	751	856	23	0	1	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	310	ARG	MET	conflict	UNP P32456
B	310	ARG	MET	conflict	UNP P32456


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	10	Total	O	0	0
			10	10		
2	B	27	Total	O	0	0
			27	27		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Guanylate-binding protein 2



- Molecule 1: Guanylate-binding protein 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	42.21Å 124.16Å 355.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.03 – 2.60 47.03 – 2.60	Depositor EDS
% Data completeness (in resolution range)	93.1 (47.03-2.60) 93.1 (47.03-2.60)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.61Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.255 , 0.272 0.255 , 0.271	Depositor DCC
R_{free} test set	2779 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	52.2	Xtrriage
Anisotropy	0.794	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 34.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	17718	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/4474	0.40	0/6015
1	B	0.25	0/4508	0.40	0/6059
All	All	0.25	0/8982	0.40	0/12074

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4402	4416	4428	19	0
1	B	4436	4427	4459	11	1
2	A	10	0	0	1	0
2	B	27	0	0	1	0
All	All	8875	8843	8887	30	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:448:VAL:O	1:A:451:LYS:NZ	2.02	0.92

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:181:ARG:NH1	2:A:601:HOH:O	2.22	0.72
1:A:12:SER:OG	1:A:15:ASP:OD1	2.12	0.66
1:B:387:GLU:OE2	2:B:601:HOH:O	2.13	0.66
1:A:168:ASP:OD1	1:A:580:ARG:NH2	2.32	0.62
1:B:154:ALA:O	1:B:310[B]:ARG:NH2	2.32	0.62
1:A:206:LEU:HD11	1:A:222:LEU:HD21	1.83	0.61
1:A:16:ASN:OD1	1:A:111:ASN:ND2	2.30	0.58
1:B:50:GLY:O	1:B:181:ARG:NH1	2.40	0.55
1:B:448:VAL:HG12	1:B:451:LYS:HG3	1.88	0.55
1:A:163:VAL:HG11	1:A:580:ARG:HH21	1.71	0.53
1:A:44:VAL:HG21	1:A:126:PHE:CE2	2.45	0.52
1:A:450:ARG:CG	1:A:450:ARG:O	2.60	0.50
1:A:206:LEU:HD11	1:A:222:LEU:CD2	2.43	0.48
1:A:337:MET:HA	1:A:337:MET:HE2	1.97	0.47
1:B:23:VAL:HG13	1:B:299:VAL:HG11	1.95	0.47
1:A:46:LEU:H	1:A:46:LEU:HD12	1.80	0.47
1:A:485:LYS:O	1:A:489:VAL:HG23	2.16	0.46
1:B:359:GLU:HG3	1:B:386:LEU:HD13	1.97	0.46
1:A:155:ASN:ND2	1:A:168:ASP:OD2	2.50	0.43
1:B:44:VAL:HG21	1:B:126:PHE:CE2	2.54	0.43
1:A:332:HIS:O	1:A:336:GLN:HG2	2.19	0.43
1:A:46:LEU:HD12	1:A:46:LEU:N	2.34	0.42
1:A:359:GLU:HG3	1:A:386:LEU:HD13	2.00	0.42
1:B:129:ASN:OD1	1:B:181:ARG:HD2	2.20	0.42
1:B:451:LYS:HE3	1:B:451:LYS:HB2	1.75	0.42
1:B:259:ASP:O	1:B:263:GLN:HG3	2.19	0.41
1:B:450:ARG:O	1:B:450:ARG:CG	2.68	0.41
1:A:219:ASP:HB3	1:A:220:PRO:HD3	2.03	0.41
1:A:337:MET:HA	1:A:337:MET:CE	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:212:LYS:HZ3	1:B:262:GLU:OE2[1_455]	1.60	0.00

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	539/591 (91%)	535 (99%)	4 (1%)	0	100	100
1	B	543/591 (92%)	538 (99%)	5 (1%)	0	100	100
All	All	1082/1182 (92%)	1073 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	488/525 (93%)	465 (95%)	23 (5%)	26	50
1	B	491/525 (94%)	478 (97%)	13 (3%)	46	72
All	All	979/1050 (93%)	943 (96%)	36 (4%)	35	60

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	SER
1	A	17	THR
1	A	76	LYS
1	A	87	LYS
1	A	88	LYS
1	A	194	ILE
1	A	207	ARG
1	A	213	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	310	ARG
1	A	335	GLN
1	A	406	MET
1	A	444	LYS
1	A	461	LYS
1	A	464	GLU
1	A	473	LEU
1	A	504	LEU
1	A	537	ARG
1	A	540	LEU
1	A	550	LEU
1	A	554	GLU
1	A	567	SER
1	A	569	ARG
1	A	570	LEU
1	B	15	ASP
1	B	207	ARG
1	B	310[A]	ARG
1	B	310[B]	ARG
1	B	335	GLN
1	B	381	LYS
1	B	410	GLN
1	B	411	ASP
1	B	432	ARG
1	B	448	VAL
1	B	491	ARG
1	B	518	LYS
1	B	579	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	91	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	549/591 (92%)	0.49	18 (3%) 46 39	52, 76, 111, 142	0
1	B	552/591 (93%)	0.48	12 (2%) 62 56	43, 60, 88, 125	0
All	All	1101/1182 (93%)	0.49	30 (2%) 54 48	43, 70, 104, 142	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	577	ILE	4.5
1	A	575	TRP	3.7
1	A	194	ILE	3.6
1	A	157	SER	3.6
1	A	164	ASP	3.1
1	A	261	ILE	3.0
1	B	309	CYS	3.0
1	A	110	GLU	3.0
1	A	163	VAL	2.5
1	B	212	LYS	2.5
1	B	310[A]	ARG	2.4
1	A	253	GLU	2.4
1	B	433	LEU	2.4
1	A	187	LEU	2.4
1	A	204	LEU	2.4
1	B	110	GLU	2.3
1	A	559	LEU	2.3
1	B	148	LEU	2.3
1	A	579	MET	2.2
1	B	187	LEU	2.2
1	B	410	GLN	2.1
1	B	308	PRO	2.1
1	A	572	LYS	2.1
1	B	437	LYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	153	LYS	2.1
1	A	450	ARG	2.1
1	A	564	GLU	2.1
1	B	213	LYS	2.1
1	B	166	SER	2.1
1	A	254	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.