



wwPDB EM Validation Summary Report ⓘ

Aug 5, 2024 – 08:52 AM EDT

PDB ID : 8E6Q
EMDB ID : EMD-27922
Title : Human TRPM2 ion channel in 1 mM ADPR
Authors : Wang, L.; Fu, T.M.; Xia, S.; Wu, H.
Deposited on : 2022-08-23
Resolution : 3.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev92
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1503	
1	B	1503	
1	C	1503	
1	D	1503	

2 Entry composition [i](#)


There are 2 unique types of molecules in this entry. The entry contains 40888 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transient receptor potential cation channel subfamily M member 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	1348	Total 10150	C 6560	N 1776	O 1772	S 42	0	0
1	B	1348	Total 10150	C 6560	N 1776	O 1772	S 42	0	0
1	C	1348	Total 10150	C 6560	N 1776	O 1772	S 42	0	0
1	D	1348	Total 10150	C 6560	N 1776	O 1772	S 42	0	0

- Molecule 2 is ADENOSINE-5-DIPHOSPHORIBOSE (three-letter code: APR) (formula: $C_{15}H_{23}N_5O_{14}P_2$).



Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
2	A	1	Total 36	C 15	N 5	O 14	P 2	0
2	A	1	Total 36	C 15	N 5	O 14	P 2	0

Continued on next page...


Continued from previous page...


Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
2	B	1	Total 36	15	5	14	2	0
2	B	1	Total 36	15	5	14	2	0
2	C	1	Total 36	15	5	14	2	0
2	C	1	Total 36	15	5	14	2	0
2	D	1	Total 36	15	5	14	2	0
2	D	1	Total 36	15	5	14	2	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transient receptor potential cation channel subfamily M member 2






• Molecule 1: Transient receptor potential cation channel subfamily M member 2






• Molecule 1: Transient receptor potential cation channel subfamily M member 2



Q381	K382	V385	F386	F387	Q388	E389	M390	F391	E392	T393	F394	T403	K404	D408	V410	R411	R412	R413	Q414	L415	L416	T417	V418	F419	R420	R421	G422	K423	D424	G425	Q426	Q427	D428	V429	D430	V431	A432	Q435	A436	L437	S441	R442	G448	H449	E450	M451	W452	Q455	A459										
W462	M463	R464	E471	M474	D475	E476	W477	Q478	M479	K480	P481	S482	L484	H485	P486	M488	M495	K496	V500	K501	L504	E512	L519	L526	L536	E542	D543	P544	E545	R546	P547	A550	P551	A552	A553	L556	Q557	M558	H559	H560	V561	A562	Q563	W564	L565														
R566	Q574	P575	L576	P578	R579	F580	R581	H582	N583	D584	R585	L586	R587	L588	L589	L590	F591	V592	P593	H594	V595	K596	L597	N598	V599	Q600	G601	V602	S603	L604	R605	S606	L607	V608	K609	R610	S611	S612	G613	H614	V615	T616	F617	T618	M619	R623	D624	L625	L626	I627	Q632	N633	R634	R635	E636				
I641	S645	C655	I658	L659	K660	E666	T669	D670	S671	L678	A679	E680	E692	K703	L704	R707	V708	S709	W712	G713	K714	T715	W730	S731	I735	T740	W743	W744	D750	N751	L758	C759	F763	P764	L765	L766	L767	T768	G769	L770																			
R774	E775	K776	R777	Q779	D780	W781	G782	A785	A786	R787	R789	A790	F791	F792	W795	V796	V797	F799	H800	L801	N802	L803	Y806	F807	L810	C811	A814	Y815	M818	V819	Q822	C830	A831	I832	Y833	F837	C841	E842	Q846	L847	F848	Y849	D850	P851	D852														
E853	C854	G855	L856	H857	K858	X859	D866	F867	W868	N869	K870	L871	D872	V873	G874	A875	L876	L878	F879	V880	A881	G882	L883	T884	C885	R886	P894	V897	L898	L899	S900	L901	D902	F903	Y904	C907	L908	R909	L910	F914	S917	K918	T919	L920	K923	L924	I925	V926	K928	R929									
M930	M931	X932	D933	V934	F935	F936	F937	L938	F939	L940	L941	A942	V943	V944	V945	S947	F948	V950	A951	K952	G953	A954	L955	L956	L957	H958	N959	E960	R961	R962	V963	D964	X965	L966	F967	R968	G969	A970	V971	Y972	H973	S974	V975	L976	V977	L978	F979	G980	Q981	L982	P983	G984	V985	D986	D987	G988	VAL		
ASN	PHE	ASN	PRO	GLU	HIS	CYS	SER	PRO	ASN	GLY	THR	ASP	PRO	TYR	LYS	LYS	LYS	CYS	PRO	GLU	SER	ASP	ALA	THR	GLN	GLN	ARG	PRO	ALA	F1020	P1021	E1022	W1023	L1024	T1025	V1026	L1027	L1028	L1029	C1030	L1031	Y1032	L1033	L1034	F1035	T1036	N1037	I1038	L1039	L1040	L1041	N1042	L1043	L1044	I1045	A1046	M1047	F1048	N1049
Y1050	T1051	F1052	Q1053	V1055	Q1056	E1057	H1058	T1059	D1060	K1064	R1067	E1073	Y1074	A1079	P1081	F1084	I1085	H1089	L1090	Q1091	L1092	F1093	I1094	V1097	V1098	L1099	K1100	T1101	P1102	A1103	K1104	H1106	K1112	K1115	K1129	E1130	N1131	Y1132	L1133	Q1134	N1135	R1136	Q1137	F1138	Q1139	Q1140													
K1141	Q1142	R1143	P1144	E1145	E1149	D1150	K1154	V1155	D1156	L1164	D1165	LEU	LEU	ARG	ARG	SER	GLY	SER	MET	GLU	GLN	ARG	LEU	ALA	SER	ALA	GLN	ALA	ALA	GLN	ALA	ALA	GLN	THR	ALA	GLN	LEU	ARG	THR	LEU	ARG	ALA	ALA	SER	GLY	PHE	SER	SER	GLU	ALA	ASP	VAL	PRO						
THR	LEU	ALA	SER	GLN	ALA	ALA	GLU	GLU	PRO	GLU	GLY	ARG	LYS	LYS	THR	GLU	GLU	PRO	G1235	D1236	S1237	V1240	M1241	A1242	R1243	H1244	L1245	L1246	M1249	R1254	F1255	P1256	V1257	P1258	M1259	E1260	K1261	P1263	M1264	E1265	T1266	E1267	F1268	L1269	I1270	Y1271	D1272	P1273	Y1276										
E1279	R1280	K1281	D1282	A1283	A1284	A1285	M1286	D1287	P1288	M1289	G1290	D1291	L1292	L1293	E1294	P1295	L1296	S1297	T1298	I1299	Q1300	V1301	V1304	S1367	G1368	K1369	K1370	M1371	L1372	E1373	V1374	K1378	L1381	S1382	W1385	G1389	E1393	P1394	G1395	E1396	M1397	R1400	K1401	R1404	I1405	L1406	R1407	M1422	E1423	K1426	G1427	Y1428	M1429	D1430	W1439	I1440			



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	135036	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	49	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	2.187	Depositor
Minimum map value	-1.194	Depositor
Average map value	0.005	Depositor
Map value standard deviation	0.059	Depositor
Recommended contour level	0.246	Depositor
Map size (\AA)	317.99997, 317.99997, 317.99997	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.06, 1.06, 1.06	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: APR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/10402	0.52	6/14191 (0.0%)
1	B	0.26	0/10402	0.52	6/14191 (0.0%)
1	C	0.26	0/10402	0.52	6/14191 (0.0%)
1	D	0.26	0/10402	0.52	6/14191 (0.0%)
All	All	0.26	0/41608	0.52	24/56764 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5
1	B	0	5
1	C	0	5
1	D	0	4
All	All	0	19

There are no bond length outliers.

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	930	MET	CB-CG-SD	6.88	133.03	112.40
1	C	930	MET	CB-CG-SD	6.87	133.02	112.40
1	A	930	MET	CB-CG-SD	6.87	133.02	112.40
1	D	930	MET	CB-CG-SD	6.86	132.97	112.40
1	D	1115	LYS	CB-CG-CD	5.99	127.17	111.60

There are no chirality outliers.

5 of 19 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	421	GLU	Peptide
1	A	422	GLY	Peptide
1	A	666	GLU	Peptide
1	A	669	THR	Peptide
1	A	713	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	10150	0	9654	242	0
1	B	10150	0	9654	243	0
1	C	10150	0	9654	250	0
1	D	10150	0	9654	250	0
2	A	72	0	42	2	0
2	B	72	0	42	3	0
2	C	72	0	42	3	0
2	D	72	0	42	3	0
All	All	40888	0	38784	965	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 965 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:930:MET:HG2	1:B:1039:LEU:HD11	1.43	0.98
1:A:1039:LEU:HD11	1:D:930:MET:HG2	1.48	0.96
1:B:930:MET:HG2	1:C:1039:LEU:HD11	1.53	0.88
1:C:930:MET:HG2	1:D:1039:LEU:HD11	1.60	0.84
1:C:1356:ARG:NH2	1:C:1371:MET:SD	2.54	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1342/1503 (89%)	1232 (92%)	107 (8%)	3 (0%)	47	78
1	B	1342/1503 (89%)	1231 (92%)	108 (8%)	3 (0%)	47	78
1	C	1342/1503 (89%)	1233 (92%)	106 (8%)	3 (0%)	47	78
1	D	1342/1503 (89%)	1231 (92%)	108 (8%)	3 (0%)	47	78
All	All	5368/6012 (89%)	4927 (92%)	429 (8%)	12 (0%)	50	78

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1455	VAL
1	B	1455	VAL
1	C	1455	VAL
1	D	1455	VAL
1	A	670	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	981/1318 (74%)	963 (98%)	18 (2%)	59	79
1	B	981/1318 (74%)	963 (98%)	18 (2%)	59	79
1	C	981/1318 (74%)	963 (98%)	18 (2%)	59	79
1	D	981/1318 (74%)	962 (98%)	19 (2%)	57	78
All	All	3924/5272 (74%)	3851 (98%)	73 (2%)	59	78

5 of 73 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	423	LYS
1	D	1149	GLU
1	D	636	GLU
1	D	923	LYS
1	B	703	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	1137	GLN
1	D	958	HIS
1	D	1131	ASN
1	D	953	GLN
1	B	1131	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	APR	A	1602	-	34,39,39	0.61	0	40,60,60	0.83	1 (2%)
2	APR	C	1601	-	34,39,39	0.61	0	40,60,60	0.82	2 (5%)
2	APR	C	1602	-	34,39,39	0.62	0	40,60,60	0.83	1 (2%)
2	APR	D	1601	-	34,39,39	0.61	0	40,60,60	0.82	2 (5%)
2	APR	B	1602	-	34,39,39	0.62	0	40,60,60	0.83	1 (2%)
2	APR	D	1602	-	34,39,39	0.61	0	40,60,60	0.83	1 (2%)
2	APR	A	1601	-	34,39,39	0.61	0	40,60,60	0.82	2 (5%)
2	APR	B	1601	-	34,39,39	0.62	0	40,60,60	0.82	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	APR	A	1602	-	-	4/18/54/54	0/4/4/4
2	APR	C	1601	-	-	1/18/54/54	0/4/4/4
2	APR	C	1602	-	-	4/18/54/54	0/4/4/4
2	APR	D	1601	-	-	1/18/54/54	0/4/4/4
2	APR	B	1602	-	-	4/18/54/54	0/4/4/4
2	APR	D	1602	-	-	4/18/54/54	0/4/4/4
2	APR	A	1601	-	-	1/18/54/54	0/4/4/4
2	APR	B	1601	-	-	1/18/54/54	0/4/4/4

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1601	APR	C5-C6-N6	2.31	123.87	120.35
2	A	1602	APR	C5-C6-N6	2.30	123.84	120.35
2	D	1601	APR	C5-C6-N6	2.30	123.84	120.35
2	A	1601	APR	C5-C6-N6	2.29	123.83	120.35
2	C	1602	APR	C5-C6-N6	2.28	123.82	120.35

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1602	APR	C5'-O5'-PA-O2A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	1602	APR	C5'-O5'-PA-O3A
2	B	1602	APR	C5'-O5'-PA-O2A
2	B	1602	APR	C5'-O5'-PA-O3A
2	C	1602	APR	C5'-O5'-PA-O2A




There are no ring outliers.




7 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1601	APR	2	0
2	C	1602	APR	1	0
2	D	1601	APR	2	0
2	B	1602	APR	1	0
2	D	1602	APR	1	0
2	A	1601	APR	2	0
2	B	1601	APR	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-27922. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 147




Y Index: 153




Z Index: 157


6.3.2 Raw map



X Index: 147



Y Index: 153




Z Index: 157


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.246. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 322 nm³; this corresponds to an approximate mass of 291 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)




*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.08	6.67	4.19

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.08 differs from the reported value 3.4 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-27922 and PDB model 8E6Q. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.246 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.246).











9.4 Atom inclusion [i](#)



At the recommended contour level, 70% of all backbone atoms, 59% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.246) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5930	 0.2510
A	 0.6030	 0.2580
B	 0.5890	 0.2480
C	 0.5910	 0.2470
D	 0.5900	 0.2520

