



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2023 – 09:15 PM EDT

PDB ID : 3E7B
Title : Crystal Structure of Protein Phosphatase-1 Bound to the natural toxin inhibitor Tautomycin
Authors : Kelker, M.S.; Page, R.; Peti, W.
Deposited on : 2008-08-18
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	299	 4% 91% 7% ..
1	B	299	 6% 90% 7% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	AZI	A	2	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 5370 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein phosphatase PP1-alpha catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	294	2382	1535	393	434	20	0	12	0
1	B	294	2361	1529	386	427	19	0	11	0


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP P62136
A	-3	HIS	-	expression tag	UNP P62136
A	-2	MET	-	expression tag	UNP P62136
A	-1	GLY	-	expression tag	UNP P62136
A	0	SER	-	expression tag	UNP P62136
B	-4	GLY	-	expression tag	UNP P62136
B	-3	HIS	-	expression tag	UNP P62136
B	-2	MET	-	expression tag	UNP P62136
B	-1	GLY	-	expression tag	UNP P62136
B	0	SER	-	expression tag	UNP P62136

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
2	A	2	2	2	0	0
2	B	2	2	2	0	0

- Molecule 3 is (2Z)-2-[(1R)-3-[[[(1R,2S,3R,6S,7S,10R)-10-{(2S,3S,6R,8S,9R)-3,9-dimethyl-8-[(3S)-3-methyl-4-oxopentyl]-1,7-dioxaspiro[5.5]undec-2-yl}]-3,7-dihydroxy-2-methoxy-6-methyl-1-(1-methylethyl)-5-oxoundecyl]oxy]-1-hydroxy-3-oxopropyl]-3-methylbut-2-enedioic acid (three-letter code: E7B) (formula: C₄₁H₆₈O₁₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			55	41	14		
3	B	1	Total	C	O	0	0
			55	41	14		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Na 1 1	0	0

- Molecule 7 is AZIDE ION (three-letter code: AZI) (formula: N₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total N 3 3	0	0
7	A	1	Total N 3 3	0	0
7	A	1	Total N 3 3	0	0
7	A	1	Total N 3 3	0	0
7	B	1	Total N 3 3	0	0
7	B	1	Total N 3 3	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	283	Total 283	O 283	0	0
8	B	204	Total 204	O 204	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit



- Molecule 1: Serine/threonine-protein phosphatase PP1-alpha catalytic subunit



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.76Å 78.52Å 130.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.63 – 1.70 19.63 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.2 (19.63-1.70) 99.2 (19.63-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.81 (at 1.70Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.153 , 0.175 0.159 , 0.178	Depositor DCC
R_{free} test set	3750 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	22.5	Xtrriage
Anisotropy	0.079	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5370	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AZI, NA, GOL, CL, E7B, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.60	0/2472	0.72	3/3343 (0.1%)
1	B	0.57	0/2449	0.70	2/3317 (0.1%)
All	All	0.58	0/4921	0.71	5/6660 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0
1	B	1	0
All	All	2	0

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	23	ARG	C-N-CD	-5.39	108.73	120.60
1	A	7	LEU	CA-CB-CG	5.31	127.50	115.30
1	A	36[A]	ARG	CG-CD-NE	-5.19	100.91	111.80
1	A	36[B]	ARG	CG-CD-NE	-5.19	100.91	111.80
1	B	23	ARG	N-CA-C	5.03	124.58	111.00

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	273	CYS	CA
1	B	23	ARG	CA

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2382	0	2349	14	0
1	B	2361	0	2323	21	0
2	A	2	0	0	0	0
2	B	2	0	0	1	0
3	A	55	0	66	1	0
3	B	55	0	66	3	0
4	A	1	0	0	0	0
5	A	6	0	8	1	0
6	A	1	0	0	0	0
7	A	12	0	0	0	0
7	B	6	0	0	0	0
8	A	283	0	0	1	0
8	B	204	0	0	5	0
All	All	5370	0	4812	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139[A]:GLU:OE1	8:B:714:HOH:O	1.69	1.10
1:B:153[A]:THR:HG22	8:B:715:HOH:O	1.85	0.77
1:B:153[A]:THR:CG2	8:B:715:HOH:O	2.35	0.73
1:A:32:GLU:O	1:A:36[A]:ARG:HG3	1.91	0.71
3:B:500:E7B:H29A	3:B:500:E7B:H21	1.72	0.70
1:B:36:ARG:O	1:B:40[B]:LEU:HD13	1.92	0.68
1:A:116[B]:GLU:OE2	5:A:501:GOL:O3	2.13	0.66
1:B:24:PRO:CB	1:B:25:GLY:HA2	2.25	0.66
1:B:62[A]:CYS:SG	1:B:65:ILE:HD11	2.36	0.65
1:B:24:PRO:HB3	1:B:25:GLY:HA2	1.77	0.65
2:B:401:MN:MN	8:B:700:HOH:O	1.57	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:195:VAL:HB	3:B:500:E7B:H1A	1.83	0.60
1:A:62[B]:CYS:SG	1:A:65:ILE:HD11	2.45	0.57
1:B:23:ARG:CB	1:B:24:PRO:O	2.54	0.55
1:A:70:TYR:HE1	1:A:99:GLN:HE22	1.56	0.54
1:B:161[B]:ILE:HG21	1:B:201:LEU:HD11	1.91	0.53
1:A:281:ALA:HB2	1:A:295[B]:ILE:CD1	2.40	0.52
1:A:58:PRO:HD3	1:A:287:GLU:HG3	1.91	0.52
1:A:9:LEU:HD11	1:A:112:ILE:HG22	1.93	0.50
1:A:113:LYS:HE2	8:A:760:HOH:O	2.14	0.47
1:B:161[B]:ILE:HG21	1:B:201:LEU:CD1	2.45	0.46
1:A:236:LEU:HD21	1:A:244:ILE:HG13	1.98	0.46
1:B:60:LYS:NZ	1:B:284[B]:SER:OG	2.49	0.46
1:A:281:ALA:CB	1:A:295[B]:ILE:HD13	2.46	0.45
1:B:24:PRO:CB	1:B:25:GLY:CA	2.93	0.45
1:B:18:GLU:OE2	8:B:631:HOH:O	2.21	0.45
1:A:181:GLN:N	1:A:185:GLN:OE1	2.42	0.45
1:B:210:ASP:HB3	1:B:213:VAL:HG12	1.99	0.44
1:A:281:ALA:CB	1:A:295[B]:ILE:CD1	2.96	0.44
1:A:236:LEU:CD1	1:A:262:GLN:HB3	2.48	0.43
1:B:195:VAL:O	3:B:500:E7B:C1	2.67	0.42
3:A:500:E7B:H29B	3:A:500:E7B:H24	1.90	0.42
1:B:161[B]:ILE:CD1	1:B:201:LEU:HD13	2.49	0.42
1:B:51:ILE:HD13	1:B:189:ILE:HB	2.02	0.41
1:B:255:TYR:HA	1:B:265:THR:O	2.20	0.41
1:B:161[A]:ILE:HD12	1:B:201:LEU:HD11	2.02	0.41
1:A:255:TYR:HA	1:A:265:THR:O	2.21	0.40
1:B:208:ASP:O	1:B:226:THR:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	304/299 (102%)	294 (97%)	10 (3%)	0	100	100
1	B	303/299 (101%)	288 (95%)	13 (4%)	2 (1%)	22	8
All	All	607/598 (102%)	582 (96%)	23 (4%)	2 (0%)	41	24

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	23	ARG
1	B	24	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/262 (99%)	258 (99%)	2 (1%)	81	74
1	B	255/262 (97%)	254 (100%)	1 (0%)	91	87
All	All	515/524 (98%)	512 (99%)	3 (1%)	86	80

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	113	LYS
1	A	246	ARG
1	B	246	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	99	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 6 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	E7B	B	500	-	54,56,56	1.14	3 (5%)	68,80,80	1.24	7 (10%)
5	GOL	A	501	-	5,5,5	0.40	0	5,5,5	0.22	0
3	E7B	A	500	-	54,56,56	1.22	5 (9%)	68,80,80	1.72	6 (8%)
7	AZI	B	2	-	0,2,2	-	-	0,1,1	-	-
7	AZI	A	504	-	0,2,2	-	-	0,1,1	-	-
7	AZI	B	501	-	0,2,2	-	-	0,1,1	-	-
7	AZI	A	505	-	0,2,2	-	-	0,1,1	-	-
7	AZI	A	2	-	0,2,2	-	-	0,1,1	-	-
7	AZI	A	503	-	0,2,2	-	-	0,1,1	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	501	-	-	4/4/4/4	-
3	E7B	B	500	-	-	9/66/98/98	0/2/2/2
3	E7B	A	500	-	-	12/66/98/98	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	500	E7B	C7'-C4'	3.52	1.54	1.48
3	B	500	E7B	C7'-C4'	3.33	1.54	1.48
3	A	500	E7B	C24-C23	3.27	1.58	1.52
3	B	500	E7B	O6-C23	3.22	1.50	1.42
3	B	500	E7B	C24-C23	3.13	1.58	1.52
3	A	500	E7B	O6-C23	3.06	1.50	1.42
3	A	500	E7B	O9-C7'	-2.06	1.24	1.30
3	A	500	E7B	C2'-C1'	2.02	1.54	1.50

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	E7B	O7-C24-C31	-9.07	91.66	107.31
3	A	500	E7B	C31-C24-C23	7.09	123.71	113.55
3	B	500	E7B	O7-C24-C31	-4.60	99.37	107.31
3	B	500	E7B	O7-C1'-C2'	3.21	117.35	111.46
3	A	500	E7B	C17-C18-C19	-2.74	109.48	113.25
3	B	500	E7B	C31-C24-C23	2.74	117.47	113.55
3	B	500	E7B	C17-C18-C19	-2.56	109.73	113.25
3	A	500	E7B	O7-C1'-C2'	2.56	116.15	111.46
3	B	500	E7B	O12-C1'-C2'	-2.32	119.59	124.73
3	B	500	E7B	C17-C16-C15	-2.28	109.38	115.34
3	B	500	E7B	O10-C6'-C5'	-2.23	116.66	122.64
3	A	500	E7B	C24-O7-C1'	2.13	121.08	117.53
3	A	500	E7B	O12-C1'-C2'	-2.01	120.28	124.73

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	500	E7B	C24-C23-O6-C29
3	A	500	E7B	C23-C24-C31-C33
3	A	500	E7B	C23-C24-C31-C32
3	A	500	E7B	O7-C24-C31-C33
3	A	500	E7B	O7-C24-C31-C32
3	A	500	E7B	C41-C5'-C6'-O10
3	A	500	E7B	C41-C5'-C6'-O8
3	B	500	E7B	C22-C23-O6-C29
3	B	500	E7B	C23-C24-C31-C33
3	B	500	E7B	C23-C24-C31-C32
3	B	500	E7B	O7-C24-C31-C33
3	B	500	E7B	O7-C24-C31-C32

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	500	E7B	C41-C5'-C6'-O10
3	B	500	E7B	C41-C5'-C6'-O8
5	A	501	GOL	O1-C1-C2-O2
5	A	501	GOL	O1-C1-C2-C3
5	A	501	GOL	C1-C2-C3-O3
5	A	501	GOL	O2-C2-C3-O3
3	A	500	E7B	C1'-C2'-C3'-O11
3	B	500	E7B	C1'-C2'-C3'-O11
3	A	500	E7B	C22-C23-C24-C31
3	A	500	E7B	C1-C2-C3-C30
3	A	500	E7B	C4'-C5'-C6'-O10
3	B	500	E7B	C4'-C5'-C6'-O10
3	A	500	E7B	O3-C2-C3-C30

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	500	E7B	3	0
5	A	501	GOL	1	0
3	A	500	E7B	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	294/299 (98%)	-0.11	11 (3%) 41 46	17, 22, 35, 46	8 (2%)
1	B	294/299 (98%)	0.23	17 (5%) 23 25	17, 26, 37, 57	10 (3%)
All	All	588/598 (98%)	0.06	28 (4%) 30 34	17, 24, 36, 57	18 (3%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	23	ARG	6.9
1	B	0	SER	6.9
1	B	24	PRO	6.5
1	B	176	LEU	4.6
1	A	21	GLY	4.1
1	A	23	ARG	3.8
1	B	237	HIS	3.7
1	B	22	SER	3.6
1	B	215	GLY	3.5
1	B	25	GLY	3.4
1	A	24	PRO	3.4
1	B	273	CYS	3.3
1	B	214	GLN	3.0
1	A	176	LEU	3.0
1	A	213	VAL	2.7
1	B	7	LEU	2.7
1	A	273	CYS	2.7
1	B	21	GLY	2.6
1	B	190	MET	2.5
1	A	212	ASP	2.4
1	A	22[A]	SER	2.1
1	B	213	VAL	2.1
1	A	0	SER	2.1
1	A	170	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	274	GLY	2.1
1	A	237	HIS	2.1
1	B	170	PHE	2.1
1	B	204	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	AZI	A	2	3/3	0.57	0.64	44,44,52,59	0
7	AZI	B	501	3/3	0.77	0.16	63,63,67,71	0
7	AZI	A	505	3/3	0.81	0.23	47,47,51,53	0
4	CL	A	1	1/1	0.85	0.86	48,48,48,48	0
7	AZI	B	2	3/3	0.88	0.21	56,56,57,59	0
3	E7B	B	500	55/55	0.89	0.16	25,31,43,53	55
7	AZI	A	504	3/3	0.91	0.19	41,41,52,54	0
3	E7B	A	500	55/55	0.91	0.15	24,31,37,47	0
7	AZI	A	503	3/3	0.92	0.21	34,34,45,48	0
5	GOL	A	501	6/6	0.96	0.17	31,46,52,53	0
2	MN	B	400	1/1	0.98	0.07	23,23,23,23	1
2	MN	B	401	1/1	0.98	0.08	23,23,23,23	0
2	MN	A	400	1/1	0.99	0.08	19,19,19,19	0
2	MN	A	401	1/1	0.99	0.06	19,19,19,19	0
6	NA	A	502	1/1	0.99	0.10	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different

orientation to approximate a three-dimensional view.




Electron density around E7B B 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around E7B A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.