

Full wwPDB X-ray Structure Validation Report (i)

Nov 21, 2023 – 12:57 AM JST

PDB ID	:	7EBW
Title	:	Crystal structure of Aedes aegypti Noppera-bo, glutathione S-transferase ep-
		silon 8, in desmethylglycitein and glutathione-bound form
Authors	:	Inaba, K.; Koiwai, K.; Senda, M.; Senda, T.; Niwa, R.
Deposited on	:	2021-03-11
Resolution	:	1.94 Å(reported)
resolution	·	1.94 A(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity Mogul Xtriage (Phenix) EDS	:	4.02b-467 1.8.5 (274361), CSD as541be (2020) 1.13 2.36
buster-report Percentile statistics Refmac	: : :	1.1.7 (2018) 20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Percentile relative to X-ray structures of similar re	solution
---	----------

Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$		
R _{free}	130704	4310 (1.96-1.92)		
Clashscore	141614	1023 (1.94-1.94)		
Ramachandran outliers	138981	1007 (1.94-1.94)		
Sidechain outliers	138945	1007 (1.94-1.94)		
RSRZ outliers	127900	4250 (1.96-1.92)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	227	87%	10%	·
1	В	227	% 94%		•••
1	С	227	85%	10%	5%
1	D	227	89%	7%	•



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7670 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	٨	210	Total	С	Ν	0	\mathbf{S}	0	2	0
	А	219	1701	1100	286	302	13	0		0
1	D	221	Total	С	Ν	0	S	0	3	0
	ГБ	221	1705	1099	289	302	15	0		0
1	С	216	Total	С	Ν	0	S	0	1	0
		210	1668	1077	282	297	12	0	1	0
1	1 D	210	Total	С	Ν	0	S	0	9	0
		219	1689	1086	285	304	14	0	Δ	0

• Molecule 1 is a protein called Glutathione transferase.


There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-6	MET	-	initiating methionine	UNP A0A1S4FIB3
А	-5	ASN	-	expression tag	UNP A0A1S4FIB3
A	-4	HIS	-	expression tag	UNP A0A1S4FIB3
А	-3	LYS	-	expression tag	UNP A0A1S4FIB3
А	-2	VAL	-	expression tag	UNP A0A1S4FIB3
А	-1	HIS	-	expression tag	UNP A0A1S4FIB3
A	0	MET	-	expression tag	UNP A0A1S4FIB3
В	-6	MET	-	initiating methionine	UNP A0A1S4FIB3
В	-5	ASN	-	expression tag	UNP A0A1S4FIB3
В	-4	HIS	-	expression tag	UNP A0A1S4FIB3
В	-3	LYS	-	expression tag	UNP A0A1S4FIB3
В	-2	VAL	-	expression tag	UNP A0A1S4FIB3
В	-1	HIS	-	expression tag	UNP A0A1S4FIB3
В	0	MET	-	expression tag	UNP A0A1S4FIB3
С	-6	MET	-	initiating methionine	UNP A0A1S4FIB3
C	-5	ASN	-	expression tag	UNP A0A1S4FIB3
С	-4	HIS	-	expression tag	UNP A0A1S4FIB3
С	-3	LYS	-	expression tag	UNP A0A1S4FIB3
С	-2	VAL	-	expression tag	UNP A0A1S4FIB3
С	-1	HIS	-	expression tag	UNP A0A1S4FIB3
С	0	MET	-	expression tag	UNP A0A1S4FIB3



Chain	Residue	Modelled	Actual	Comment	Reference
D	-6	MET	-	initiating methionine	UNP A0A1S4FIB3
D	-5	ASN	-	expression tag	UNP A0A1S4FIB3
D	-4	HIS	-	expression tag	UNP A0A1S4FIB3
D	-3	LYS	-	expression tag	UNP A0A1S4FIB3
D	-2	VAL	-	expression tag	UNP A0A1S4FIB3
D	-1	HIS	-	expression tag	UNP A0A1S4FIB3
D	0	MET	-	expression tag	UNP A0A1S4FIB3


• Molecule 2 is GLUTATHIONE (three-letter code: GSH) (formula: $C_{10}H_{17}N_3O_6S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	\mathbf{ms}			ZeroOcc	AltConf	
2	Δ	1	Total	С	Ν	0	\mathbf{S}	0	0	
	Π	T	20	10	3	6	1	0	0	
2	2 B	В	1	Total	С	Ν	0	S	0	0
		1	20	10	3	6	1	0	0	
2	С	1	Total	С	Ν	0	S	0	0	
		1	20	10	3	6	1	0		
2	Л	1	Total	С	Ν	Ο	S	0	0	
				10	3	6	1	0	0	

• Molecule 3 is 6,7-dihydroxy-3-(4-hydroxyphenyl)-4H-chromen-4-one (three-letter code: 674) (formula: $C_{15}H_{10}O_5$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C O 20 15 5	0	0
3	В	1	Total C O 20 15 5	0	0
3	С	1	Total C O 20 15 5	0	0
3	D	1	Total C O 20 15 5	0	0

• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	2	Total Ca 2 2	0	0
4	В	1	Total Ca 1 1	0	0
4	С	3	Total Ca 3 3	0	0
4	D	2	Total Ca 2 2	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	204	Total O 204 204	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	185	Total O 185 185	0	0
5	С	182	Total O 182 182	0	0
5	D	168	Total O 168 168	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Glutathione transferase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants	151.31Å 151.31Å 146.15Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.85 - 1.94	Depositor
Resolution (A)	47.85 - 1.94	EDS
% Data completeness	98.0 (47.85-1.94)	Depositor
(in resolution range)	$98.1 \ (47.85 - 1.94)$	EDS
R _{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.48 (at 1.94 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.19_4092	Depositor
D D	0.172 , 0.219	Depositor
R, R_{free}	0.172 , 0.218	DCC
R_{free} test set	2960 reflections (4.83%)	wwPDB-VP
Wilson B-factor $(Å^2)$	27.4	Xtriage
Anisotropy	0.114	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34, 64.4	EDS
L-test for twinning ²	$< L > = 0.50, < L^2 > = 0.33$	Xtriage
Estimated twinning fraction	0.000 for l,-k,h	Vtriago
Estimated twinning fraction	0.000 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7670	wwPDB-VP
Average B, all atoms $(Å^2)$	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 43.51 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.7457e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GSH, 674

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.39	0/1738	0.60	0/2367	
1	В	0.39	0/1742	0.58	0/2374	
1	С	0.40	0/1704	0.59	0/2325	
1	D	0.40	0/1724	0.60	0/2351	
All	All	0.40	0/6908	0.59	0/9417	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1701	0	1666	13	0
1	В	1705	0	1648	5	0
1	С	1668	0	1633	12	0
1	D	1689	0	1641	9	0
2	А	20	0	15	0	0
2	В	20	0	15	0	0
2	С	20	0	15	0	0
2	D	20	0	15	0	0
3	А	20	0	0	0	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	20	0	0	0	0
3	С	20	0	0	0	0
3	D	20	0	0	0	0
4	А	2	0	0	0	0
4	В	1	0	0	0	0
4	С	3	0	0	0	0
4	D	2	0	0	0	0
5	А	204	0	0	0	0
5	В	185	0	0	0	0
5	С	182	0	0	1	0
5	D	168	0	0	2	0
All	All	7670	0	6648	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:GLY:HA3	1:A:125:ILE:HD12	1.81	0.62
1:A:78:CYS:SG	1:A:160:VAL:HG11	2.39	0.61
1:B:119:GLY:HA3	1:B:125:ILE:HG13	1.82	0.61
1:B:78:CYS:SG	1:B:160:VAL:HG11	2.41	0.61
1:C:134:LYS:O	1:C:138:GLU:HG2	2.05	0.56
1:C:171:THR:O	1:C:174:MET:HG2	2.06	0.56
1:A:117:MET:SD	1:A:214:LEU:HD11	2.48	0.53
1:A:174:MET:CE	1:A:214:LEU:HD12	2.39	0.53
1:A:109:LEU:HG	1:A:168[B]:CYS:SG	2.48	0.53
1:C:109:LEU:HG	1:C:168[B]:CYS:SG	2.51	0.51
1:C:27:ILE:HG21	1:C:81:PHE:HB3	1.93	0.50
1:A:7:TYR:CE2	1:A:34:LYS:HD3	2.47	0.49
1:D:178:LEU:HD11	1:D:186:THR:HG22	1.95	0.48
1:C:9:ASP:OD1	1:C:11:ILE:HG12	2.14	0.47
1:D:165:ILE:O	1:D:169:MET:HG2	2.15	0.46
1:D:97[B]:ARG:NH1	5:D:407:HOH:O	2.48	0.46
1:A:178:LEU:HD21	1:A:186:THR:HG22	1.97	0.46
1:C:129:GLU:HA	5:C:546:HOH:O	2.16	0.45
1:C:76:TYR:CZ	1:C:80:ILE:HD13	2.51	0.45
1:C:178:LEU:HD21	1:C:186:THR:HG22	1.98	0.45
1:B:109:LEU:HG	1:B:168[B]:CYS:SG	2.57	0.44
1:D:9:ASP:HB2	1:D:36:VAL:O	2.18	0.44



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:175:VAL:C	1:D:177:PRO:HD3	2.39	0.43
1:A:119:GLY:HA3	1:A:125:ILE:CD1	2.48	0.43
1:C:166:VAL:HG21	1:C:193:MET:HB3	2.00	0.42
1:A:28:LYS:HE3	1:A:28:LYS:HB3	1.86	0.42
1:A:22:VAL:HG13	1:A:27:ILE:HG13	2.01	0.42
1:C:9:ASP:CG	1:C:11:ILE:HG12	2.40	0.41
1:C:78:CYS:SG	1:C:160:VAL:HG11	2.60	0.41
1:D:40:GLU:OE2	5:D:401:HOH:O	2.21	0.41
1:C:116:VAL:HG13	1:C:128:LEU:HD21	2.02	0.41
1:A:80:ILE:HG13	1:A:81:PHE:CD2	2.56	0.41
1:D:13:PRO:HB2	1:D:14:PRO:HD3	2.03	0.41
1:D:9:ASP:H	1:D:15:VAL:HG21	1.86	0.41
1:D:114:SER:O	1:D:118:ARG:HB2	2.21	0.41
1:A:16:ARG:HD2	1:A:203:ASN:OD1	2.20	0.40
1:B:174:MET:HE3	1:B:214:LEU:HD22	2.03	0.40
1:A:114[A]:SER:O	1:A:118:ARG:HB2	2.21	0.40
1:B:119:GLY:CA	1:B:125:ILE:HG13	2.49	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	219/227~(96%)	215~(98%)	4(2%)	0	100	100
1	В	222/227~(98%)	217~(98%)	5(2%)	0	100	100
1	С	215/227~(95%)	211 (98%)	4(2%)	0	100	100
1	D	219/227~(96%)	213~(97%)	6 (3%)	0	100	100
All	All	875/908~(96%)	856~(98%)	19~(2%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	А	177/206~(86%)	177~(100%)	0	100	100
1	В	175/206~(85%)	175 (100%)	0	100	100
1	С	175/206~(85%)	174 (99%)	1 (1%)	86	85
1	D	176/206~(85%)	174 (99%)	2(1%)	73	67
All	All	703/824~(85%)	700 (100%)	3~(0%)	91	91

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	88	SER
1	D	178	LEU
1	D	181	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	111	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Chain		Chain	ain Res Link		Bo	Bond lengths			Bond angles		
MOI	Iol Type Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2		
3	674	А	302	-	22,22,22	<mark>3.79</mark>	5 (22%)	32,32,32	1.71	<mark>6 (18%)</mark>	
2	GSH	С	301	-	18,19,19	0.72	0	23,24,24	0.74	0	
3	674	D	302	-	22,22,22	3.91	4 (18%)	32,32,32	1.70	8 (25%)	
3	674	С	302	-	22,22,22	4.02	6 (27%)	32,32,32	1.80	8 (25%)	
3	674	В	302	-	22,22,22	3.84	6 (27%)	32,32,32	1.43	7 (21%)	
2	GSH	В	301	-	18,19,19	0.66	0	23,24,24	0.82	0	
2	GSH	D	301	-	18,19,19	0.72	0	23,24,24	0.68	0	
2	GSH	А	301	-	18,19,19	0.67	0	23,24,24	0.84	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	674	А	302	-	-	0/4/4/4	0/3/3/3
2	GSH	С	301	-	-	0/24/24/24	-
3	674	D	302	-	-	0/4/4/4	0/3/3/3
3	674	С	302	-	-	0/4/4/4	0/3/3/3
3	674	В	302	-	-	0/4/4/4	0/3/3/3
2	GSH	В	301	-	-	1/24/24/24	-
2	GSH	D	301	-	-	1/24/24/24	-
2	GSH	А	301	-	-	1/24/24/24	-

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	302	674	C2-C3	12.86	1.49	1.34
3	С	302	674	C2-C3	12.58	1.48	1.34
3	В	302	674	C2-C3	12.02	1.48	1.34



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	А	302	674	C2-C3	11.46	1.47	1.34
3	С	302	674	01-C8A	11.30	1.51	1.37
3	А	302	674	O1-C8A	10.50	1.50	1.37
3	D	302	674	01-C8A	10.26	1.50	1.37
3	В	302	674	01-C8A	10.08	1.50	1.37
3	В	302	674	O1-C2	6.38	1.46	1.35
3	А	302	674	O1-C2	6.36	1.46	1.35
3	С	302	674	O1-C2	6.06	1.45	1.35
3	D	302	674	O1-C2	5.68	1.45	1.35
3	D	302	674	C3-C4	-3.18	1.42	1.46
3	А	302	674	C1'-C3	3.07	1.53	1.48
3	В	302	674	C1'-C3	3.00	1.53	1.48
3	С	302	674	C1'-C3	2.80	1.52	1.48
3	В	302	674	C3-C4	-2.70	1.42	1.46
3	А	302	674	C3-C4	-2.55	1.43	1.46
3	С	302	674	C3-C4	-2.30	1.43	1.46
3	В	302	674	O3-C6	2.11	1.40	1.36
3	С	302	674	O3-C6	2.08	1.40	1.36

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	Observed(°)	$Ideal(^{o})$
3	С	302	674	C1'-C3-C2	-5.32	114.27	119.72
3	А	302	674	C1'-C3-C2	-4.71	114.89	119.72
3	D	302	674	O1-C2-C3	-4.58	121.21	125.81
3	С	302	674	O1-C2-C3	-4.06	121.73	125.81
3	А	302	674	C8A-O1-C2	-3.67	114.39	118.28
3	D	302	674	C1'-C3-C2	-3.57	116.06	119.72
3	В	302	674	C8A-O1-C2	-3.35	114.73	118.28
3	В	302	674	O1-C2-C3	-3.35	122.45	125.81
3	D	302	674	C2'-C1'-C3	-3.27	116.36	120.98
3	D	302	674	C2-C3-C4	3.08	121.56	118.60
3	С	302	674	C8A-O1-C2	-3.01	115.09	118.28
3	В	302	674	C1'-C3-C2	-2.98	116.67	119.72
3	А	302	674	O1-C2-C3	-2.97	122.83	125.81
3	А	302	674	C4A-C4-C3	2.93	118.94	115.35
3	А	302	674	C1'-C3-C4	2.92	124.74	121.65
3	С	302	674	C1'-C3-C4	2.88	124.70	121.65
3	С	302	674	C4A-C4-C3	2.81	118.79	115.35
3	А	302	674	O1-C8A-C4A	2.71	123.22	121.50
3	С	302	674	O1-C8A-C4A	2.54	123.12	121.50
3	С	302	674	C2-C3-C4	2.52	121.03	118.60



Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
3	D	302	674	C3'-C2'-C1'	-2.49	117.88	120.78
3	В	302	674	C4A-C4-C3	2.45	118.34	115.35
3	В	302	674	O1-C8A-C4A	2.42	123.05	121.50
3	D	302	674	C6'-C1'-C2'	2.37	121.97	118.59
3	D	302	674	C8A-O1-C2	-2.32	115.82	118.28
3	В	302	674	C2-C3-C4	2.27	120.78	118.60
3	В	302	674	O2-C4-C3	-2.26	120.16	122.51
3	С	302	674	C6'-C1'-C3	-2.13	117.98	120.98
3	D	302	674	C4A-C4-C3	2.08	117.89	115.35

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	301	GSH	C3-CA3-N3-C2
2	В	301	GSH	C3-CA3-N3-C2
2	А	301	GSH	C3-CA3-N3-C2

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2		$OWAB(Å^2)$	Q < 0.9
1	А	219/227~(96%)	-0.20	1 (0%) 91	93	18, 28, 54, 65	0
1	В	221/227~(97%)	-0.26	2 (0%) 84	87	19, 28, 47, 61	0
1	С	216/227~(95%)	-0.18	1 (0%) 91	93	20, 31, 48, 67	0
1	D	219/227~(96%)	-0.16	1 (0%) 91	93	20, 31, 52, 67	0
All	All	875/908~(96%)	-0.20	5 (0%) 89	92	18, 29, 51, 67	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	176[A]	CYS	2.8
1	В	131	HIS	2.7
1	А	0	MET	2.3
1	В	127	THR	2.1
1	С	130	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CA	С	303	1/1	0.90	0.10	60,60,60,60	0
4	CA	А	304	1/1	0.94	0.07	$53,\!53,\!53,\!53$	0
3	674	D	302	20/20	0.94	0.10	24,31,35,35	0
4	CA	С	304	1/1	0.94	0.07	49,49,49,49	0
3	674	В	302	20/20	0.96	0.08	20,27,29,33	0
4	CA	В	303	1/1	0.96	0.11	47,47,47,47	0
3	674	С	302	20/20	0.96	0.09	22,29,34,40	0
3	674	А	302	20/20	0.96	0.09	20,26,30,30	0
2	GSH	С	301	20/20	0.97	0.09	18,24,29,33	0
4	CA	А	303	1/1	0.97	0.09	52,52,52,52	0
2	GSH	В	301	20/20	0.98	0.08	18,22,27,34	0
2	GSH	А	301	20/20	0.98	0.09	18,21,33,33	0
2	GSH	D	301	20/20	0.98	0.09	18,25,31,43	0
4	CA	D	304	1/1	0.98	0.04	38,38,38,38	0
4	CA	D	303	1/1	0.99	0.04	31,31,31,31	0
4	CA	С	305	1/1	0.99	0.04	40,40,40,40	0


median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





















6.5 Other polymers (i)

There are no such residues in this entry.

