

wwPDB X-ray Structure Validation Summary Report (i)

Nov 21, 2023 – 01:51 AM JST

PDB ID : 7EDB

Title: EcoT38I restriction endonuclease complexed with DNA

Authors: Kita, K.; Mikami, B.

Deposited on : 2021-03-15

Resolution : 2.39 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},\ {\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	351	94%	5%
1	В	351	5% 91%	7% ••
2	Е	13	92%	8%
3	F	13	85%	8% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-



ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	TRS	В	405[A]	-	X	-	-
5	TRS	В	405[B]	-	X	-	-



2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 6405 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called EcoT38I restriction endonuclease.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	Δ	350	Total	С	N	О	S	0	1	0
1	Λ	330	2736	1741	471	516	8	0		
1	В	347	Total	С	N	О	S	0	2	0
1	Ъ	941	2721	1733	465	515	8	0	3	

• Molecule 2 is a DNA chain called DNA (5'-D(*GP*GP*CP*AP*GP*AP*GP*CP*TP*CP* AP*CP*G)-3').

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
2	Е	13	Total 285	C 136	N 59	O 78	P 12	0	1	0

• Molecule 3 is a DNA chain called DNA (5'-D(P*CP*GP*TP*GP*AP*GP*CP*TP*CP*TP *GP*C)-3').


\mathbf{N}	Iol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
	3	F	12	Total 245	C 116	N 43	O 74	P 12	0	0	0

• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Ca 2 2	0	0
4	В	1	Total Ca 1 1	0	0
4	F	1	Total Ca 1 1	0	0


• Molecule 5 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: C₄H₁₂NO₃) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 9 4 1 4	0	1
5	A	1	Total C N O 8 4 1 3	0	0
5	В	1	Total C N O 8 4 1 3	0	0
5	В	1	Total C N O 9 4 1 4	0	1

• Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	A	1	Total C O 6 3 3	0	0
6	В	1	Total C O 6 3 3	0	0
6	В	1	Total C O 6 3 3	0	0
6	В	1	Total C O 6 3 3	0	0
6	Е	1	Total C O 6 3 3	0	0

• Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total Cl 1 1	0	0
7	F	1	Total Cl 1 1	0	0

• Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	В	1	Total Na 1 1	0	0

• Molecule 9 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	В	1	Total 10	C 6	O 4	0	0

• Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	135	Total O 135 135	0	0
10	В	119	Total O 119 119	0	0
10	Е	31	Total O 31 31	0	0
10	F	34	Total O 34 34	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: EcoT38I restriction endonuclease Chain A: 94% • Molecule 1: EcoT38I restriction endonuclease Chain B: 91% 7% ... • Molecule 2: DNA (5'-D(*GP*GP*CP*AP*GP*AP*GP*CP*TP*CP*AP*CP*G)-3') Chain E: 92% 8% • Molecule 3: DNA (5'-D(P*CP*GP*TP*GP*AP*GP*CP*TP*CP*TP*GP*C)-3') Chain F: 85%



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 64 2 2	Depositor
Cell constants	156.29Å 156.29Å 173.05Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.41 - 2.39	Depositor
rtesolution (A)	46.41 - 2.39	EDS
% Data completeness	99.9 (46.41-2.39)	Depositor
(in resolution range)	99.9 (46.41-2.39)	EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.18 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874	Depositor
D D.	0.162 , 0.197	Depositor
R, R_{free}	0.161 , 0.196	DCC
R_{free} test set	2484 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	39.5	Xtriage
Anisotropy	0.485	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 42.3	EDS
L-test for twinning ²	$ < L >=0.51, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6405	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.76% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, TRS, CL, NA, GOL, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
Mol	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.39	0/2788	0.52	1/3791 (0.0%)
1	В	0.37	0/2778	0.51	0/3781
2	Е	1.05	0/321	1.01	0/495
3	F	0.97	0/273	1.12	1/419 (0.2%)
All	All	0.48	0/6160	0.60	2/8486 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	F	3	DT	O4'-C1'-N1	5.72	112.01	108.00
1	A	242	LEU	CA-CB-CG	5.20	127.25	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2736	0	2762	12	0
1	В	2721	0	2738	15	0
2	Е	285	0	158	1	0
3	F	245	0	136	0	0
4	A	2	0	0	0	0

Continued on next page...



Continued from previous page.					
	C'	ontinued	from	previous	page

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	В	1	0	0	0	0
4	F	1	0	0	0	0
5	A	17	0	18	0	0
5	В	17	0	18	0	0
6	A	24	0	32	0	0
6	В	18	0	24	0	0
6	${ m E}$	6	0	8	0	0
7	A	1	0	0	0	0
7	F	1	0	0	0	0
8	В	1	0	0	0	0
9	В	10	0	14	0	0
10	A	135	0	0	0	0
10	В	119	0	0	0	0
10	Е	31	0	0	0	0
10	F	34	0	0	0	0
All	All	6405	0	5908	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 25 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} \text{Interatomic} \\ \text{distance (Å)} \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:242:LEU:HD21	1:A:272:ILE:HG13	1.71	0.70
1:B:7:HIS:H	2:E:1[A]:DG:H1	1.40	0.69
1:A:11:TYR:O	1:A:15:ILE:HG12	2.09	0.52
1:B:203:LEU:HD11	1:B:244:ILE:HD12	1.90	0.52
1:B:208:GLU:OE2	1:B:323:ARG:NH2	2.40	0.51

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	349/351 (99%)	344 (99%)	5 (1%)	0	100	100
1	В	346/351 (99%)	339 (98%)	7 (2%)	0	100	100
All	All	$695/702 \ (99\%)$	683 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	303/305 (99%)	296 (98%)	7 (2%)	50 70	
1	В	302/305 (99%)	293 (97%)	9 (3%)	41 61	
All	All	605/610 (99%)	589 (97%)	16 (3%)	47 66	

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	273	PHE
1	В	172[B]	ASN
1	В	8	GLU
1	В	172[A]	ASN
1	В	2	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 22 ligands modelled in this entry, 7 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

N / L 1	D	Cl:	D	T : 1-	В	ond leng	$_{ m gths}$	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	A	407	-	5,5,5	1.06	0	5,5,5	0.94	0
6	GOL	В	406	-	5,5,5	0.96	0	5,5,5	1.01	0
9	PGE	В	403	-	9,9,9	0.43	0	8,8,8	0.27	0
6	GOL	В	407	-	5,5,5	0.78	0	5,5,5	0.97	0
5	TRS	A	403[A]	-	7,7,7	0.25	0	9,9,9	0.78	0
5	TRS	В	405[A]	-	7,7,7	0.39	0	9,9,9	1.53	1 (11%)
5	TRS	В	404	-	7,7,7	0.32	0	9,9,9	0.47	0
6	GOL	Е	101	-	5,5,5	1.22	0	5,5,5	1.13	0
6	GOL	A	406	-	5,5,5	1.00	0	5,5,5	0.98	0
5	TRS	A	404	-	7,7,7	0.31	0	9,9,9	0.32	0
5	TRS	A	403[B]	-	7,7,7	0.25	0	9,9,9	0.84	0
5	TRS	В	405[B]	-	7,7,7	0.39	0	9,9,9	1.47	1 (11%)
6	GOL	A	405	-	5,5,5	1.02	0	5,5,5	0.87	0
6	GOL	В	408	-	5,5,5	1.47	1 (20%)	5,5,5	0.71	0
6	GOL	A	408	-	5,5,5	0.97	0	5,5,5	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	407	-	-	2/4/4/4	-
6	GOL	В	406	-	-	2/4/4/4	-
9	PGE	В	403	-	-	0/7/7/7	-

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	В	407	-	-	2/4/4/4	1
5	TRS	A	403[A]	-	-	3/9/9/9	-
5	TRS	В	405[A]	-	-	9/9/9/9	-
5	TRS	В	404	-	-	3/9/9/9	-
6	GOL	Е	101	-	-	1/4/4/4	-
6	GOL	A	406	-	-	4/4/4/4	-
5	TRS	A	404	-	-	3/9/9/9	1
5	TRS	A	403[B]	-	-	0/9/9/9	-
5	TRS	В	405[B]	-	-	9/9/9/9	-
6	GOL	A	405	-	-	2/4/4/4	-
6	GOL	В	408	-	-	4/4/4/4	-
6	GOL	A	408	-	-	0/4/4/4	-

All (1) bond length outliers are listed below:

	Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$Ideal(\AA)$
Ī	6	В	408	GOL	C3-C2	2.37	1.61	1.51

All (2) bond angle outliers are listed below:

	Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
	5	В	405[A]	TRS	O1-C1-C	3.06	120.71	111.00
ſ	5	В	405[B]	TRS	O1-C1-C	3.06	120.71	111.00

There are no chirality outliers.

5 of 44 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	403[A]	TRS	C2-C-C1-O1
5	A	403[A]	TRS	C3-C-C1-O1
5	A	403[A]	TRS	N-C-C1-O1
5	A	404	TRS	C1-C-C3-O3
5	A	404	TRS	C2-C-C3-O3


There are no ring outliers.

No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will



also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









Rings

Torsions























































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	350/351 (99%)	-0.26	9 (2%) 56 54	30, 41, 74, 128	0
1	В	347/351 (98%)	-0.16	17 (4%) 29 28	26, 45, 91, 140	0
2	E	13/13 (100%)	-0.22	0 100 100	30, 34, 47, 48	0
3	F	12/13 (92%)	-0.38	0 100 100	30, 33, 40, 42	0
All	All	722/728 (99%)	-0.21	26 (3%) 42 42	26, 42, 84, 140	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	174	ILE	5.7
1	В	222	ALA	5.2
1	В	283	ASP	5.0
1	A	172	ASN	4.6
1	В	167	ARG	4.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$ m B ext{-}factors(\AA^2)$	Q < 0.9
6	GOL	В	408	6/6	0.57	0.25	51,63,65,70	0
6	GOL	A	407	6/6	0.62	0.24	66,72,75,84	0
7	CL	A	409	1/1	0.66	0.14	74,74,74,74	0
6	GOL	Е	101	6/6	0.70	0.34	40,52,55,56	0
5	TRS	В	404	8/8	0.71	0.19	57,71,74,75	0
5	TRS	A	404	8/8	0.76	0.24	62,66,74,79	0
6	GOL	A	406	6/6	0.78	0.24	71,78,78,81	0
6	GOL	В	406	6/6	0.80	0.24	57,70,72,76	0
6	GOL	A	408	6/6	0.81	0.40	73,77,79,80	0
5	TRS	В	405[B]	8/8	0.82	0.17	36,44,48,49	1
5	TRS	В	405[A]	8/8	0.82	0.17	36,44,48,49	1
5	TRS	A	403[B]	8/8	0.86	0.25	42,53,55,57	1
5	TRS	A	403[A]	8/8	0.86	0.25	46,53,55,57	1
6	GOL	В	407	6/6	0.87	0.30	51,58,65,66	0
7	CL	F	102	1/1	0.87	0.18	62,62,62,62	0
6	GOL	A	405	6/6	0.88	0.30	59,68,73,77	0
9	PGE	В	403	10/10	0.94	0.12	41,47,55,57	0
4	CA	В	401	1/1	0.96	0.09	75,75,75,75	0
4	CA	F	101	1/1	0.98	0.07	38,38,38,38	0
8	NA	В	402	1/1	0.98	0.11	57,57,57,57	0
4	CA	A	401	1/1	0.98	0.11	31,31,31,31	0
4	CA	A	402	1/1	0.99	0.07	68,68,68,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.














Electron density around CL A 409:


 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)


























Electron density around GOL A 406:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around GOL B 406: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)


























































Electron density around NA B 402: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

