



wwPDB EM Validation Summary Report ⓘ

Dec 9, 2024 – 03:03 PM EST

PDB ID : 9EH0
EMDB ID : EMD-48042
Title : RNA polymerase II-DSIF-SPT6-PAF1c-TFIIS-IWS1-SETD2-nucleosome, 30 bp upstream
Authors : Markert, J.; Farnung, L.
Deposited on : 2024-11-21
Resolution : 3.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415
RNA backbone	6643	2191

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1984	
2	B	1251	
3	C	275	
4	D	142	
5	E	210	
6	F	127	
7	G	172	




Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
8	H	150	 75% 24%
9	I	125	 69% 24% 7%
10	J	67	 78% 21%
11	K	117	 79% 19%
12	L	58	 55% 24% 19%
13	M	1729	 57% 42%
14	N	198	 16% 72% 9%
15	O	821	 14% 84%
16	P	13	 8% 46% 31% 15%
17	Q	1179	 7% 58% 18% 25%
18	R	713	 29% 6% 66%
19	S	304	 52% 47%
20	T	198	 37% 58% 5% 2%
21	U	666	 16% 81%
22	V	531	 7% 40% 6% 54%
23	W	305	 72% 26%
24	X	531	 6% 92%
25	Y	121	 78% 18%
26	Z	1087	 40% 7% 53%
27	a	136	 73% 26%
27	e	136	 69% 31%
28	b	103	 75% 24%
28	f	103	 76% 24%
29	c	130	 80% 20%
29	g	130	 79% 21%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
30	d	123	 75% 25%
30	h	123	 72% 28%
31	l	1133	 36% 64%

2 Entry composition [i](#)

There are 33 unique types of molecules in this entry. The entry contains 73438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	N	O	P			S
1	A	1426	11255	7074	2014	2095	2	70	0	0

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	1122	8980	5684	1576	1656	64	0	0

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	258	2072	1300	356	410	6	0	0

- Molecule 4 is a protein called RNA polymerase Rpb4/RPC9 core domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	D	126	1004	630	170	200	4	0	0

- Molecule 5 is a protein called DNA-directed RNA polymerase II subunit E.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	E	209	1720	1089	300	323	8	0	0

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	F	78	626	401	106	114	5	0	0

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	G	171	1333	866	214	245	8	0	0

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	H	149	1197	759	195	238	5	0	0

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	I	116	942	582	168	181	11	0	0

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	J	66	524	339	88	91	6	0	0

- Molecule 11 is a protein called RNA polymerase II subunit J.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	K	115	920	593	152	173	2	0	0

- Molecule 12 is a protein called RNA polymerase II subunit K.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	L	47	397	246	77	68	6	0	0

- Molecule 13 is a protein called Transcription elongation factor SPT6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	M	1002	4883	2708	1072	1096	7	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	-2	SER	-	expression tag	UNP Q7KZ85
M	-1	ASN	-	expression tag	UNP Q7KZ85
M	0	ALA	-	expression tag	UNP Q7KZ85

- Molecule 14 is a DNA chain called Non-template.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	181	Total	C	N	O	P	0	0
			3695	1757	664	1093	181		

- Molecule 15 is a protein called Protein IWS1 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	132	Total	C	N	O	S	0	0
			1046	663	181	196	6		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
O	-1	SER	-	expression tag	UNP Q96ST2
O	0	ASN	-	expression tag	UNP Q96ST2
O	1	ALA	-	expression tag	UNP Q96ST2

- Molecule 16 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	P	11	Total	C	N	O	P	0	0
			233	105	42	75	11		

- Molecule 17 is a protein called RNA polymerase-associated protein CTR9 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	Q	890	Total	C	N	O	S	0	0
			7226	4579	1264	1352	31		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Q	1174	GLU	-	expression tag	UNP Q6PD62
Q	1175	ASN	-	expression tag	UNP Q6PD62
Q	1176	LEU	-	expression tag	UNP Q6PD62
Q	1177	TYR	-	expression tag	UNP Q6PD62

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Q	1178	PHE	-	expression tag	UNP Q6PD62
Q	1179	GLN	-	expression tag	UNP Q6PD62

- Molecule 18 is a protein called RNA polymerase-associated protein RTF1 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
18	R	244	1836	1152	340	337	7	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
R	-2	SER	-	expression tag	UNP Q92541
R	-1	ASN	-	expression tag	UNP Q92541
R	0	ALA	-	expression tag	UNP Q92541

- Molecule 19 is a protein called Transcription elongation factor A protein 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
19	S	161	657	334	161	162	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
S	-2	SER	-	expression tag	UNP P23193
S	-1	ASN	-	expression tag	UNP P23193
S	0	ALA	-	expression tag	UNP P23193

- Molecule 20 is a DNA chain called Template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
20	T	192	3955	1874	745	1144	192	0	0

- Molecule 21 is a protein called RNA polymerase-associated protein LEO1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
21	U	125	856	538	151	166	1	0	0

- Molecule 22 is a protein called RNA polymerase II-associated factor 1 homolog.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
22	V	244	1703	1061	305	333	4	0	0

- Molecule 23 is a protein called WDR61.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
23	W	300	2333	1483	392	454	4	0	0

- Molecule 24 is a protein called Parafibromin.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
24	X	43	353	220	69	64	0	0

- Molecule 25 is a protein called Transcription elongation factor SPT4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
25	Y	116	911	570	159	173	9	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
Y	-3	GLY	-	expression tag	UNP P63272
Y	-2	PRO	-	expression tag	UNP P63272
Y	-1	GLY	-	expression tag	UNP P63272
Y	0	SER	-	expression tag	UNP P63272

- Molecule 26 is a protein called Transcription elongation factor SPT5.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	N	O	P	S		
26	Z	510	4025	2552	709	745	1	18	0	0

- Molecule 27 is a protein called Histone H3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
27	a	101	823	520	157	142	4	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
27	e	94	776	491	149	133	3	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
a	36	MET	LYS	engineered mutation	UNP A0A310TTQ1
e	36	MET	LYS	engineered mutation	UNP A0A310TTQ1

- Molecule 28 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
28	b	78	622	393	120	108	1	0	0
28	f	78	622	393	120	108	1	0	0

- Molecule 29 is a protein called Histone H2A type 1.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
29	c	104	800	504	156	140	0	0
29	g	103	795	501	155	139	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
c	99	ARG	GLY	conflict	UNP P06897
c	123	SER	ALA	conflict	UNP P06897
g	99	ARG	GLY	conflict	UNP P06897
g	123	SER	ALA	conflict	UNP P06897

- Molecule 30 is a protein called Histone H2B 1.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
30	d	92	721	454	129	136	2	0	0
30	h	89	694	438	122	132	2	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
d	3	MET	-	initiating methionine	UNP P02281
d	32	THR	SER	engineered mutation	UNP P02281
h	3	MET	-	initiating methionine	UNP P02281
h	32	THR	SER	engineered mutation	UNP P02281

- Molecule 31 is a protein called Histone-lysine N-methyltransferase SETD2.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	1	412	Total	C	N	O	S	0	0
			2893	1772	545	555	21		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
1	1432	SER	-	expression tag	UNP Q9BYW2
1	1433	ASN	-	expression tag	UNP Q9BYW2
1	1434	ALA	-	expression tag	UNP Q9BYW2
1	1962	LEU	PRO	conflict	UNP Q9BYW2

- Molecule 32 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
32	A	2	Total	Zn	0
			2	2	
32	B	1	Total	Zn	0
			1	1	
32	C	1	Total	Zn	0
			1	1	
32	I	2	Total	Zn	0
			2	2	
32	J	1	Total	Zn	0
			1	1	
32	L	1	Total	Zn	0
			1	1	
32	Y	1	Total	Zn	0
			1	1	


- Molecule 33 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
33	A	1	Total	Mg	0
			1	1	

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase subunit





- Molecule 3: DNA-directed RNA polymerase II subunit RPB3

Chain C: 77% 16% 6%




- Molecule 4: RNA polymerase Rpb4/RPC9 core domain-containing protein

Chain D: 74% 15% 11%




- Molecule 5: DNA-directed RNA polymerase II subunit E

Chain E: 80% 20%



- Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2

Chain F: 49% 13% 39%




- Molecule 7: DNA-directed RNA polymerase II subunit RPB7


Chain G: 82% 17%





- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3




• Molecule 19: Transcription elongation factor A protein 1




• Molecule 20: Template DNA





• Molecule 21: RNA polymerase-associated protein LEO1







● Molecule 22: RNA polymerase II-associated factor 1 homolog




● Molecule 23: WDR61




● Molecule 28: Histone H4




● Molecule 29: Histone H2A type 1




● Molecule 29: Histone H2A type 1




● Molecule 30: Histone H2B 1.1



● Molecule 30: Histone H2B 1.1



● Molecule 31: Histone-lysine N-methyltransferase SETD2



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	301837	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	900	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.683	Depositor
Minimum map value	-0.193	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.010	Depositor
Recommended contour level	0.01	Depositor
Map size (Å)	550.0, 550.0, 550.0	wwPDB
Map dimensions	500, 500, 500	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.1, 1.1, 1.1	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SEP, TPO, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.28	0/11437	0.50	0/15433
2	B	0.30	0/9158	0.50	0/12360
3	C	0.31	0/2115	0.49	0/2873
4	D	0.26	0/1017	0.45	0/1368
5	E	0.28	0/1751	0.50	0/2366
6	F	0.29	0/636	0.50	0/859
7	G	0.29	0/1364	0.49	0/1853
8	H	0.33	0/1219	0.51	0/1644
9	I	0.29	0/964	0.48	0/1305
10	J	0.33	0/533	0.49	0/719
11	K	0.30	0/939	0.46	0/1271
12	L	0.31	0/403	0.58	0/536
13	M	0.24	0/4944	0.46	0/6387
14	N	0.58	0/4138	1.01	7/6379 (0.1%)
15	O	0.24	0/1062	0.42	0/1428
16	P	0.20	0/260	0.81	0/402
17	Q	0.27	0/7365	0.47	0/9927
18	R	0.25	0/1866	0.47	0/2519
19	S	0.24	0/659	0.44	0/827
20	T	0.61	1/4443 (0.0%)	0.97	7/6859 (0.1%)
21	U	0.27	0/870	0.47	0/1183
22	V	0.25	0/1728	0.49	0/2357
23	W	0.26	0/2392	0.47	0/3257
24	X	0.27	0/356	0.56	0/478
25	Y	0.25	0/927	0.49	0/1250
26	Z	0.25	0/4084	0.49	0/5498
27	a	0.33	0/835	0.58	0/1120
27	e	0.31	0/786	0.59	0/1053
28	b	0.31	0/629	0.59	0/843
28	f	0.31	0/629	0.59	0/843
29	c	0.29	0/810	0.57	0/1095
29	g	0.29	0/805	0.57	0/1088

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
30	d	0.30	0/732	0.50	0/986
30	h	0.30	0/705	0.49	0/951
31	l	0.29	0/2932	0.54	0/3963
All	All	0.33	1/75493 (0.0%)	0.58	14/103280 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
31	l	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
20	T	11	DG	P-OP2	-5.17	1.40	1.49

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	N	45	DC	OP1-P-O3'	7.13	120.90	105.20
20	T	-46	DG	OP1-P-OP2	-6.90	109.25	119.60
20	T	-49	DC	OP1-P-OP2	-6.76	109.45	119.60
14	N	46	DC	OP1-P-OP2	-6.75	109.48	119.60
20	T	-48	DC	OP1-P-OP2	-6.56	109.76	119.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
31	l	2018	ASP	Peptide
31	l	2021	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11255	0	11374	184	0
2	B	8980	0	9019	153	0
3	C	2072	0	2020	39	0
4	D	1004	0	980	16	0
5	E	1720	0	1737	32	0
6	F	626	0	657	10	0
7	G	1333	0	1321	23	0
8	H	1197	0	1156	27	0
9	I	942	0	872	25	0
10	J	524	0	541	12	0
11	K	920	0	942	16	0
12	L	397	0	405	15	0
13	M	4883	0	2550	10	0
14	N	3695	0	2038	226	0
15	O	1046	0	1096	10	0
16	P	233	0	118	15	0
17	Q	7226	0	7169	155	0
18	R	1836	0	1699	26	0
19	S	657	0	199	2	0
20	T	3955	0	2154	142	0
21	U	856	0	680	14	0
22	V	1703	0	1426	28	0
23	W	2333	0	2246	48	0
24	X	353	0	371	9	0
25	Y	911	0	907	15	0
26	Z	4025	0	4041	56	0
27	a	823	0	864	0	0
27	e	776	0	815	0	0
28	b	622	0	660	0	0
28	f	622	0	660	0	0
29	c	800	0	851	0	0
29	g	795	0	846	0	0
30	d	721	0	742	0	0
30	h	694	0	709	0	0
31	l	2893	0	2422	0	0
32	A	2	0	0	0	0
32	B	1	0	0	0	0
32	C	1	0	0	0	0
32	I	2	0	0	0	0
32	J	1	0	0	0	0
32	L	1	0	0	0	0
32	Y	1	0	0	0	0
33	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	73438	0	66287	1232	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 1232 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
17:Q:276:ALA:HB1	17:Q:288:VAL:HG23	1.31	1.12
1:A:1172:ASN:N	1:A:1215:GLU:OE2	1.99	0.95
20:T:9:DG:H2''	20:T:10:DC:H5'	1.50	0.94
2:B:790:GLN:O	2:B:968:ASN:ND2	2.01	0.93
20:T:7:DG:H2'	20:T:8:DT:C6	2.03	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1408/1984 (71%)	1319 (94%)	88 (6%)	1 (0%)	48	79
2	B	1112/1251 (89%)	1024 (92%)	88 (8%)	0	100	100
3	C	254/275 (92%)	229 (90%)	25 (10%)	0	100	100
4	D	124/142 (87%)	120 (97%)	4 (3%)	0	100	100
5	E	207/210 (99%)	198 (96%)	9 (4%)	0	100	100
6	F	76/127 (60%)	74 (97%)	2 (3%)	0	100	100
7	G	169/172 (98%)	161 (95%)	8 (5%)	0	100	100
8	H	147/150 (98%)	134 (91%)	13 (9%)	0	100	100
9	I	114/125 (91%)	103 (90%)	11 (10%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
10	J	64/67 (96%)	57 (89%)	7 (11%)	0	100	100
11	K	113/117 (97%)	110 (97%)	3 (3%)	0	100	100
12	L	45/58 (78%)	41 (91%)	3 (7%)	1 (2%)	5	32
13	M	976/1729 (56%)	920 (94%)	55 (6%)	1 (0%)	48	79
15	O	130/821 (16%)	127 (98%)	3 (2%)	0	100	100
17	Q	888/1179 (75%)	847 (95%)	41 (5%)	0	100	100
18	R	240/713 (34%)	231 (96%)	9 (4%)	0	100	100
19	S	157/304 (52%)	154 (98%)	3 (2%)	0	100	100
21	U	117/666 (18%)	106 (91%)	10 (8%)	1 (1%)	14	48
22	V	234/531 (44%)	216 (92%)	18 (8%)	0	100	100
23	W	298/305 (98%)	288 (97%)	10 (3%)	0	100	100
24	X	41/531 (8%)	41 (100%)	0	0	100	100
25	Y	114/121 (94%)	108 (95%)	6 (5%)	0	100	100
26	Z	497/1087 (46%)	479 (96%)	18 (4%)	0	100	100
27	a	99/136 (73%)	96 (97%)	3 (3%)	0	100	100
27	e	90/136 (66%)	89 (99%)	1 (1%)	0	100	100
28	b	76/103 (74%)	76 (100%)	0	0	100	100
28	f	76/103 (74%)	75 (99%)	1 (1%)	0	100	100
29	c	102/130 (78%)	101 (99%)	1 (1%)	0	100	100
29	g	101/130 (78%)	100 (99%)	1 (1%)	0	100	100
30	d	90/123 (73%)	89 (99%)	1 (1%)	0	100	100
30	h	87/123 (71%)	83 (95%)	4 (5%)	0	100	100
31	l	406/1133 (36%)	374 (92%)	30 (7%)	2 (0%)	25	59
All	All	8652/14782 (58%)	8170 (94%)	476 (6%)	6 (0%)	50	79

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
12	L	39	CYS
13	M	700	HIS
31	l	2018	ASP
31	l	2022	VAL
1	A	1343	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1245/1761 (71%)	1241 (100%)	4 (0%)	91	96
2	B	986/1084 (91%)	982 (100%)	4 (0%)	89	95
3	C	235/252 (93%)	234 (100%)	1 (0%)	89	95
4	D	109/126 (86%)	108 (99%)	1 (1%)	75	87
5	E	191/192 (100%)	189 (99%)	2 (1%)	73	85
6	F	68/111 (61%)	67 (98%)	1 (2%)	60	78
7	G	146/153 (95%)	146 (100%)	0	100	100
8	H	130/131 (99%)	130 (100%)	0	100	100
9	I	104/112 (93%)	103 (99%)	1 (1%)	73	85
10	J	55/56 (98%)	55 (100%)	0	100	100
11	K	104/106 (98%)	104 (100%)	0	100	100
12	L	44/55 (80%)	44 (100%)	0	100	100
13	M	196/1524 (13%)	196 (100%)	0	100	100
15	O	118/737 (16%)	116 (98%)	2 (2%)	56	75
17	Q	761/1011 (75%)	752 (99%)	9 (1%)	67	82
18	R	170/625 (27%)	168 (99%)	2 (1%)	67	82
19	S	4/268 (2%)	4 (100%)	0	100	100
21	U	65/590 (11%)	65 (100%)	0	100	100
22	V	144/462 (31%)	142 (99%)	2 (1%)	62	79
23	W	255/260 (98%)	251 (98%)	4 (2%)	58	76
24	X	40/467 (9%)	40 (100%)	0	100	100
25	Y	102/105 (97%)	102 (100%)	0	100	100
26	Z	435/939 (46%)	433 (100%)	2 (0%)	86	93
27	a	87/111 (78%)	85 (98%)	2 (2%)	45	68
27	e	82/111 (74%)	82 (100%)	0	100	100
28	b	64/79 (81%)	63 (98%)	1 (2%)	58	76

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
28	f	64/79 (81%)	64 (100%)	0	100	100
29	c	82/102 (80%)	82 (100%)	0	100	100
29	g	82/102 (80%)	82 (100%)	0	100	100
30	d	79/103 (77%)	79 (100%)	0	100	100
30	h	76/103 (74%)	75 (99%)	1 (1%)	65	81
31	l	237/1017 (23%)	231 (98%)	6 (2%)	42	66
All	All	6560/12934 (51%)	6515 (99%)	45 (1%)	80	90

5 of 45 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
22	V	132	ARG
27	a	36	MET
23	W	43	LEU
23	W	272	ASP
28	b	31	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 44 such sidechains are listed below:

Mol	Chain	Res	Type
17	Q	466	ASN
22	V	72	HIS
17	Q	527	HIS
17	Q	775	ASN
23	W	273	HIS

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
16	P	11/13 (84%)	2 (18%)	2 (18%)

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
16	P	17	A
16	P	19	A

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
16	P	16	A
16	P	18	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SEP	A	1547	1	8,9,10	1.61	1 (12%)	7,12,14	1.28	1 (14%)
1	TPO	A	1525	1	8,10,11	1.12	0	10,14,16	2.13	1 (10%)
26	TPO	Z	775	26	8,10,11	1.13	0	10,14,16	1.90	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	1547	1	-	0/6/8/10	-
1	TPO	A	1525	1	-	0/9/11/13	-
26	TPO	Z	775	26	-	2/9/11/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1547	SEP	P-O1P	3.51	1.61	1.50

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1525	TPO	P-OG1-CB	-6.08	106.81	123.33
26	Z	775	TPO	P-OG1-CB	-5.19	109.22	123.33
1	A	1547	SEP	OG-CB-CA	2.76	110.83	108.14

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
26	Z	775	TPO	C-CA-CB-CG2
26	Z	775	TPO	CB-OG1-P-O2P

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1525	TPO	1	0
26	Z	775	TPO	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
21	U	1

Continued on next page...

Continued from previous page...

Mol	Chain	Number of breaks
22	V	1
13	M	1
31	l	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	U	497:ASP	C	505:SER	N	27.02
1	V	299:GLU	C	310:ASN	N	12.71
1	M	1334:ASN	C	1338:ILE	N	5.60
1	l	1709:SER	C	1710:ARG	N	4.41


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-48042. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 221



Y Index: 253



Z Index: 275


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 5148 nm³; this corresponds to an approximate mass of 4651 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.278 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-48042 and PDB model 9EH0. Per-residue inclusion information can be found in section 3 on page 12.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).









































































9.4 Atom inclusion [i](#)



At the recommended contour level, 98% of all backbone atoms, 98% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9800	 0.2400
A	 0.9930	 0.3520
B	 0.9910	 0.3820
C	 0.9970	 0.4190
D	 0.9880	 0.1880
E	 0.9940	 0.3350
F	 0.9870	 0.3930
G	 0.9890	 0.2060
H	 0.9840	 0.3950
I	 0.9890	 0.3460
J	 0.9900	 0.3930
K	 0.9970	 0.4010
L	 1.0000	 0.3580
M	 0.9950	 0.1470
N	 1.0000	 0.2300
O	 1.0000	 0.0150
P	 1.0000	 0.2090
Q	 0.8850	 0.0790
R	 0.9670	 0.0920
S	 1.0000	 0.2400
T	 1.0000	 0.2210
U	 0.9930	 0.0160
V	 0.8710	 0.0570
W	 1.0000	 0.1550
X	 0.9940	 0.1760
Y	 0.9970	 -0.0020
Z	 0.9990	 0.0700
a	 0.9870	 0.3390
b	 0.9980	 0.3680
c	 0.9870	 0.3220
d	 0.9890	 0.3130
e	 0.9810	 0.3320
f	 0.9800	 0.3550
g	 0.9920	 0.3260
h	 0.9880	 0.3260
l	 0.9970	 0.1220

