



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 11:26 am GMT

PDB ID : 1EJ7
Title : CRYSTAL STRUCTURE OF UNACTIVATED TOBACCO RUBISCO WITH
BOUND PHOSPHATE IONS
Authors : Duff, A.P.; Andrews, T.J.; Curmi, P.M.G.
Deposited on : 2000-03-01
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1259 (2.48-2.44)
Clashscore	122126	1323 (2.48-2.44)
Ramachandran outliers	120053	1314 (2.48-2.44)
Sidechain outliers	120020	1314 (2.48-2.44)
RSRZ outliers	108989	1238 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	L	475	
2	S	123	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PO4	L	641	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4790 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RUBISCO (LARGE SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	457	3579	2274	631	658	16	32	0	0


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
L	19	ASP	GLU	CONFLICT	UNP P00876
L	377	VAL	GLU	CONFLICT	UNP P00876

- Molecule 2 is a protein called RUBISCO (SMALL SUBUNIT).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	S	123	1029	672	163	188	6	0	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	L	1	Total O P 5 4 1	0	0
3	L	1	Total O P 5 4 1	0	0
3	L	1	Total O P 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	139	Total O 139 139	0	0
4	S	28	Total O 28 28	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RUBISCO (LARGE SUBUNIT)



- Molecule 2: RUBISCO (SMALL SUBUNIT)



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	149.18Å 149.18Å 138.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.45 19.61 – 2.45	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.45) 93.2 (19.61-2.45)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.46Å)	Xtrriage
Refinement program	X-PLOR 3.843	Depositor
R, R_{free}	0.149 , 0.197 0.157 , 0.202	Depositor DCC
R_{free} test set	1407 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	37.0	Xtrriage
Anisotropy	0.028	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 56.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4790	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	L	0.60	5/3666 (0.1%)	0.85	16/4973 (0.3%)
2	S	0.58	3/1062 (0.3%)	0.96	7/1442 (0.5%)
All	All	0.60	8/4728 (0.2%)	0.88	23/6415 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	L	0	3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	471	ALA	N-CA	-17.63	1.11	1.46
1	L	470	ALA	N-CA	-13.71	1.19	1.46
1	L	470	ALA	CA-C	-8.71	1.30	1.52
2	S	606	ASN	N-CA	7.44	1.61	1.46
2	S	605	ASP	CA-C	7.04	1.71	1.52
1	L	470	ALA	C-N	-6.22	1.19	1.34
1	L	472	VAL	C-N	-5.80	1.20	1.34
2	S	610	VAL	C-N	5.30	1.46	1.34

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	S	610	VAL	O-C-N	16.92	149.77	122.70
1	L	470	ALA	O-C-N	16.36	148.87	122.70
1	L	469	PHE	CB-CA-C	-15.56	79.28	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	S	610	VAL	CA-C-N	-13.63	87.20	117.20
1	L	470	ALA	CA-C-N	-13.53	87.44	117.20
1	L	469	PHE	CB-CG-CD2	-12.85	111.81	120.80
1	L	336	GLU	O-C-N	-10.47	105.40	123.20
2	S	610	VAL	C-N-CA	-9.35	98.34	121.70
1	L	336	GLU	N-CA-C	-8.25	88.73	111.00
1	L	469	PHE	CA-C-N	-7.27	101.20	117.20
1	L	336	GLU	CA-C-N	7.25	130.70	116.20
2	S	608	ARG	NE-CZ-NH2	6.74	123.67	120.30
1	L	469	PHE	O-C-N	6.54	133.16	122.70
1	L	469	PHE	N-CA-CB	6.39	122.09	110.60
1	L	470	ALA	CB-CA-C	-6.02	101.07	110.10
1	L	469	PHE	CD1-CG-CD2	6.02	126.12	118.30
2	S	605	ASP	O-C-N	-6.00	113.09	122.70
1	L	469	PHE	N-CA-C	5.75	126.53	111.00
1	L	336	GLU	C-N-CA	-5.61	110.53	122.30
1	L	471	ALA	CB-CA-C	-5.37	102.04	110.10
1	L	472	VAL	C-N-CA	5.24	134.79	121.70
2	S	605	ASP	CB-CA-C	-5.20	100.00	110.40
2	S	609	GLN	CB-CA-C	-5.03	100.35	110.40

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	L	471	ALA	Peptide
1	L	472	VAL	Peptide
1	L	473	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	3579	0	3500	48	0
2	S	1029	0	991	19	0
3	L	15	0	0	5	0
4	L	139	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	S	28	0	0	0	0
All	All	4790	0	4491	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:S:609:GLN:HG3	2:S:609:GLN:O	1.33	1.12
1:L:339:ARG:NH2	1:L:393:ILE:HG12	1.70	1.07
1:L:339:ARG:NH2	1:L:393:ILE:CG1	2.41	0.83
1:L:383:HIS:H	1:L:386:HIS:HD2	1.28	0.81
2:S:609:GLN:CG	2:S:609:GLN:O	2.17	0.79
1:L:335:LEU:C	1:L:336:GLU:O	2.17	0.74
1:L:75:THR:HG22	1:L:76:SER:H	1.54	0.71
1:L:339:ARG:NH1	1:L:393:ILE:HD11	2.07	0.70
2:S:606:ASN:OD1	2:S:606:ASN:N	2.25	0.69
2:S:608:ARG:HB3	2:S:610:VAL:HG22	1.77	0.66
2:S:556:ASN:ND2	2:S:558:SER:H	1.95	0.63
1:L:60:GLU:HB2	1:L:124:VAL:HG12	1.81	0.63
2:S:608:ARG:O	2:S:610:VAL:HG13	1.98	0.62
2:S:510:LYS:HB3	2:S:550:PHE:CE1	2.37	0.59
1:L:335:LEU:O	1:L:336:GLU:O	2.21	0.58
1:L:383:HIS:H	1:L:386:HIS:CD2	2.17	0.56
1:L:170:LEU:HD22	1:L:402:PHE:HE2	1.72	0.55
2:S:511:LYS:HG3	2:S:517:TYR:CE1	2.41	0.54
1:L:414:ALA:HB3	1:L:415:PRO:HD3	1.89	0.54
1:L:403:GLY:HA3	3:L:641:PO4:P	2.48	0.54
1:L:328:SER:HB2	1:L:345:PHE:HE1	1.73	0.53
1:L:339:ARG:CZ	1:L:393:ILE:HD11	2.40	0.52
1:L:172:CYS:HB3	1:L:197:LEU:CD1	2.40	0.51
1:L:383:HIS:N	1:L:386:HIS:HD2	2.05	0.51
1:L:403:GLY:HA3	3:L:641:PO4:O1	2.11	0.51
1:L:339:ARG:HH21	1:L:393:ILE:HG12	1.71	0.51
1:L:201:LYS:HG2	1:L:202:ASP:O	2.12	0.50
1:L:305:LYS:HE2	3:L:643:PO4:O4	2.13	0.49
1:L:94:LYS:O	1:L:95:ASP:HB2	2.13	0.49
1:L:155:ILE:HG12	1:L:375:LEU:HD13	1.95	0.48
1:L:339:ARG:CZ	1:L:393:ILE:CD1	2.92	0.48
1:L:172:CYS:SG	1:L:192:CYS:SG	3.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:75:THR:HG22	1:L:76:SER:N	2.28	0.48
2:S:608:ARG:O	2:S:609:GLN:HB3	2.12	0.48
1:L:339:ARG:HH12	1:L:393:ILE:HD11	1.76	0.47
1:L:407:LEU:HD13	1:L:413:ASN:OD1	2.14	0.47
1:L:172:CYS:HB3	1:L:197:LEU:HD13	1.97	0.47
1:L:79:ARG:HH11	1:L:79:ARG:HG2	1.80	0.47
1:L:463:LYS:HB3	1:L:469:PHE:O	2.15	0.47
1:L:298:HIS:NE2	3:L:642:PO4:O1	2.35	0.47
1:L:197:LEU:HG	1:L:417:ALA:HB1	1.97	0.47
1:L:70:TRP:HZ3	1:L:74:LEU:HD11	1.80	0.47
2:S:608:ARG:O	2:S:609:GLN:C	2.53	0.46
1:L:431:ARG:HH21	1:L:432:ASN:HD21	1.64	0.46
1:L:403:GLY:HA3	3:L:641:PO4:O4	2.16	0.45
2:S:540:PRO:HG2	2:S:574:MET:HB2	1.97	0.45
2:S:531:GLU:O	2:S:535:LYS:HD3	2.17	0.45
1:L:293:ILE:HG13	1:L:318:LEU:HD21	1.99	0.45
2:S:606:ASN:C	2:S:608:ARG:N	2.68	0.45
2:S:608:ARG:O	2:S:610:VAL:N	2.49	0.45
1:L:339:ARG:HH22	1:L:393:ILE:CG1	2.25	0.45
1:L:89:ARG:HH22	1:L:94:LYS:CB	2.30	0.45
1:L:371:LEU:HD22	1:L:372:PRO:HD2	1.98	0.44
2:S:514:THR:HG22	2:S:515:LEU:HG	1.98	0.43
1:L:336:GLU:HA	1:L:339:ARG:HB3	2.00	0.43
1:L:128:LYS:HD3	1:L:128:LYS:HA	1.87	0.43
1:L:421:ARG:O	1:L:425:GLU:HG3	2.19	0.43
1:L:117:PHE:O	1:L:121:VAL:HG22	2.19	0.43
1:L:335:LEU:O	1:L:336:GLU:C	2.56	0.42
1:L:239:TYR:HB3	1:L:266:MET:HB2	1.99	0.42
1:L:178:LEU:HD21	1:L:205:ASN:HB3	2.01	0.42
1:L:431:ARG:HE	1:L:432:ASN:ND2	2.17	0.42
2:S:607:VAL:HG23	2:S:608:ARG:HB2	2.01	0.42
2:S:556:ASN:HD21	2:S:558:SER:CB	2.33	0.41
2:S:533:LEU:HD11	2:S:613:ILE:HG21	2.02	0.41
1:L:339:ARG:NH2	1:L:393:ILE:CD1	2.83	0.41
2:S:606:ASN:C	2:S:608:ARG:H	2.19	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	L	455/475 (96%)	428 (94%)	23 (5%)	4 (1%)	19	22
2	S	121/123 (98%)	113 (93%)	8 (7%)	0	100	100
All	All	576/598 (96%)	541 (94%)	31 (5%)	4 (1%)	24	28

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	471	ALA
1	L	469	PHE
1	L	336	GLU
1	L	92	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	369/384 (96%)	360 (98%)	9 (2%)	52	65
2	S	110/110 (100%)	103 (94%)	7 (6%)	19	24
All	All	479/494 (97%)	463 (97%)	16 (3%)	41	53

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	28	GLU
1	L	66	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	L	172	CYS
1	L	241	ASN
1	L	258	ARG
1	L	300	VAL
1	L	371	LEU
1	L	446	ARG
1	L	469	PHE
2	S	568	THR
2	S	584	LEU
2	S	596	GLN
2	S	606	ASN
2	S	608	ARG
2	S	609	GLN
2	S	613	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	30	GLN
1	L	153	HIS
1	L	207	ASN
1	L	229	GLN
1	L	241	ASN
1	L	356	GLN
1	L	386	HIS
1	L	420	ASN
1	L	432	ASN
2	S	556	ASN
2	S	596	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	L	641	-	4,4,4	0.76	0	6,6,6	0.72	0
3	PO4	L	642	-	4,4,4	0.75	0	6,6,6	0.73	0
3	PO4	L	643	-	4,4,4	0.87	0	6,6,6	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PO4	L	641	-	-	0/0/0/0	0/0/0/0
3	PO4	L	642	-	-	0/0/0/0	0/0/0/0
3	PO4	L	643	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	L	641	PO4	3	0
3	L	642	PO4	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	L	643	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	L	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	L	470:ALA	C	471:ALA	N	1.19

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	L	452/475 (95%)	-0.38	24 (5%) 26 24	19, 35, 86, 100	0
2	S	123/123 (100%)	-0.28	6 (4%) 29 27	23, 50, 75, 92	0
All	All	575/598 (96%)	-0.36	30 (5%) 27 24	19, 39, 86, 100	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	66	TRP	11.3
1	L	469	PHE	10.3
1	L	67	THR	9.5
1	L	22	LEU	6.7
1	L	68	THR	6.1
2	S	607	VAL	5.6
1	L	91	VAL	4.5
1	L	65	THR	3.5
1	L	468	ASN	3.5
1	L	335	LEU	3.4
1	L	75	THR	3.4
1	L	405	GLY	3.4
1	L	407	LEU	3.1
2	S	606	ASN	3.1
2	S	508	ASN	3.0
1	L	78	ASP	3.0
1	L	19	ASP	3.0
1	L	94	LYS	2.9
1	L	18	LYS	2.8
1	L	70	TRP	2.8
2	S	576	GLY	2.8
2	S	621	GLU	2.8
1	L	46	PRO	2.8
1	L	69	VAL	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	L	47	GLY	2.7
1	L	92	GLY	2.5
1	L	380	GLY	2.3
1	L	336	GLU	2.3
2	S	578	THR	2.2
1	L	74	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	L	641	5/5	0.92	0.21	62,63,66,66	5
3	PO4	L	642	5/5	0.94	0.13	52,54,56,58	5
3	PO4	L	643	5/5	0.94	0.17	37,39,44,49	5

6.5 Other polymers [i](#)

There are no such residues in this entry.