



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 20, 2018 – 01:55 pm GMT

PDB ID : 1ENY
Title : CRYSTAL STRUCTURE AND FUNCTION OF THE ISONIAZID TARGET OF MYCOBACTERIUM TUBERCULOSIS
Authors : Dessen, A.; Quemard, A.; Blanchard, J.S.; Jacobs Jr., W.R.; Sacchettini, J.C.; TB Structural Genomics Consortium (TBSGC)
Deposited on : 1995-01-27
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30686


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	122078	5026 (2.20-2.20)
Ramachandran outliers	120005	4951 (2.20-2.20)
Sidechain outliers	119972	4952 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	268	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2079 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENOYL-ACYL CARRIER PROTEIN (ACP) REDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	268	1994	1263	348	373	10	0	0	0

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: $C_{21}H_{27}N_7O_{14}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	44	21	7	14	2	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	41	Total	O	0	0
			41	41		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: ENOYL-ACYL CARRIER PROTEIN (ACP) REDUCTASE



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	100.14Å 100.14Å 140.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	(Not available) – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.20)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	TNT, X-PLOR	Depositor
R, R_{free}	0.196 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2079	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.22	6/2032 (0.3%)	1.32	19/2758 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	1	0

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	171	VAL	CB-CG1	-6.20	1.39	1.52
1	A	158	TYR	CB-CG	-5.72	1.43	1.51
1	A	158	TYR	CD2-CE2	-5.53	1.31	1.39
1	A	220	GLU	CG-CD	5.50	1.60	1.51
1	A	68	GLU	CG-CD	5.36	1.59	1.51
1	A	153	ARG	CZ-NH1	5.05	1.39	1.33

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	173	ARG	NE-CZ-NH2	-9.89	115.35	120.30
1	A	16	ILE	CB-CA-C	8.11	127.82	111.60
1	A	225	ARG	NE-CZ-NH1	-8.05	116.27	120.30
1	A	185	ARG	NE-CZ-NH1	7.44	124.02	120.30
1	A	225	ARG	NE-CZ-NH2	7.24	123.92	120.30
1	A	145	VAL	CB-CA-C	-7.24	97.64	111.40
1	A	173	ARG	NE-CZ-NH1	7.24	123.92	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	11	LEU	CB-CG-CD1	6.06	121.30	111.00
1	A	11	LEU	CA-CB-CG	5.93	128.95	115.30
1	A	185	ARG	NE-CZ-NH2	-5.87	117.36	120.30
1	A	3	GLY	N-CA-C	-5.69	98.88	113.10
1	A	88	LEU	CA-CB-CG	-5.68	102.24	115.30
1	A	43	ARG	NE-CZ-NH2	-5.50	117.55	120.30
1	A	254	THR	CB-CA-C	-5.37	97.11	111.60
1	A	153	ARG	NE-CZ-NH1	5.27	122.94	120.30
1	A	37	VAL	N-CA-C	-5.23	96.87	111.00
1	A	50	ILE	CB-CA-C	-5.15	101.30	111.60
1	A	268	LEU	CB-CG-CD2	5.14	119.73	111.00
1	A	195	ARG	NE-CZ-NH1	5.12	122.86	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	236	THR	CA

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1994	0	2011	81	0
2	A	44	0	26	3	0
3	A	41	0	0	2	0
All	All	2079	0	2037	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186:SER:H	1:A:254:THR:HG22	1.27	0.99

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:LEU:HD13	1:A:134:LEU:HD21	1.45	0.98
1:A:234:ASP:OD1	1:A:236:THR:HG23	1.78	0.83
1:A:98:MET:HE2	1:A:160:TRP:HB2	1.61	0.83
1:A:43:ARG:O	1:A:47:ILE:HG13	1.83	0.79
1:A:104:GLY:CA	1:A:207:LEU:HD23	2.14	0.78
1:A:101:THR:HG21	1:A:115:ASP:OD2	1.85	0.77
1:A:193:PRO:HD2	1:A:230:TRP:NE1	2.05	0.72
1:A:185:ARG:HA	1:A:254:THR:CG2	2.19	0.71
1:A:61:LEU:HD23	1:A:61:LEU:N	2.05	0.71
1:A:16:ILE:HG22	1:A:20:SER:OG	1.90	0.71
1:A:104:GLY:HA2	1:A:207:LEU:HD23	1.72	0.70
1:A:87:LYS:HB3	1:A:137:ILE:O	1.91	0.70
1:A:4:LEU:H	1:A:32:GLN:HG3	1.56	0.69
1:A:78:VAL:O	1:A:82:ILE:HG12	1.91	0.69
1:A:202:ILE:HD12	1:A:202:ILE:O	1.97	0.65
1:A:51:THR:O	1:A:54:LEU:HB2	1.97	0.64
1:A:105:ILE:HG23	1:A:106:ASN:N	2.13	0.64
1:A:47:ILE:O	1:A:51:THR:HG23	1.96	0.64
1:A:186:SER:N	1:A:254:THR:HG22	2.08	0.63
1:A:46:LEU:HD12	1:A:46:LEU:O	1.99	0.62
1:A:44:LEU:HD21	1:A:62:GLU:HB2	1.83	0.61
1:A:145:VAL:HG12	1:A:146:GLY:N	2.15	0.61
1:A:54:LEU:HD12	1:A:55:PRO:HD3	1.82	0.60
1:A:4:LEU:HB3	1:A:32:GLN:HG2	1.82	0.60
1:A:185:ARG:HA	1:A:254:THR:HG23	1.85	0.59
1:A:4:LEU:N	1:A:32:GLN:HG3	2.18	0.59
1:A:191:ALA:HB3	2:A:500:NAD:C5N	2.34	0.57
1:A:199:MET:O	1:A:203:VAL:HG12	2.04	0.56
1:A:60:LEU:C	1:A:61:LEU:HD23	2.25	0.56
1:A:105:ILE:HG23	1:A:106:ASN:H	1.69	0.56
1:A:103:MET:O	1:A:157:ALA:O	2.24	0.56
1:A:43:ARG:HB3	1:A:46:LEU:HB3	1.88	0.56
1:A:28:VAL:HA	1:A:31:GLU:HG3	1.87	0.55
1:A:49:ARG:HG2	1:A:53:ARG:HH21	1.73	0.54
1:A:196:THR:HG22	1:A:199:MET:H	1.74	0.53
1:A:48:GLN:O	1:A:52:ASP:OD1	2.28	0.52
1:A:202:ILE:HD11	1:A:215:ILE:HG13	1.91	0.52
1:A:67:ASN:HD22	1:A:70:HIS:CD2	2.28	0.51
1:A:15:ILE:HG13	3:A:538:HOH:O	2.10	0.51
1:A:46:LEU:HD12	1:A:46:LEU:C	2.32	0.50
1:A:191:ALA:HB3	2:A:500:NAD:H5N	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:245:LEU:HD21	1:A:258:ILE:HD13	1.92	0.50
1:A:187:ASN:ND2	1:A:256:ASP:H	2.09	0.49
1:A:177:ARG:CG	1:A:177:ARG:HH11	2.24	0.49
1:A:98:MET:HE2	1:A:160:TRP:CB	2.39	0.49
1:A:135:LEU:N	1:A:136:PRO:CD	2.75	0.49
1:A:39:THR:HA	1:A:61:LEU:O	2.13	0.49
1:A:74:LEU:CD1	1:A:134:LEU:HD21	2.32	0.48
1:A:105:ILE:CG2	1:A:106:ASN:H	2.28	0.47
1:A:61:LEU:N	1:A:61:LEU:CD2	2.78	0.46
1:A:105:ILE:CG2	1:A:106:ASN:N	2.77	0.45
1:A:74:LEU:HD22	1:A:74:LEU:O	2.16	0.45
1:A:157:ALA:C	1:A:159:ASN:H	2.20	0.45
1:A:49:ARG:HG2	1:A:53:ARG:NH2	2.31	0.45
1:A:145:VAL:CG1	1:A:146:GLY:N	2.78	0.44
1:A:236:THR:N	1:A:237:PRO:HD2	2.32	0.44
1:A:67:ASN:ND2	1:A:70:HIS:CD2	2.86	0.44
1:A:44:LEU:CD2	1:A:62:GLU:HB2	2.47	0.44
1:A:130:MET:HE3	1:A:130:MET:HB3	1.79	0.44
1:A:187:ASN:ND2	1:A:256:ASP:CB	2.81	0.43
1:A:197:LEU:HD22	1:A:197:LEU:HA	1.53	0.43
1:A:259:TYR:CG	1:A:265:HIS:CE1	3.06	0.43
1:A:218:LEU:HD23	3:A:510:HOH:O	2.18	0.43
1:A:165:LYS:O	1:A:169:GLU:HG3	2.19	0.43
1:A:135:LEU:CB	1:A:136:PRO:HD3	2.49	0.42
1:A:268:LEU:HD12	1:A:268:LEU:HA	1.59	0.42
1:A:177:ARG:CG	1:A:177:ARG:NH1	2.82	0.42
1:A:111:ALA:HA	1:A:112:PRO:HD3	1.76	0.42
1:A:225:ARG:HD3	1:A:225:ARG:HH11	1.60	0.42
1:A:98:MET:CE	1:A:160:TRP:HB2	2.40	0.41
1:A:202:ILE:HD12	1:A:202:ILE:C	2.40	0.41
1:A:48:GLN:C	1:A:48:GLN:NE2	2.74	0.41
1:A:98:MET:CE	1:A:160:TRP:CB	2.99	0.41
1:A:30:GLN:OE1	1:A:55:PRO:HD2	2.21	0.41
1:A:57:LYS:HB2	1:A:57:LYS:HE2	1.61	0.41
1:A:185:ARG:CA	1:A:254:THR:HG23	2.50	0.41
1:A:63:LEU:C	1:A:63:LEU:HD12	2.41	0.41
1:A:100:GLN:HB3	1:A:100:GLN:HE21	1.73	0.41
1:A:101:THR:HG21	1:A:115:ASP:CG	2.42	0.41
1:A:148:ASP:O	2:A:500:NAD:H6N	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	266/268 (99%)	243 (91%)	21 (8%)	2 (1%)	21 20

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	159	ASN
1	A	104	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	203/203 (100%)	157 (77%)	46 (23%)	1 1

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	LEU
1	A	6	ASP
1	A	11	LEU
1	A	31	GLU
1	A	35	GLN
1	A	42	ASP
1	A	43	ARG
1	A	44	LEU
1	A	48	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	54	LEU
1	A	57	LYS
1	A	61	LEU
1	A	65	VAL
1	A	74	LEU
1	A	79	THR
1	A	87	LYS
1	A	95	ILE
1	A	100	GLN
1	A	101	THR
1	A	118	LYS
1	A	121	HIS
1	A	122	ILE
1	A	130	MET
1	A	137	ILE
1	A	153	ARG
1	A	156	PRO
1	A	168	LEU
1	A	177	ARG
1	A	188	LEU
1	A	196	THR
1	A	197	LEU
1	A	199	MET
1	A	202	ILE
1	A	203	VAL
1	A	210	GLU
1	A	218	LEU
1	A	220	GLU
1	A	222	TRP
1	A	232	MET
1	A	233	LYS
1	A	236	THR
1	A	250	LEU
1	A	253	THR
1	A	254	THR
1	A	256	ASP
1	A	258	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	35	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	48	GLN
1	A	66	GLN
1	A	67	ASN
1	A	100	GLN
1	A	187	ASN
1	A	224	GLN
1	A	265	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	A	500	-	40,48,48	1.30	1 (2%)	44,73,73	3.29	13 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	500	-	-	0/22/62/62	0/5/5/5

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500	NAD	O7N-C7N	5.06	1.34	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	NAD	C2N-C3N-C4N	-9.11	107.75	118.26
2	A	500	NAD	C1B-N9A-C4A	-8.38	112.16	126.64
2	A	500	NAD	O7N-C7N-N7N	-7.43	111.79	122.60
2	A	500	NAD	C4D-O4D-C1D	-5.29	104.31	109.83
2	A	500	NAD	O5D-PN-O1N	-3.56	95.15	109.07
2	A	500	NAD	C5N-C6N-N1N	-3.44	115.14	120.39
2	A	500	NAD	C4B-O4B-C1B	-3.25	106.44	109.83
2	A	500	NAD	O2N-PN-O1N	2.04	122.49	112.14
2	A	500	NAD	C2N-C3N-C7N	3.16	128.53	119.31
2	A	500	NAD	O7N-C7N-C3N	3.26	123.70	119.62
2	A	500	NAD	C3N-C7N-N7N	5.54	124.20	117.76
2	A	500	NAD	C5N-C4N-C3N	6.55	128.03	120.35
2	A	500	NAD	C3N-C2N-N1N	8.91	129.33	120.41

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	NAD	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.