



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 21, 2024 – 01:55 am BST

PDB ID : 9ENT  
Title : SSX structure of Autotaxin in cryogenic conditions  
Authors : Eymery, M.C.; McCarthy, A.A.; Foos, N.; Basu, S.  
Deposited on : 2024-03-13  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	862	
2	B	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	IOD	A	911	-	-	X	-

## 2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 6968 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ectonucleotide pyrophosphatase/phosphodiesterase family member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	785	6424	4066	1110	1197	51	0	12	0

There are 25 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	GLU	deletion	UNP Q64610
A	?	-	ALA	deletion	UNP Q64610
A	?	-	GLU	deletion	UNP Q64610
A	?	-	THR	deletion	UNP Q64610
A	?	-	GLY	deletion	UNP Q64610
A	?	-	LYS	deletion	UNP Q64610
A	?	-	PHE	deletion	UNP Q64610
A	?	-	ARG	deletion	UNP Q64610
A	?	-	GLY	deletion	UNP Q64610
A	?	-	SER	deletion	UNP Q64610
A	?	-	LYS	deletion	UNP Q64610
A	?	-	HIS	deletion	UNP Q64610
A	?	-	GLU	deletion	UNP Q64610
A	?	-	ASN	deletion	UNP Q64610
A	?	-	LYS	deletion	UNP Q64610
A	?	-	LYS	deletion	UNP Q64610
A	?	-	ASN	deletion	UNP Q64610
A	?	-	LEU	deletion	UNP Q64610
A	?	-	ASN	deletion	UNP Q64610
A	?	-	GLY	deletion	UNP Q64610
A	?	-	SER	deletion	UNP Q64610
A	?	-	VAL	deletion	UNP Q64610
A	?	-	GLU	deletion	UNP Q64610
A	?	-	PRO	deletion	UNP Q64610
A	?	-	ARG	deletion	UNP Q64610

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	3	42	24	3	15	0	0	0

- Molecule 3 is 7alpha-hydroxycholesterol (three-letter code: 5JK) (formula: C<sub>27</sub>H<sub>46</sub>O<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
3	A	1	29	27	2	0	0

- Molecule 4 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
4	A	1	Total	C	N	S	0	0
			3	1	1	1		

- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		

- Molecule 6 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	15	Total	I	0	0
			15	15		

- Molecule 8 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	3	Total	Na	0	0
			3	3		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	449	Total 449	O 449	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ectonucleotide pyrophosphatase/phosphodiesterase family member 2



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	53.51Å 61.19Å 65.01Å 102.57° 99.13° 93.67°	Depositor
Resolution (Å)	37.51 – 2.20 44.26 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.0 (37.51-2.20) 98.9 (44.26-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.99 (at 2.00Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.194 , 0.236 0.193 , 0.235	Depositor DCC
$R_{free}$ test set	2644 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.0	Xtrriage
Anisotropy	0.125	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 44.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6968	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.83% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, 5JK, NA, IOD, SCN, NAG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.26	0/6603	0.51	0/8955

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6424	0	6126	105	0
2	B	42	0	37	1	0
3	A	29	0	0	0	0
4	A	3	0	0	0	0
5	A	2	0	0	0	0
6	A	1	0	0	0	0
7	A	15	0	0	4	0
8	A	3	0	0	0	0
9	A	449	0	0	28	0
All	All	6968	0	6163	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (106) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:678:GLY:O	9:A:1001:HOH:O	2.00	0.79
1:A:550:PRO:O	9:A:1002:HOH:O	2.04	0.75
1:A:589:GLY:HA2	1:A:592:LYS:HB2	1.68	0.75
1:A:380:ASP:OD1	9:A:1003:HOH:O	2.07	0.73
1:A:621:TYR:HA	1:A:628:PRO:HA	1.72	0.71
1:A:456:LYS:O	9:A:1004:HOH:O	2.09	0.69
1:A:393:ARG:NH1	9:A:1013:HOH:O	2.25	0.68
1:A:638:GLN:NE2	9:A:1015:HOH:O	2.25	0.68
1:A:208:LYS:NZ	9:A:1017:HOH:O	2.28	0.67
1:A:682:PRO:HB3	1:A:716:GLN:HB3	1.77	0.67
1:A:618:GLU:OE2	9:A:1005:HOH:O	2.13	0.66
1:A:565:GLY:HA3	1:A:667:LEU:HG	1.77	0.65
1:A:702:VAL:O	9:A:1001:HOH:O	2.13	0.65
1:A:149[B]:GLU:HG3	1:A:349:ARG:NH1	2.14	0.63
1:A:770:ILE:N	9:A:1037:HOH:O	2.32	0.63
1:A:549:ARG:HH11	1:A:550:PRO:HD2	1.63	0.63
1:A:199:PRO:HG2	1:A:502:PRO:HG3	1.82	0.62
1:A:549:ARG:NH1	9:A:1002:HOH:O	2.33	0.61
1:A:430:LYS:NZ	9:A:1044:HOH:O	2.34	0.60
1:A:678:GLY:N	9:A:1001:HOH:O	2.34	0.60
1:A:481:ASN:HA	1:A:484:GLN:HG2	1.85	0.59
1:A:121[A]:GLU:OE2	9:A:1006:HOH:O	2.17	0.59
1:A:104:LYS:NZ	9:A:1049:HOH:O	2.36	0.58
1:A:820:LYS:NZ	9:A:1045:HOH:O	2.34	0.58
1:A:309:GLN:HG3	1:A:310:PRO:HA	1.86	0.58
1:A:817[A]:GLU:OE1	9:A:1007:HOH:O	2.17	0.57
1:A:558:LEU:HB2	1:A:561:GLU:OE2	2.05	0.56
1:A:144:VAL:HG21	1:A:186:PRO:HB2	1.87	0.56
1:A:77:ASN:HD21	1:A:291:TRP:HH2	1.54	0.56
1:A:596:LEU:HD21	1:A:600:ARG:HB3	1.87	0.56
1:A:535:ARG:HG3	1:A:536:THR:HG23	1.89	0.55
1:A:455:ARG:NH2	9:A:1003:HOH:O	2.39	0.54
1:A:550:PRO:HB2	1:A:611:ILE:HG12	1.89	0.54
1:A:793:PHE:CD1	1:A:795:LEU:HG	2.43	0.54
1:A:702:VAL:N	9:A:1001:HOH:O	2.32	0.54
1:A:595:HIS:HB3	1:A:633:TYR:HA	1.90	0.54
1:A:804:SER:HB2	1:A:807:SER:HB3	1.88	0.54
1:A:820:LYS:NZ	9:A:1007:HOH:O	2.35	0.54
1:A:207:THR:HA	1:A:212:ASN:HD21	1.73	0.54
1:A:415:LYS:HG2	1:A:416:PRO:HD2	1.90	0.53

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:ILE:HG12	1:A:349:ARG:NH1	2.23	0.53
1:A:222:PRO:HA	1:A:225:HIS:CE1	2.44	0.52
1:A:312:PHE:CE2	1:A:316:LYS:HE2	2.44	0.52
1:A:546[B]:GLU:OE1	1:A:600:ARG:NH2	2.43	0.52
1:A:78:LEU:HD11	1:A:274:PHE:HB2	1.92	0.51
1:A:457:PRO:HD2	7:A:913:IOD:I	2.80	0.51
1:A:535:ARG:NH1	9:A:1029:HOH:O	2.38	0.51
1:A:550:PRO:HB3	1:A:609:TYR:CZ	2.46	0.51
1:A:589:GLY:HA2	1:A:592:LYS:HE2	1.93	0.50
1:A:74:ARG:NH2	1:A:76:ASP:OD1	2.45	0.50
1:A:636:SER:HA	7:A:917:IOD:I	2.82	0.50
1:A:658:VAL:HG21	1:A:699:THR:HG21	1.93	0.50
1:A:704:MET:HA	1:A:797:HIS:NE2	2.28	0.49
1:A:355:PHE:HB3	1:A:488:VAL:HB	1.94	0.49
1:A:715:PHE:HA	1:A:719:LEU:HB2	1.94	0.48
1:A:549:ARG:NH1	1:A:550:PRO:HD2	2.27	0.48
1:A:733:ILE:N	9:A:1037:HOH:O	2.43	0.48
1:A:737:ILE:HB	1:A:766:HIS:HB2	1.95	0.48
1:A:377:THR:HG23	7:A:911:IOD:I	2.84	0.48
1:A:622:SER:N	1:A:627[B]:MET:O	2.42	0.48
1:A:406:THR:O	1:A:410:ASN:ND2	2.45	0.47
1:A:596:LEU:O	9:A:1008:HOH:O	2.20	0.47
1:A:588:LYS:HE2	9:A:1030:HOH:O	2.13	0.47
1:A:182:SER:H	1:A:189:GLU:HG3	1.80	0.47
1:A:243:LEU:HD12	1:A:244:ARG:HG3	1.97	0.47
1:A:548:SER:O	9:A:1009:HOH:O	2.20	0.46
1:A:281:HIS:NE2	1:A:308:GLU:OE2	2.48	0.46
1:A:508:LEU:HD23	1:A:508:LEU:HA	1.80	0.46
1:A:200:TYR:HB2	1:A:484:GLN:HB3	1.97	0.46
1:A:635:ILE:HG13	1:A:685:LEU:HD13	1.96	0.46
1:A:134:GLN:HB3	1:A:140:GLU:HG3	1.98	0.45
1:A:243:LEU:O	1:A:248:LYS:NZ	2.37	0.45
1:A:677:TYR:HB2	9:A:1001:HOH:O	2.16	0.45
1:A:597:LEU:HD12	1:A:731:ASN:HB2	1.99	0.45
1:A:769:SER:HB3	1:A:793:PHE:CZ	2.52	0.45
1:A:540[A]:ARG:HA	1:A:540[A]:ARG:HE	1.82	0.44
1:A:704:MET:HE2	1:A:709:LYS:HG2	1.99	0.44
1:A:77:ASN:HB3	1:A:274:PHE:O	2.17	0.44
1:A:795:LEU:HD21	1:A:819:MET:HG2	1.99	0.44
1:A:124:LEU:HD22	1:A:129:CYS:SG	2.58	0.44
1:A:682:PRO:HG2	1:A:685:LEU:HD12	2.00	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:3:NAG:H83	2:B:3:NAG:O3	2.17	0.44
1:A:269:ARG:CZ	1:A:269:ARG:HB2	2.48	0.44
1:A:176:SER:HA	1:A:179:LYS:HE3	2.00	0.44
1:A:553:PRO:HD3	1:A:610:ASP:HB3	2.00	0.44
1:A:717:ARG:HG3	1:A:718:VAL:HG23	1.99	0.44
1:A:564:LEU:HD21	1:A:663:SER:HB2	2.00	0.44
1:A:460:VAL:HG13	9:A:1004:HOH:O	2.18	0.43
1:A:371:PHE:CE1	1:A:457:PRO:HA	2.53	0.43
1:A:205:TYR:CD1	1:A:206:PRO:HA	2.54	0.43
1:A:379:VAL:HG23	9:A:1003:HOH:O	2.16	0.43
1:A:286:LEU:HD12	1:A:286:LEU:HA	1.93	0.42
1:A:70:PRO:HA	1:A:71:PRO:HA	1.80	0.42
1:A:370:GLU:OE2	1:A:415:LYS:NZ	2.33	0.42
1:A:376:LEU:HD22	7:A:911:IOD:I	2.90	0.42
1:A:310:PRO:HB2	1:A:324:MET:HE1	2.01	0.42
1:A:92:LEU:O	1:A:95:LYS:HE2	2.20	0.42
1:A:231:SER:HA	1:A:241:PHE:O	2.20	0.42
1:A:393:ARG:HG3	1:A:441:GLU:HG3	2.02	0.41
1:A:646:GLU:HA	1:A:649:THR:HG23	2.02	0.41
1:A:680:LEU:O	1:A:716:GLN:NE2	2.54	0.41
1:A:682:PRO:CG	1:A:685:LEU:HD12	2.50	0.41
1:A:382:ILE:HG23	1:A:392:ILE:HG23	2.02	0.41
1:A:452:HIS:NE2	1:A:473:ASP:OD2	2.48	0.41
1:A:798:ARG:NH1	1:A:799:PRO:HD2	2.37	0.40
1:A:77:ASN:ND2	1:A:272:THR:HG23	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	791/862 (92%)	758 (96%)	30 (4%)	3 (0%)	34 37

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	468[A]	CYS
1	A	468[B]	CYS
1	A	464	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	716/780 (92%)	701 (98%)	15 (2%)	53 67

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	ARG
1	A	74	ARG
1	A	147	ASP
1	A	152	LYS
1	A	286	LEU
1	A	389	LEU
1	A	417	ASP
1	A	479	LYS
1	A	540[A]	ARG
1	A	540[B]	ARG
1	A	651	CYS
1	A	665	ASN
1	A	746	ARG
1	A	775	LEU
1	A	813	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	A	671	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	1,2	14,14,15	0.25	0	17,19,21	0.64	0
2	NAG	B	2	2	14,14,15	0.35	0	17,19,21	1.24	3 (17%)
2	NAG	B	3	2	14,14,15	0.29	0	17,19,21	1.91	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	B	3	2	-	5/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	NAG	C2-N2-C7	7.03	132.92	122.90
2	B	2	NAG	C4-C3-C2	2.95	115.34	111.02

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	O4-C4-C3	-2.54	104.48	110.35
2	B	2	NAG	O4-C4-C5	-2.51	103.07	109.30

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	3	NAG	C3-C2-N2-C7
2	B	3	NAG	C8-C7-N2-C2
2	B	3	NAG	O7-C7-N2-C2
2	B	3	NAG	O5-C5-C6-O6
2	B	3	NAG	C4-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 21 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SCN	A	902	-	1,2,2	0.88	0	0,1,1	-	-
3	5JK	A	901	-	32,32,32	0.11	0	47,50,50	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5JK	A	901	-	-	1/10/71/71	0/4/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	901	5JK	C21-C20-C22-C23

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	785/862 (91%)	0.27	33 (4%) 36 34	25, 43, 73, 134	1 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	466	GLY	29.9
1	A	465	SER	17.9
1	A	589	GLY	13.7
1	A	397	ILE	8.6
1	A	467	LYS	7.1
1	A	464	PRO	6.8
1	A	461	TYR	5.1
1	A	110	VAL	5.0
1	A	588	LYS	4.6
1	A	468[A]	CYS	4.5
1	A	126	ARG	4.3
1	A	68	VAL	4.3
1	A	646	GLU	4.2
1	A	558	LEU	3.9
1	A	396	SER	3.9
1	A	538	THR	3.7
1	A	158	ALA	3.4
1	A	571	LYS	3.2
1	A	570	ASP	3.1
1	A	658	VAL	2.9
1	A	559	GLN	2.8
1	A	469	PHE	2.7
1	A	563	ASP	2.7
1	A	554	GLY	2.6
1	A	83	SER	2.5
1	A	67	GLU	2.5
1	A	59	LYS	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	416	PRO	2.3
1	A	562	PHE	2.2
1	A	103	THR	2.2
1	A	591	THR	2.2
1	A	463	LYS	2.1
1	A	417	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	3	14/15	0.65	0.22	73,93,98,103	0
2	NAG	B	2	14/15	0.86	0.17	49,58,70,71	0
2	NAG	B	1	14/15	0.95	0.16	30,40,47,57	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	IOD	A	910	1/1	0.39	0.19	169,169,169,169	0
7	IOD	A	909	1/1	0.42	0.13	176,176,176,176	0
4	SCN	A	902	3/3	0.60	0.23	68,68,71,90	0
7	IOD	A	919	1/1	0.63	0.08	164,164,164,164	0
7	IOD	A	918	1/1	0.75	0.17	143,143,143,143	0
5	ZN	A	903	1/1	0.83	0.11	48,48,48,48	0
7	IOD	A	906	1/1	0.84	0.13	152,152,152,152	0
7	IOD	A	920	1/1	0.90	0.27	183,183,183,183	0
3	5JK	A	901	29/29	0.92	0.17	25,37,50,66	0
7	IOD	A	911	1/1	0.93	0.11	124,124,124,124	0
7	IOD	A	907	1/1	0.94	0.16	138,138,138,138	0
8	NA	A	921	1/1	0.94	0.08	58,58,58,58	0
8	NA	A	923	1/1	0.94	0.14	59,59,59,59	0
5	ZN	A	904	1/1	0.95	0.04	77,77,77,77	0

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
7	IOD	A	908	1/1	0.95	0.15	58,58,58,58	1
7	IOD	A	913	1/1	0.96	0.09	59,59,59,59	1
8	NA	A	922	1/1	0.97	0.05	40,40,40,40	0
7	IOD	A	914	1/1	0.97	0.09	57,57,57,57	1
7	IOD	A	917	1/1	0.98	0.13	51,51,51,51	1
7	IOD	A	915	1/1	0.99	0.06	68,68,68,68	1
7	IOD	A	916	1/1	0.99	0.12	47,47,47,47	1
6	CA	A	905	1/1	0.99	0.16	38,38,38,38	0
7	IOD	A	912	1/1	1.00	0.15	37,37,37,37	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.