



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 31, 2020 – 09:53 AM BST

PDB ID : 1EO2
Title : CRYSTAL STRUCTURE OF ACINETOBACTER SP. ADP1 PROTOCAT-
ECHUATE 3,4-DIOXYGENASE
Authors : Vetting, M.W.; D'Argenio, D.A.; Ornston, L.N.; Ohlendorf, D.H.
Deposited on : 2000-03-21
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.13
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	209	 72% 23% . .
2	B	241	 74% 23% . .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3712 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTOCATECHUATE 3,4-DIOXYGENASE ALPHA CHAIN.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	202	1602	1010	285	307	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	60	GLY	SER	SEE REMARK 999	UNP P20371

- Molecule 2 is a protein called PROTOCATECHUATE 3,4-DIOXYGENASE BETA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	238	1926	1224	345	352	5	0	0	0

- Molecule 3 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Fe	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	66	Total	O	0	0
			66	66		
4	B	117	Total	O	0	0
			117	117		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PROTOCATECHUATE 3,4-DIOXYGENASE ALPHA CHAIN

Chain A: 



- Molecule 2: PROTOCATECHUATE 3,4-DIOXYGENASE BETA CHAIN

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	145.50Å 145.50Å 145.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.25 25.72 – 2.27	Depositor EDS
% Data completeness (in resolution range)	86.8 (20.00-2.25) 86.8 (25.72-2.27)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.48 (at 2.26Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.176 , 0.218 0.272 , 0.295	Depositor DCC
R_{free} test set	1020 reflections (4.26%)	wwPDB-VP
Wilson B-factor (Å ²)	18.1	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.035 for -l,-k,-h	Xtrriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	3712	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/1638	0.63	0/2223
2	B	0.33	0/1981	0.69	2/2691 (0.1%)
All	All	0.32	0/3619	0.66	2/4914 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	451	ASN	N-CA-C	-6.40	93.71	111.00
2	B	452	ARG	N-CA-C	-6.02	94.74	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1602	0	1548	43	0
2	B	1926	0	1857	49	0
3	B	1	0	0	0	0
4	A	66	0	0	1	0
4	B	117	0	0	0	0
All	All	3712	0	3405	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:369:LYS:H	2:B:422:ASN:HD22	1.21	0.84
2:B:410:HIS:HD2	2:B:412:ASN:HB2	1.44	0.80
1:A:64:ARG:HD3	1:A:99:PHE:O	1.88	0.74
2:B:369:LYS:H	2:B:422:ASN:ND2	1.86	0.74
1:A:37:ASN:HB3	1:A:95:THR:HB	1.73	0.70
1:A:68:ILE:HD12	1:A:130:ILE:HD13	1.72	0.69
2:B:428:ARG:C	2:B:429:MET:HG3	2.15	0.66
1:A:28:ILE:HG22	1:A:30:VAL:HG23	1.79	0.65
2:B:453:ILE:HD12	2:B:454:ASN:H	1.64	0.63
1:A:133:ARG:HG2	2:B:326:THR:HG21	1.81	0.61
1:A:5:LEU:N	1:A:5:LEU:HD12	2.17	0.60
1:A:95:THR:HG22	1:A:106:PHE:CZ	2.37	0.59
2:B:403:ASN:HD22	2:B:403:ASN:C	2.05	0.59
2:B:376:GLU:OE1	2:B:519:ARG:NH2	2.36	0.59
2:B:309:ALA:HB3	2:B:387:GLN:OE1	2.03	0.59
1:A:21:LEU:HD12	1:A:67:LEU:HD22	1.85	0.58
2:B:453:ILE:HD12	2:B:454:ASN:N	2.19	0.57
1:A:68:ILE:HD12	1:A:130:ILE:CD1	2.34	0.57
1:A:67:LEU:HD21	1:A:94:ARG:HD2	1.84	0.57
2:B:315:ASP:HB3	2:B:500:GLN:NE2	2.21	0.56
2:B:428:ARG:O	2:B:429:MET:HG3	2.07	0.54
1:A:177:ARG:O	1:A:178:ASP:HB3	2.08	0.54
2:B:387:GLN:HG2	2:B:529:GLY:N	2.23	0.54
2:B:403:ASN:ND2	2:B:405:SER:H	2.07	0.53
2:B:486:ILE:HG23	2:B:492:LEU:CD2	2.39	0.53
1:A:69:GLU:OE2	1:A:94:ARG:HD3	2.08	0.52
1:A:9:PRO:HG2	2:B:504:LEU:HD21	1.91	0.52
1:A:39:LEU:HD11	1:A:93:GLY:HA3	1.90	0.52
2:B:315:ASP:HB3	2:B:500:GLN:HE21	1.75	0.52
2:B:432:ASP:OD2	2:B:436:TYR:HB3	2.10	0.51
1:A:131:PHE:O	1:A:132:ALA:HB2	2.10	0.51
1:A:16:TYR:HD1	1:A:19:ILE:HD12	1.76	0.51
1:A:130:ILE:HD11	1:A:187:ILE:HD13	1.93	0.50
1:A:40:VAL:HG22	1:A:109:ILE:HD13	1.93	0.50
2:B:400:TRP:HA	2:B:425:GLY:O	2.12	0.49
1:A:194:GLU:OE1	2:B:333:LYS:HG3	2.12	0.49
2:B:410:HIS:CD2	2:B:412:ASN:HB2	2.35	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:467:ALA:HB3	2:B:472:GLN:NE2	2.29	0.47
2:B:429:MET:HE2	2:B:439:PHE:HB3	1.95	0.47
1:A:19:ILE:HG23	1:A:26:ALA:CB	2.45	0.47
2:B:507:LEU:N	2:B:507:LEU:HD22	2.30	0.47
1:A:130:ILE:HD11	1:A:187:ILE:CD1	2.45	0.46
1:A:108:THR:OG1	1:A:109:ILE:N	2.49	0.46
2:B:448:PRO:HD3	2:B:456:TRP:CZ3	2.51	0.46
2:B:403:ASN:C	2:B:403:ASN:ND2	2.69	0.46
2:B:429:MET:HE1	2:B:439:PHE:CD1	2.51	0.45
2:B:490:PRO:HA	2:B:493:LYS:HZ2	1.81	0.45
1:A:175:GLU:OE1	1:A:177:ARG:NH2	2.50	0.45
2:B:403:ASN:ND2	2:B:407:ARG:H	2.15	0.45
1:A:118:LYS:HE3	4:A:851:HOH:O	2.17	0.44
2:B:510:LYS:HA	2:B:513:PHE:CE1	2.52	0.44
2:B:370:ASP:H	2:B:422:ASN:HD21	1.66	0.44
1:A:21:LEU:HD12	1:A:67:LEU:CD2	2.46	0.44
1:A:196:VAL:HG13	2:B:337:ILE:HD12	2.00	0.44
1:A:177:ARG:N	1:A:177:ARG:HD2	2.33	0.43
1:A:30:VAL:HG22	2:B:367:TYR:CD1	2.54	0.43
2:B:495:ILE:HA	2:B:496:PRO:HD3	1.87	0.43
1:A:55:VAL:O	1:A:62:PRO:HA	2.18	0.43
2:B:312:ASN:C	2:B:312:ASN:HD22	2.22	0.43
1:A:98:ASP:OD2	1:A:101:THR:HG23	2.19	0.43
1:A:128:LEU:HD23	1:A:129:ILE:N	2.33	0.43
2:B:468:ASP:H	2:B:472:GLN:HE22	1.66	0.43
1:A:11:GLN:HE22	2:B:492:LEU:HA	1.84	0.42
2:B:368:ALA:HB1	2:B:371:GLY:O	2.19	0.42
2:B:363:LEU:N	2:B:363:LEU:HD12	2.35	0.42
2:B:410:HIS:CD2	2:B:412:ASN:H	2.37	0.42
2:B:441:THR:OG1	2:B:442:ILE:N	2.53	0.42
1:A:38:ASN:ND2	1:A:40:VAL:H	2.17	0.42
1:A:137:ILE:HD11	2:B:317:PRO:O	2.20	0.41
1:A:143:VAL:HG22	1:A:185:PHE:CE2	2.54	0.41
1:A:129:ILE:HD13	1:A:131:PHE:CZ	2.55	0.41
1:A:155:ASP:HA	1:A:156:PRO:HD3	1.79	0.41
2:B:343:LEU:HA	2:B:346:VAL:HG22	2.02	0.41
1:A:188:ARG:HH11	1:A:188:ARG:HG3	1.85	0.41
2:B:510:LYS:HA	2:B:513:PHE:CD1	2.55	0.41
1:A:79:TYR:O	1:A:81:SER:N	2.51	0.41
1:A:155:ASP:O	1:A:159:ASN:ND2	2.54	0.41
2:B:469:GLY:H	2:B:472:GLN:NE2	2.19	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48:ARG:NH1	1:A:109:ILE:HD11	2.36	0.40
2:B:387:GLN:HG2	2:B:529:GLY:H	1.84	0.40
1:A:53:GLY:HA3	1:A:185:PHE:O	2.21	0.40
2:B:400:TRP:CE2	2:B:462:HIS:CB	3.04	0.40
2:B:489:CYS:HA	2:B:490:PRO:HD3	1.83	0.40
1:A:5:LEU:CD1	1:A:5:LEU:N	2.83	0.40
2:B:353:ALA:HA	2:B:356:PHE:HD2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	200/209 (96%)	191 (96%)	8 (4%)	1 (0%)	29 29
2	B	236/241 (98%)	221 (94%)	15 (6%)	0	100 100
All	All	436/450 (97%)	412 (94%)	23 (5%)	1 (0%)	47 55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	178	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/175 (97%)	167 (99%)	2 (1%)	71	80
2	B	201/204 (98%)	195 (97%)	6 (3%)	41	50
All	All	370/379 (98%)	362 (98%)	8 (2%)	52	61

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	A	177	ARG
2	B	312	ASN
2	B	333	LYS
2	B	381	HIS
2	B	403	ASN
2	B	412	ASN
2	B	429	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	18	HIS
1	A	33	HIS
1	A	38	ASN
1	A	72	GLN
2	B	312	ASN
2	B	403	ASN
2	B	410	HIS
2	B	412	ASN
2	B	422	ASN
2	B	472	GLN
2	B	499	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.