



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 23, 2024 – 04:04 PM EST

PDB ID : 8EO0
Title : Crystal structure of alpha-COPI WD40 domain R300A mutant.
Authors : Dey, D.; Hasan, S.S.
Deposited on : 2022-10-01
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	2% 83% 7% 10%
1	B	337	3% 84% 8% 8%
1	C	337	4% 79% 10% 11%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 15177 atoms, of which 7203 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative coatomer subunit alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	303	4852	1577	2385	443	430	17	0	0	0
1	B	311	4964	1610	2443	453	440	18	0	0	0
1	C	301	4827	1569	2375	441	425	17	0	0	0

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	181	LYS	LEU	conflict	UNP Q96WV5
A	185	LYS	LEU	conflict	UNP Q96WV5
A	192	LYS	ILE	conflict	UNP Q96WV5
A	196	LYS	LEU	conflict	UNP Q96WV5
A	197	LYS	PHE	conflict	UNP Q96WV5
A	300	ALA	ARG	engineered mutation	UNP Q96WV5
A	328	SER	-	expression tag	UNP Q96WV5
A	329	ALA	-	expression tag	UNP Q96WV5
A	330	TRP	-	expression tag	UNP Q96WV5
A	331	SER	-	expression tag	UNP Q96WV5
A	332	HIS	-	expression tag	UNP Q96WV5
A	333	PRO	-	expression tag	UNP Q96WV5
A	334	GLN	-	expression tag	UNP Q96WV5
A	335	PHE	-	expression tag	UNP Q96WV5
A	336	GLU	-	expression tag	UNP Q96WV5
A	337	LYS	-	expression tag	UNP Q96WV5
B	181	LYS	LEU	conflict	UNP Q96WV5
B	185	LYS	LEU	conflict	UNP Q96WV5
B	192	LYS	ILE	conflict	UNP Q96WV5
B	196	LYS	LEU	conflict	UNP Q96WV5
B	197	LYS	PHE	conflict	UNP Q96WV5
B	300	ALA	ARG	engineered mutation	UNP Q96WV5
B	328	SER	-	expression tag	UNP Q96WV5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	329	ALA	-	expression tag	UNP Q96WV5
B	330	TRP	-	expression tag	UNP Q96WV5
B	331	SER	-	expression tag	UNP Q96WV5
B	332	HIS	-	expression tag	UNP Q96WV5
B	333	PRO	-	expression tag	UNP Q96WV5
B	334	GLN	-	expression tag	UNP Q96WV5
B	335	PHE	-	expression tag	UNP Q96WV5
B	336	GLU	-	expression tag	UNP Q96WV5
B	337	LYS	-	expression tag	UNP Q96WV5
C	181	LYS	LEU	conflict	UNP Q96WV5
C	185	LYS	LEU	conflict	UNP Q96WV5
C	192	LYS	ILE	conflict	UNP Q96WV5
C	196	LYS	LEU	conflict	UNP Q96WV5
C	197	LYS	PHE	conflict	UNP Q96WV5
C	300	ALA	ARG	engineered mutation	UNP Q96WV5
C	328	SER	-	expression tag	UNP Q96WV5
C	329	ALA	-	expression tag	UNP Q96WV5
C	330	TRP	-	expression tag	UNP Q96WV5
C	331	SER	-	expression tag	UNP Q96WV5
C	332	HIS	-	expression tag	UNP Q96WV5
C	333	PRO	-	expression tag	UNP Q96WV5
C	334	GLN	-	expression tag	UNP Q96WV5
C	335	PHE	-	expression tag	UNP Q96WV5
C	336	GLU	-	expression tag	UNP Q96WV5
C	337	LYS	-	expression tag	UNP Q96WV5

- Molecule 2 is ACETYL GROUP (three-letter code: ACE) (formula: C₂H₄O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 3 2 1	0	0
2	B	1	Total C O 3 2 1	0	0
2	C	1	Total C O 3 2 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	186	Total O 186 186	0	0
3	B	176	Total O 176 176	0	0
3	C	163	Total O 163 163	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	37.94Å 171.33Å 71.58Å 90.00° 100.28° 90.00°	Depositor
Resolution (Å)	28.97 – 1.80 28.97 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (28.97-1.80) 99.9 (28.97-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.18	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.80Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.177 , 0.218 0.177 , 0.218	Depositor DCC
R_{free} test set	4061 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtrriage
Anisotropy	0.087	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.48 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	15177	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.34 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.8181e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/2542	0.67	0/3454
1	B	0.36	0/2597	0.66	0/3528
1	C	0.34	0/2527	0.64	0/3433
All	All	0.35	0/7666	0.66	0/10415

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2467	2385	2385	11	0
1	B	2521	2443	2442	14	0
1	C	2452	2375	2374	21	0
2	A	3	0	3	0	0
2	B	3	0	3	0	0
2	C	3	0	3	0	0
3	A	186	0	0	1	1
3	B	176	0	0	0	2
3	C	163	0	0	1	1
All	All	7974	7203	7210	46	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:34:ARG:NH2	3:C:501:HOH:O	2.07	0.83
1:B:136:HIS:CE1	1:B:162:ARG:HD3	2.27	0.70
1:A:228:LEU:HD21	1:A:271:ILE:HG23	1.74	0.70
1:B:228:LEU:HD11	1:B:271:ILE:HG23	1.80	0.63
1:C:34:ARG:NE	1:C:50:ASP:OD1	2.33	0.61
1:A:272:LEU:HD21	1:A:326:LEU:HD11	1.89	0.55
1:C:33:GLY:HA2	1:C:56:VAL:HG23	1.89	0.55
1:C:107:TYR:OH	1:C:170:ARG:HD2	2.06	0.55
1:A:240:ARG:HG2	1:A:247:TRP:CZ2	2.42	0.54
1:B:4:LEU:HG	1:B:6:LYS:HE3	1.88	0.54
1:C:240:ARG:HG2	1:C:247:TRP:CZ2	2.43	0.54
1:A:202:ALA:O	1:A:203:ILE:HD13	2.09	0.53
1:B:66:PRO:HB3	1:B:83:LYS:HD3	1.91	0.53
1:C:34:ARG:CD	1:C:50:ASP:OD1	2.59	0.51
1:C:34:ARG:HA	1:C:34:ARG:HE	1.76	0.50
1:A:202:ALA:C	1:A:203:ILE:HD13	2.31	0.50
1:C:236:VAL:HB	1:C:252:CYS:HB2	1.94	0.50
1:C:228:LEU:HG	1:C:264:PHE:CZ	2.48	0.48
1:A:327:GLU:OE2	1:A:327:GLU:HA	2.13	0.47
1:C:34:ARG:CZ	1:C:50:ASP:OD1	2.62	0.47
1:B:311:ASN:O	1:B:325:LYS:HG2	2.15	0.47
1:B:6:LYS:HD2	1:B:325:LYS:HD2	1.98	0.45
1:A:162:ARG:NH2	3:A:509:HOH:O	2.44	0.45
1:C:101:CYS:HA	1:C:111:LEU:O	2.16	0.45
1:B:6:LYS:HA	1:B:6:LYS:HE2	1.98	0.44
1:B:279:THR:HG22	1:B:295:ARG:HB3	1.98	0.44
1:C:111:LEU:HB2	1:C:145:PHE:CE2	2.53	0.44
1:C:62:HIS:CG	1:C:63:PRO:HD2	2.52	0.44
1:B:101:CYS:HA	1:B:111:LEU:O	2.17	0.43
1:C:143:ALA:HA	1:C:153:VAL:O	2.19	0.43
1:B:20:HIS:CE1	1:B:23:GLN:HG2	2.54	0.43
1:A:101:CYS:HA	1:A:111:LEU:O	2.18	0.43
1:B:42:MET:HB3	1:B:44:THR:HG23	2.01	0.43
1:C:3:MET:SD	1:C:322:MET:CE	3.07	0.42
1:C:269:GLU:OE1	1:C:286:ASN:ND2	2.51	0.42
1:C:36:GLN:HE21	1:C:48:ARG:NH2	2.18	0.41
1:B:104:HIS:CE1	1:B:107:TYR:CD2	3.09	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:47:ASP:OD2	1:B:85:ARG:HD2	2.21	0.41
1:C:263:LEU:C	1:C:263:LEU:HD12	2.41	0.41
1:A:34:ARG:NH2	1:A:48:ARG:HD2	2.36	0.40
1:B:236:VAL:HB	1:B:252:CYS:HB2	2.02	0.40
1:C:68:PHE:CE1	1:C:80:TRP:CD1	3.10	0.40
1:C:230:ALA:HB1	1:C:259:VAL:HB	2.02	0.40
1:C:314:ALA:HA	1:C:322:MET:O	2.22	0.40
1:A:146:HIS:CG	1:A:147:PRO:HD2	2.56	0.40
1:A:263:LEU:C	1:A:263:LEU:HD12	2.42	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:640:HOH:O	3:C:519:HOH:O[2_646]	2.10	0.10
3:A:543:HOH:O	3:B:620:HOH:O[1_454]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/337 (89%)	287 (96%)	12 (4%)	0	100	100
1	B	307/337 (91%)	292 (95%)	15 (5%)	0	100	100
1	C	297/337 (88%)	285 (96%)	12 (4%)	0	100	100
All	All	903/1011 (89%)	864 (96%)	39 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/297 (90%)	263 (98%)	5 (2%)	57	46
1	B	274/297 (92%)	270 (98%)	4 (2%)	65	56
1	C	266/297 (90%)	262 (98%)	4 (2%)	65	56
All	All	808/891 (91%)	795 (98%)	13 (2%)	62	54

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ARG
1	A	50	ASP
1	A	75	TYR
1	A	170	ARG
1	A	288	ARG
1	B	201	ASP
1	B	240	ARG
1	B	245	LYS
1	B	288	ARG
1	C	10	ARG
1	C	12	SER
1	C	288	ARG
1	C	331	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	128	ASN
1	C	128	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACE	B	401	1	1,2,2	0.83	0	1,1,1	0.32	0
2	ACE	C	401	1	1,2,2	0.70	0	1,1,1	0.28	0
2	ACE	A	401	1	1,2,2	0.74	0	1,1,1	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	303/337 (89%)	-0.05	6 (1%) 65 61	15, 22, 37, 61	0
1	B	311/337 (92%)	0.05	11 (3%) 44 38	15, 23, 42, 59	0
1	C	301/337 (89%)	0.01	12 (3%) 38 32	14, 25, 41, 61	0
All	All	915/1011 (90%)	0.01	29 (3%) 47 41	14, 24, 41, 61	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	171	MET	5.4
1	A	330	TRP	4.9
1	B	171	MET	4.7
1	A	200	THR	4.7
1	B	177	VAL	4.6
1	A	171	MET	4.4
1	B	174	ALA	4.4
1	B	10	ARG	3.6
1	B	4	LEU	3.1
1	C	85	ARG	3.0
1	B	176	PRO	2.9
1	A	170	ARG	2.9
1	C	86	LYS	2.9
1	A	201	ASP	2.8
1	C	170	ARG	2.7
1	B	170	ARG	2.7
1	B	85	ARG	2.5
1	C	34	ARG	2.5
1	C	50	ASP	2.4
1	C	169	LEU	2.3
1	A	4	LEU	2.3
1	B	217	TRP	2.3
1	C	17	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	101	CYS	2.1
1	C	10	ARG	2.1
1	C	287	ARG	2.1
1	C	168	GLY	2.1
1	B	173	ASN	2.0
1	B	42	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ACE	A	401	3/3	0.95	0.10	22,22,24,27	0
2	ACE	B	401	3/3	0.97	0.11	25,25,28,29	0
2	ACE	C	401	3/3	0.97	0.07	26,26,27,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ACE A 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ACE B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.