



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 28, 2023 – 08:05 PM EDT

PDB ID : 8EO5
Title : Crystal structure of the class A beta-lactamase precursor LRA-5 from an Alaskan soil metagenome at 1.8 Angstrom resolution
Authors : Power, P.; D'Amico Gonzalez, G.; Centron, D.; Gutkind, G.; Handelsman, J.; Klinke, S.
Deposited on : 2022-10-02
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	 9% 84% 8% • 7%
1	B	290	 7% 86% 6% • 8%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4322 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LRA-5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	270	Total 2025	C 1268	N 370	O 383	S 4	0	0	0
1	B	268	Total 2018	C 1264	N 372	O 378	S 4	0	1	0

- Molecule 2 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	B	1	Total 8	C 4	N 1	O 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	132	Total 132	O 132	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	139	Total 139	O 139	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LRA-5



- Molecule 1: LRA-5



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	74.27Å 162.62Å 71.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.94 – 1.80 48.94 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (48.94-1.80) 99.9 (48.94-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 1.79Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.189 , 0.215 0.189 , 0.215	Depositor DCC
R_{free} test set	4029 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtrriage
Anisotropy	0.055	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.012 for l,-k,h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4322	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TRS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.79	1/2054 (0.0%)	0.86	5/2792 (0.2%)
1	B	0.73	2/2050 (0.1%)	0.88	7/2786 (0.3%)
All	All	0.76	3/4104 (0.1%)	0.87	12/5578 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	254	LYS	CE-NZ	7.66	1.68	1.49
1	B	32	GLU	CG-CD	5.68	1.60	1.51
1	A	191	VAL	CB-CG1	5.12	1.63	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	188	ARG	NE-CZ-NH2	-11.65	114.47	120.30
1	B	188	ARG	NE-CZ-NH1	9.29	124.95	120.30
1	B	90	GLN	CA-CB-CG	8.89	132.96	113.40
1	B	90	GLN	N-CA-CB	7.68	124.42	110.60
1	B	173	ARG	CA-CB-CG	6.28	127.21	113.40
1	A	269	LYS	CB-CA-C	-6.07	98.27	110.40
1	B	90	GLN	CB-CA-C	-6.04	98.33	110.40
1	B	213	ARG	NE-CZ-NH2	-5.95	117.33	120.30
1	A	222	ARG	NE-CZ-NH1	-5.75	117.42	120.30
1	A	112	ARG	NE-CZ-NH2	-5.35	117.63	120.30
1	A	188	ARG	NE-CZ-NH1	5.35	122.97	120.30
1	A	222	ARG	NE-CZ-NH2	5.08	122.84	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2025	0	2099	16	1
1	B	2018	0	2100	8	1
2	B	8	0	12	1	0
3	A	132	0	0	1	0
3	B	139	0	0	1	0
All	All	4322	0	4211	23	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:254:LYS:CE	1:B:254:LYS:NZ	1.68	1.55
1:A:108(A):ALA:HB1	1:A:112:ARG:NH2	1.58	1.17
1:A:108(A):ALA:HB1	1:A:112:ARG:HH22	1.32	0.91
1:A:33:ARG:NH2	3:A:302:HOH:O	2.29	0.66
1:A:23:LYS:HG2	1:A:25:GLU:HB3	1.79	0.64
1:B:172:ASN:OD1	1:B:174:THR:OG1	2.15	0.64
1:A:108(A):ALA:CB	1:A:112:ARG:NH2	2.52	0.52
1:B:213:ARG:HD2	3:B:499:HOH:O	2.10	0.51
1:B:103:ALA:HB3	1:B:105:TRP:CE2	2.47	0.49
1:B:189:LEU:HD23	1:B:189:LEU:C	2.34	0.48
1:A:197:GLU:HB3	2:B:301:TRS:H32	1.97	0.47
1:A:91:LEU:HD22	1:A:120:ALA:HB2	1.96	0.47
1:A:99:PRO:HG3	1:A:112:ARG:O	2.15	0.47
1:A:254:LYS:HD3	1:A:256:GLN:HE21	1.80	0.46
1:A:254:LYS:HD3	1:A:256:GLN:NE2	2.31	0.45
1:A:108(A):ALA:HB1	1:A:112:ARG:HH21	1.68	0.45
1:A:108(A):ALA:O	1:A:112:ARG:NH2	2.50	0.44
1:A:57:THR:O	1:B:153[B]:ARG:HG3	2.19	0.42
1:B:137:LYS:HA	1:B:137:LYS:HD2	1.91	0.42
1:A:137:LYS:HA	1:A:137:LYS:HD2	1.69	0.42
1:A:136:ASP:OD1	1:A:166:VAL:HG23	2.20	0.41
1:A:213:ARG:O	1:A:215:THR:HG23	2.22	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:200:GLN:OE1	1:B:202:GLN:NE2	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:HIS:CD2	1:B:254:LYS:NZ[4_456]	1.84	0.36

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/290 (92%)	263 (98%)	5 (2%)	0	100	100
1	B	267/290 (92%)	261 (98%)	6 (2%)	0	100	100
All	All	535/580 (92%)	524 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/233 (93%)	212 (98%)	5 (2%)	50	37
1	B	216/233 (93%)	213 (99%)	3 (1%)	67	59

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	433/466 (93%)	425 (98%)	8 (2%)	57 48

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	LYS
1	A	72	PHE
1	A	113	ARG
1	A	129	ARG
1	A	185	ASP
1	B	72	PHE
1	B	185	ASP
1	B	293	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TRS	B	301	-	7,7,7	0.64	0	9,9,9	1.52	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TRS	B	301	-	-	3/9/9/9	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	TRS	C1-C-N	2.91	116.65	107.98

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	301	TRS	C2-C-C3-O3
2	B	301	TRS	N-C-C3-O3
2	B	301	TRS	C1-C-C3-O3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	TRS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/290 (93%)	0.33	27 (10%) 7 5	20, 31, 67, 81	0
1	B	268/290 (92%)	0.18	21 (7%) 13 10	21, 32, 68, 85	0
All	All	538/580 (92%)	0.25	48 (8%) 9 7	20, 32, 68, 85	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	105	TRP	8.0
1	B	173	ARG	6.8
1	B	106	THR	6.4
1	B	103(A)	PRO	5.9
1	A	103(A)	PRO	5.7
1	A	171	GLU	5.6
1	B	104	GLY	5.3
1	A	174	THR	4.9
1	A	109	ALA	4.8
1	A	112	ARG	4.6
1	A	108	ASN	4.5
1	B	293	ARG	4.4
1	B	174	THR	4.3
1	A	106	THR	4.2
1	A	105	TRP	4.2
1	A	104	GLY	4.1
1	B	112	ARG	4.0
1	B	108(A)	ALA	3.9
1	B	107	ALA	3.9
1	A	24	ASN	3.8
1	B	108	ASN	3.8
1	B	109	ALA	3.6
1	A	108(A)	ALA	3.4
1	A	173	ARG	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	177	ARG	3.3
1	A	107	ALA	3.2
1	A	110	MET	3.2
1	B	110	MET	3.0
1	B	113	ARG	3.0
1	A	113	ARG	2.9
1	A	102	VAL	2.8
1	A	23	LYS	2.7
1	A	169	PHE	2.7
1	A	172	ASN	2.6
1	B	177	ARG	2.5
1	A	114	PRO	2.5
1	B	103	ALA	2.5
1	A	129	ARG	2.4
1	B	100	ALA	2.4
1	A	269	LYS	2.4
1	A	254	LYS	2.4
1	A	103	ALA	2.4
1	A	101	ASP	2.3
1	A	170	SER	2.2
1	B	29	MET	2.1
1	B	169	PHE	2.1
1	B	171	GLU	2.1
1	B	172	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	TRS	B	301	8/8	0.78	0.23	33,42,43,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.