



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 28, 2023 – 12:08 PM EDT

PDB ID : 8EO7  
Title : Crystal structure of metagenomic beta-lactamase LRA-5 Y69Q/V166E mutant at 2.15 Angstrom resolution  
Authors : Power, P.; D'Amico Gonzalez, G.; Centron, D.; Gutkind, G.; Handelsman, J.; Klinke, S.  
Deposited on : 2022-10-02  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1479 (2.16-2.16)
Clashscore	141614	1585 (2.16-2.16)
Ramachandran outliers	138981	1560 (2.16-2.16)
Sidechain outliers	138945	1559 (2.16-2.16)
RSRZ outliers	127900	1456 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	290	
1	B	290	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3858 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	255	1912	1197	348	363	4	0	0	0
1	B	248	1844	1155	334	352	3	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	69	GLN	TYR	engineered mutation	UNP B5L5W7
A	166	GLU	VAL	engineered mutation	UNP B5L5W7
B	69	GLN	TYR	engineered mutation	UNP B5L5W7
B	166	GLU	VAL	engineered mutation	UNP B5L5W7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	48	Total	O	0	0
			48	48		
2	B	54	Total	O	0	0
			54	54		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: beta-lactamase



- Molecule 1: beta-lactamase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.86Å 162.24Å 70.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.56 – 2.15 48.58 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.4 (40.56-2.15) 98.7 (48.58-2.15)	Depositor EDS
$R_{merge}$	0.25	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.67 (at 2.16Å)	Xtrriage
Refinement program	PHENIX (1.17.1_3660: ???)	Depositor
R, $R_{free}$	0.225 , 0.248 0.225 , 0.246	Depositor DCC
$R_{free}$ test set	2312 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.0	Xtrriage
Anisotropy	0.510	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 40.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.001 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3858	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.52% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/1935	0.53	0/2624
1	B	0.33	0/1865	0.65	6/2529 (0.2%)
All	All	0.31	0/3800	0.59	6/5153 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	88	ARG	CG-CD-NE	-10.07	90.66	111.80
1	B	277	LYS	CB-CG-CD	6.75	129.14	111.60
1	B	29	MET	CG-SD-CE	-5.90	90.75	100.20
1	B	167	ARG	NE-CZ-NH2	5.62	123.11	120.30
1	B	277	LYS	CD-CE-NZ	-5.33	99.44	111.70
1	B	167	ARG	N-CA-CB	-5.33	101.01	110.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1912	0	1989	21	0
1	B	1844	0	1926	9	0
2	A	48	0	0	2	0
2	B	54	0	0	1	0
All	All	3858	0	3915	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:217:GLY:HA3	1:A:235:THR:HG21	1.68	0.75
1:B:136:ASP:OD1	1:B:167:ARG:HB2	1.90	0.71
1:B:28:SER:HB2	1:B:29:MET:SD	2.36	0.66
1:A:151:ARG:NH1	2:A:301:HOH:O	2.30	0.61
1:B:33:ARG:NH2	2:B:301:HOH:O	2.20	0.58
1:A:202:GLN:HG2	1:A:203:HIS:N	2.20	0.56
1:B:83:GLU:OE2	1:B:88:ARG:NH2	2.39	0.55
1:A:125:VAL:HG23	1:A:129:ARG:HD2	1.89	0.54
1:A:136:ASP:OD2	1:A:167:ARG:HG3	2.08	0.54
1:A:91:LEU:HD22	1:A:120:ALA:HB2	1.90	0.52
1:A:202:GLN:HG2	1:A:203:HIS:H	1.73	0.52
1:A:33:ARG:NH1	2:A:303:HOH:O	2.43	0.51
1:A:124:GLU:OE2	1:A:213:ARG:NH2	2.44	0.51
1:A:78:ILE:CD1	1:A:189:LEU:HD21	2.42	0.49
1:A:194:GLN:HB2	1:A:249:ILE:HD13	1.94	0.48
1:A:26:ALA:HB1	1:A:289:ALA:HA	1.95	0.48
1:A:69:GLN:O	1:A:72:PHE:HD1	1.97	0.47
1:B:91:LEU:HD22	1:B:120:ALA:HB2	1.97	0.47
1:B:136:ASP:O	1:B:140:GLN:HG3	2.15	0.46
1:A:167:ARG:HE	1:A:167:ARG:HB2	1.55	0.46
1:A:115:ILE:HG13	1:A:116:ASP:N	2.31	0.45
1:B:81:LEU:O	1:B:85:GLU:HG3	2.18	0.44
1:A:74:LEU:O	1:A:78:ILE:HD12	2.19	0.42
1:B:194:GLN:HB2	1:B:249:ILE:HD13	2.02	0.42
1:A:25:GLU:O	1:A:28:SER:HB3	2.20	0.41
1:A:202:GLN:CG	1:A:203:HIS:N	2.82	0.41
1:A:96:ARG:HD3	1:A:116:ASP:OD1	2.22	0.40
1:A:57:THR:O	1:B:153:ARG:HG2	2.20	0.40
1:A:253:PRO:HB2	1:A:254:LYS:HG2	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/290 (86%)	241 (97%)	8 (3%)	0	100	100
1	B	242/290 (83%)	240 (99%)	2 (1%)	0	100	100
All	All	491/580 (85%)	481 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/233 (89%)	203 (98%)	4 (2%)	57	61
1	B	200/233 (86%)	194 (97%)	6 (3%)	41	40
All	All	407/466 (87%)	397 (98%)	10 (2%)	47	49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	SER
1	A	170	SER
1	A	185	ASP
1	A	254	LYS
1	B	72	PHE
1	B	88	ARG
1	B	94	LYS
1	B	153	ARG

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	B	177	ARG
1	B	185	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	ASN
1	A	192	GLN
1	B	212	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	255/290 (87%)	0.65	30 (11%) <b>4</b> <b>6</b>	31, 42, 84, 100	0
1	B	248/290 (85%)	0.44	17 (6%) <b>16</b> <b>23</b>	33, 44, 72, 102	0
All	All	503/580 (86%)	0.54	47 (9%) <b>8</b> <b>13</b>	31, 43, 79, 102	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	99	PRO	9.3
1	B	169	PHE	7.9
1	A	99	PRO	7.8
1	A	102	VAL	7.4
1	A	113	ARG	7.0
1	B	167	ARG	6.9
1	A	169	PHE	6.9
1	A	170	SER	6.6
1	A	111	TRP	6.6
1	B	100	ALA	5.8
1	B	170	SER	5.6
1	A	110	MET	5.3
1	B	98	THR	5.3
1	B	29	MET	5.3
1	A	101	ASP	4.9
1	A	112	ARG	4.5
1	A	167	ARG	4.5
1	B	114	PRO	4.4
1	A	26	ALA	4.3
1	A	25	GLU	4.3
1	A	28	SER	4.3
1	A	177	ARG	4.1
1	B	177	ARG	3.9
1	B	88	ARG	3.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	115	ILE	3.7
1	A	100	ALA	3.6
1	A	98	THR	3.6
1	A	114	PRO	3.6
1	A	168	GLU	3.5
1	B	132	ASN	3.5
1	A	216	THR	3.5
1	A	115	ILE	3.5
1	A	27	ILE	3.3
1	A	140	GLN	3.2
1	A	218	THR	3.0
1	A	214	ALA	2.9
1	A	40	ALA	2.8
1	A	292	SER	2.7
1	B	168	GLU	2.7
1	A	202	GLN	2.6
1	B	28	SER	2.5
1	B	215	THR	2.5
1	B	129	ARG	2.4
1	A	32	GLU	2.4
1	B	166	GLU	2.3
1	A	29	MET	2.2
1	A	166	GLU	2.2

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.