



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 16, 2024 – 03:06 PM EST

PDB ID : 4ETV
Title : Crystal structure of mouse ryanodine receptor 2 (2699-2904)
Authors : Yuchi, Z.; Lau, K.; Van Petegem, F.
Deposited on : 2012-04-24
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2328 (1.66-1.66)
Clashscore	180529	2515 (1.66-1.66)
Ramachandran outliers	177936	2475 (1.66-1.66)
Sidechain outliers	177891	2475 (1.66-1.66)
RSRZ outliers	164620	2328 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	
1	B	209	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2996 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ryanodine receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	191	1562	1003	262	289	8	0	4	0
1	B	135	1104	706	187	206	5	0	2	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2696	SER	-	expression tag	UNP E9Q401
A	2697	ASN	-	expression tag	UNP E9Q401
A	2698	ALA	-	expression tag	UNP E9Q401
A	2879	ALA	LYS	engineered mutation	UNP E9Q401
B	2696	SER	-	expression tag	UNP E9Q401
B	2697	ASN	-	expression tag	UNP E9Q401
B	2698	ALA	-	expression tag	UNP E9Q401
B	2879	ALA	LYS	engineered mutation	UNP E9Q401

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Cl	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	235	Total	O	0	0
			235	235		
3	B	94	Total	O	0	0
			94	94		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ryanodine receptor 2



- Molecule 1: Ryanodine receptor 2



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.90Å 88.22Å 92.07Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.05 – 1.65 49.05 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.05-1.65) 100.0 (49.05-1.65)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.42 (at 1.65Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.228 , 0.250 0.228 , 0.249	Depositor DCC
R_{free} test set	2921 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtrriage
Anisotropy	0.202	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.015 for -h,l,k	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2996	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/1604	0.68	0/2153
1	B	0.51	0/1127	0.61	0/1516
All	All	0.62	0/2731	0.65	0/3669

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1562	0	1526	5	0
1	B	1104	0	1053	14	0
2	B	1	0	0	0	0
3	A	235	0	0	1	0
3	B	94	0	0	0	0
All	All	2996	0	2579	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2867:HIS:HD2	1:B:2869:LEU:H	1.15	0.95
1:A:2721[B]:ASN:OD1	1:A:2771:ARG:NH1	2.05	0.90
1:B:2867:HIS:CD2	1:B:2869:LEU:H	2.00	0.80
1:B:2845[B]:GLU:HG2	1:B:2873:TYR:CD2	2.20	0.77
1:B:2845[B]:GLU:HG2	1:B:2873:TYR:CE2	2.35	0.61
1:B:2841:GLU:O	1:B:2845[B]:GLU:HG3	2.04	0.57
1:B:2707:THR:HB	1:B:2779:LYS:HB3	1.88	0.54
1:B:2768:GLU:OE1	1:B:2771:ARG:HD2	2.08	0.53
1:B:2855:LYS:HB3	1:B:2870:LEU:HD21	1.92	0.51
1:B:2867:HIS:HD2	1:B:2869:LEU:N	1.96	0.51
1:B:2721:ASN:HD21	1:B:2771:ARG:HH12	1.60	0.49
1:A:2843:MSE:HE3	3:A:3228:HOH:O	2.13	0.49
1:A:2880:GLU:O	1:B:2834:ARG:NH2	2.45	0.46
1:B:2771:ARG:O	1:B:2775:LYS:HG3	2.16	0.45
1:B:2768:GLU:HA	1:B:2771:ARG:HB2	1.99	0.44
1:A:2707:THR:HB	1:A:2779:LYS:HB3	1.98	0.44
1:A:2758:PRO:HD2	1:A:2761:LEU:HD12	2.01	0.43
1:B:2839:MSE:HE1	1:B:2902:VAL:HG12	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	191/209 (91%)	191 (100%)	0	0	100	100
1	B	129/209 (62%)	129 (100%)	0	0	100	100
All	All	320/418 (77%)	320 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/176 (94%)	165 (100%)	0	100	100
1	B	116/176 (66%)	115 (99%)	1 (1%)	75	63
All	All	281/352 (80%)	280 (100%)	1 (0%)	89	84

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2841	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	2699	ASN
1	B	2867	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	183/209 (87%)	0.43	13 (7%) 23 25	10, 23, 37, 49	4 (2%)
1	B	130/209 (62%)	1.62	49 (37%) 1 1	11, 35, 54, 59	2 (1%)
All	All	313/418 (74%)	0.93	62 (19%) 3 4	10, 26, 50, 59	6 (1%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2857	LEU	5.1
1	B	2763	SER	4.6
1	B	2859	LEU	4.5
1	B	2728	HIS	4.3
1	B	2792	ARG	4.2
1	B	2769	ILE	4.0
1	B	2894	PHE	4.0
1	B	2729	ASP	3.9
1	B	2720	ILE	3.8
1	B	2897	ILE	3.8
1	B	2791	THR	3.7
1	B	2860	GLU	3.6
1	B	2829	ASN	3.5
1	A	2799	LEU	3.5
1	B	2716	LEU	3.4
1	B	2854	LYS	3.4
1	B	2772	TRP	3.3
1	B	2899	GLY	3.3
1	B	2771	ARG	3.3
1	B	2828	SER	3.3
1	A	2859	LEU	3.2
1	B	2718	TYR	3.1
1	B	2711	THR	3.1
1	A	2863	GLY	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	2900	TYR	3.0
1	B	2699	ASN	3.0
1	B	2770	TYR	2.9
1	B	2904	ARG	2.9
1	B	2724	ALA	2.8
1	B	2765	LYS	2.8
1	B	2788	ILE	2.7
1	B	2709	ASN	2.7
1	B	2723	TYR	2.7
1	B	2826	ASP	2.7
1	A	2861	SER	2.6
1	B	2727	SER	2.6
1	B	2712	ILE	2.5
1	B	2830	VAL	2.5
1	A	2700	PHE	2.5
1	B	2831	THR	2.4
1	A	2816	ALA	2.4
1	A	2736	LEU	2.4
1	B	2767	LYS	2.4
1	B	2852	ALA	2.4
1	B	2766	GLU	2.3
1	B	2713	PRO	2.3
1	B	2710	ILE	2.3
1	B	2789	GLU	2.3
1	A	2822	PRO	2.3
1	A	2739	GLY	2.3
1	B	2702	PRO	2.2
1	A	2737	ALA	2.2
1	A	2870	LEU	2.2
1	B	2719	PHE	2.2
1	B	2700	PHE	2.1
1	B	2895	LEU	2.1
1	B	2790	ARG	2.1
1	B	2714	GLU	2.1
1	B	2856	LYS	2.1
1	A	2857	LEU	2.0
1	A	2821	SER	2.0
1	B	2708	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CL	B	3001	1/1	0.94	0.10	55,55,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.