

# Full wwPDB X-ray Structure Validation Report (i)

### Jun 11, 2024 – 06:06 PM EDT

PDB ID : 1EW6
Title : THE CRYSTAL STRUCTURE AND AMINO ACID SEQUENCE OF DE-HALOPEROXIDASE FROM AMPHITRITE ORNATA INDICATE COM-MON ANCESTRY WITH GLOBINS
Authors : Lebioda, L.
Deposited on : 2000-04-24
Resolution : 1.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as $543$ be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
$\mathrm{EDS}$	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber $(2001)$
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# {\rm Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
Clashscore	141614	$10184 \ (1.80-1.76)$
Ramachandran outliers	138981	10051 (1.80-1.76)
Sidechain outliers	138945	10050 (1.80-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
1	А	137	85%	14%	•
1	В	137	82%	16%	•



#### 1 EW6

# 2 Entry composition (i)


There are 4 unique types of molecules in this entry. The entry contains 2395 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called DEHALOPEROXIDASE.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	137	Total 1096	C 691	N 189	O 209	S 7	0	1	0
1	В	137	Total 1096	C 691	N 189	O 209	${f S}7$	0	1	0


• Molecule 2 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0
2	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & 1 \end{array}$	0	0

• Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
2	Δ	1	Total	С	Fe	Ν	Ο	0	0
5	A	1	43	34	1	4	4	0	0
9	D	1	Total	С	Fe	Ν	0	0	0
0	D	1	43	34	1	4	4		U

• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	47	$\begin{array}{cc} \text{Total} & \text{O} \\ 47 & 47 \end{array}$	0	0
4	В	60	Total         O           60         60	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: DEHALOPEROXIDASE





# 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	68.50Å 68.40Å 61.00Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	8.00 - 1.78	Depositor
% Data completeness	(Not available) (8.00-1.78)	Depositor
(in resolution range)	(100 available) (8.00-1.18)	Depositor
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS	Depositor
$R, R_{free}$	0.197 , $0.225$	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2395	wwPDB-VP
Average B, all atoms $(Å^2)$	25.0	wwPDB-VP



# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bo	nd lengths	Bond angles		
		RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.51	2/1122~(0.2%)	0.66	2/1504~(0.1%)	
1	В	0.36	0/1122	0.52	0/1504	
All	All	0.44	2/2244~(0.1%)	0.59	2/3008~(0.1%)	

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	Ideal(Å)
1	А	90	SER	C-N	-6.24	1.19	1.34
1	А	91	SER	CA-CB	-6.24	1.43	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	91	SER	N-CA-CB	-12.89	91.16	110.50
1	А	33	ARG	NE-CZ-NH2	5.39	123.00	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1096	0	1072	11	0
1	В	1096	0	1073	25	0

Continued on next page...



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes				
2	А	5	0	0	0	0				
2	В	5	0	0	0	0				
3	А	43	0	30	1	0				
3	В	43	0	30	6	0				
4	А	47	0	0	0	0				
4	В	60	0	0	3	0				
All	All	2395	0	2205	36	0				

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:20:ILE:HD13	1:B:104:LEU:HD13	1.68	0.75
1:B:36:LYS:HD3	1:B:36:LYS:H	1.58	0.69
1:B:113:GLN:HB3	4:B:183:HOH:O	1.96	0.64
1:B:94:THR:HB	1:B:128:VAL:HG13	1.81	0.61
1:B:55[B]:HIS:ND1	3:B:139:HEM:O2D	2.34	0.61
1:B:36:LYS:HD3	1:B:36:LYS:N	2.16	0.59
1:B:82:THR:O	1:B:86:MET:HG3	2.02	0.59
1:A:21:PHE:HZ	1:A:35:PHE:CZ	2.21	0.58
1:B:52:PHE:O	1:B:56:THR:HG23	2.03	0.58
1:B:15:THR:HA	4:B:195:HOH:O	2.03	0.58
1:B:55[B]:HIS:CE1	3:B:139:HEM:O2D	2.57	0.57
1:B:36:LYS:H	1:B:36:LYS:CD	2.19	0.55
1:A:102:VAL:O	1:A:106:GLU:HG3	2.06	0.55
1:A:83:LEU:HD13	3:A:139:HEM:HBB2	1.89	0.54
1:B:20:ILE:HD13	1:B:104:LEU:CD1	2.35	0.54
1:A:27:LYS:HE2	1:A:106:GLU:OE1	2.07	0.54
1:A:30:ASP:OD1	1:A:33:ARG:NH2	2.41	0.53
1:B:47:LYS:NZ	4:B:197:HOH:O	2.42	0.52
1:A:41:LYS:HE2	1:A:49:MET:CE	2.42	0.49
1:A:82:THR:O	1:A:86:MET:HG3	2.13	0.49
1:B:13:LEU:CD2	1:B:61:ASN:OD1	2.60	0.49
1:B:22:LEU:CD1	1:B:47:LYS:HE3	2.43	0.47
1:B:36:LYS:HG2	1:B:37:ASN:H	1.77	0.47
1:B:127:LEU:HD21	3:B:139:HEM:HBB1	1.94	0.47
1:A:94:THR:HB	1:A:128:VAL:HG13	1.95	0.47
1:A:127:LEU:O	1:A:131:LEU:HG	2.17	0.44
1:A:27:LYS:CE	1:A:106:GLU:OE1	2.65	0.44

Continued on next page...



1 EW6
-------

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:55[A]:HIS:HD1	3:B:139:HEM:CGD	2.31	0.44
1:B:22:LEU:HD11	1:B:47:LYS:HE3	1.98	0.44
1:B:13:LEU:HD23	1:B:61:ASN:OD1	2.17	0.43
1:A:52:PHE:O	1:A:56:THR:HG23	2.19	0.42
1:B:55[B]:HIS:CE1	3:B:139:HEM:CGD	3.03	0.41
1:B:88:GLN:HE21	1:B:88:GLN:H	1.67	0.41
1:B:88:GLN:H	1:B:88:GLN:NE2	2.19	0.41
1:B:83:LEU:HD13	3:B:139:HEM:HBB2	2.01	0.41
1:B:21:PHE:HZ	1:B:35:PHE:CZ	2.39	0.41

Continued from previous page...

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowe		Outliers	Percentiles
1	А	136/137~(99%)	132~(97%)	4 (3%)	0	100 100
1	В	136/137~(99%)	131 (96%)	5 (4%)	0	100 100
All	All	272/274~(99%)	263~(97%)	9~(3%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	118/117~(101%)	116~(98%)	2(2%)	60 48
1	В	118/117 (101%)	114 (97%)	4 (3%)	37 20
All	All	236/234~(101%)	230~(98%)	6(2%)	46 31

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	36	LYS
1	А	91	SER
1	В	30	ASP
1	В	36	LYS
1	В	72	ASP
1	В	88	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	44	GLN
1	А	81	ASN
1	А	85	GLN
1	В	37	ASN
1	В	88	GLN
1	В	113	GLN
1	В	118	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



## 5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res Link		Bond lengths			Bond angles		
10101	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
3	HEM	А	139	1	42,50,50	1.85	9 (21%)	46,82,82	1.31	6 (13%)
2	SO4	А	150	-	4,4,4	0.35	0	6,6,6	0.10	0
3	HEM	В	139	4,1	42,50,50	1.84	10 (23%)	46,82,82	1.26	6 (13%)
2	SO4	В	151	-	4,4,4	0.36	0	6,6,6	0.11	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	В	139	4,1	-	9/12/54/54	-
3	HEM	А	139	1	-	5/12/54/54	-

All (19)	bond length	outliers are	listed below:	
----------	-------------	--------------	---------------	--

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	А	139	HEM	CBB-CAB	5.03	1.54	1.30
3	В	139	HEM	CBB-CAB	4.85	1.53	1.30
3	В	139	HEM	C3C-C2C	-4.14	1.34	1.40
3	А	139	HEM	C3C-C2C	-4.13	1.34	1.40
3	А	139	HEM	CBC-CAC	3.97	1.54	1.29
3	В	139	HEM	CBC-CAC	3.97	1.54	1.29
3	В	139	HEM	C3C-CAC	3.34	1.55	1.47
3	В	139	HEM	C3C-C4C	3.25	1.46	1.41
3	А	139	HEM	C3C-CAC	3.22	1.54	1.47
3	А	139	HEM	C3C-C4C	2.91	1.45	1.41
3	А	139	HEM	CAB-C3B	2.88	1.55	1.47
3	В	139	HEM	CAB-C3B	2.78	1.54	1.47
3	А	139	HEM	C2C-C1C	2.71	1.48	1.42

Continued on next page...



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	А	139	HEM	O2A-CGA	-2.32	1.23	1.30
3	В	139	HEM	O2A-CGA	-2.22	1.23	1.30
3	В	139	HEM	CMD-C2D	2.20	1.55	1.50
3	В	139	HEM	C2C-C1C	2.18	1.47	1.42
3	А	139	HEM	O1A-CGA	2.10	1.28	1.22
3	В	139	HEM	CMC-C2C	2.09	1.56	1.51

Continued from previous page...

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Atoms Z		$Ideal(^{o})$
3	А	139	HEM	C4C-CHD-C1D	4.31	128.25	122.56
3	В	139	HEM	C4C-CHD-C1D	4.01	127.85	122.56
3	А	139	HEM	C4B-CHC-C1C	2.96	126.47	122.56
3	В	139	HEM	CMA-C3A-C4A	-2.90	124.20	128.46
3	А	139	HEM	CMA-C3A-C4A	-2.89	124.23	128.46
3	В	139	HEM	C4B-CHC-C1C	2.81	126.26	122.56
3	В	139	HEM	CBB-CAB-C3B	-2.49	115.06	127.53
3	А	139	HEM	CBB-CAB-C3B	-2.45	115.28	127.53
3	В	139	HEM	O1A-CGA-CBA	-2.44	115.34	123.09
3	А	139	HEM	CMC-C2C-C3C	2.20	129.08	124.68
3	В	139	HEM	CMC-C2C-C3C	2.19	129.05	124.68
3	А	139	HEM	O1A-CGA-CBA	-2.18	116.17	123.09

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	139	HEM	C3D-CAD-CBD-CGD
3	А	139	HEM	C2B-C3B-CAB-CBB
3	В	139	HEM	C2B-C3B-CAB-CBB
3	А	139	HEM	C4B-C3B-CAB-CBB
3	В	139	HEM	C3D-CAD-CBD-CGD
3	В	139	HEM	C4B-C3B-CAB-CBB
3	В	139	HEM	C2D-C3D-CAD-CBD
3	В	139	HEM	CAA-CBA-CGA-O1A
3	В	139	HEM	C4D-C3D-CAD-CBD
3	А	139	HEM	CAA-CBA-CGA-O1A
3	В	139	HEM	CAA-CBA-CGA-O2A
3	А	139	HEM	CAA-CBA-CGA-O2A
3	В	139	HEM	CAD-CBD-CGD-O2D
3	В	139	HEM	CAD-CBD-CGD-O1D




There are no ring outliers.

2 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	139	HEM	1	0
3	В	139	HEM	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	А	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	90:SER	С	91:SER	Ν	1.19



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

## 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

