



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 11:14 AM BST

PDB ID : 1EWK
Title : CRYSTAL STRUCTURE OF METABOTROPIC GLUTAMATE RECEPTOR SUBTYPE 1 COMPLEXED WITH GLUTAMATE
Authors : Kunishima, N.; Shimada, Y.; Jingami, H.; Morikawa, K.
Deposited on : 2000-04-26
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	490	 7% 79% 12% 9%
1	B	490	 3% 82% 9% 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	801	-	-	-	X
2	NAG	B	804	-	-	-	X

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 7906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called METABOTROPIC GLUTAMATE RECEPTOR SUBTYPE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	448	Total 3566	C 2261	N 621	O 663	S 21	0	1	0
1	B	449	Total 3575	C 2266	N 623	O 665	S 21	0	1	0

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 14	C 8	N 1	O 5	0	0
2	A	1	Total 14	C 8	N 1	O 5	0	0
2	B	1	Total 14	C 8	N 1	O 5	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	B	1	14	8	1	5	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	B	1	1	1	0	0
3	A	1	1	1	0	0

- Molecule 4 is GLUTAMIC ACID (three-letter code: GLU) (formula: C₅H₉NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	10	5	1	4	0	0
4	B	1	10	5	1	4	0	0

- Molecule 5 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	S			
5	A	1	Total	15	8	2	4	1	0	0
5	B	1	Total	15	8	2	4	1	0	0
5	B	1	Total	15	8	2	4	1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	246	Total	O	0	0
			246	246		
6	B	396	Total	O	0	0
			396	396		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: METABOTROPIC GLUTAMATE RECEPTOR SUBTYPE 1



- Molecule 1: METABOTROPIC GLUTAMATE RECEPTOR SUBTYPE 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	83.38Å 95.22Å 97.45Å 90.00° 114.85° 90.00°	Depositor
Resolution (Å)	17.00 – 2.20 16.99 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.9 (17.00-2.20) 98.5 (16.99-2.20)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	9.02 (at 2.21Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.196 , 0.227 0.189 , 0.220	Depositor DCC
R_{free} test set	3497 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	25.6	Xtrriage
Anisotropy	0.305	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 58.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.017 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7906	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, EPE, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/3646	0.57	1/4931 (0.0%)
1	B	0.35	0/3655	0.60	1/4943 (0.0%)
All	All	0.34	0/7301	0.58	2/9874 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	290	PHE	N-CA-C	-5.65	95.74	111.00
1	A	290	PHE	N-CA-C	-5.63	95.81	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3566	0	3487	42	0
1	B	3575	0	3495	22	0
2	A	28	0	26	0	0
2	B	28	0	26	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	10	0	5	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	10	0	5	1	0
5	A	15	0	18	0	0
5	B	30	0	36	2	0
6	A	246	0	0	1	0
6	B	396	0	0	0	0
All	All	7906	0	7098	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:213:ARG:NH2	1:A:216:LEU:HD23	1.93	0.83
1:B:39:ALA:HB3	1:B:105:ILE:HB	1.69	0.73
1:A:257:HIS:HB2	1:A:280:ARG:HG3	1.69	0.73
1:A:39:ALA:HB3	1:A:105:ILE:HB	1.77	0.66
1:A:289[B]:CYS:HB3	1:A:291:CYS:SG	2.38	0.63
1:A:377:GLN:HA	1:A:390:PHE:CE1	2.34	0.62
1:B:289[B]:CYS:HB3	1:B:291:CYS:SG	2.38	0.62
1:A:287:VAL:HB	1:A:314:LEU:HD23	1.85	0.58
1:A:216:LEU:HD21	1:A:251:GLU:HG3	1.86	0.57
1:A:379:ARG:HD3	1:A:390:PHE:O	2.05	0.57
1:A:199:LYS:HD2	1:B:63:PRO:HB3	1.87	0.56
1:B:183:ILE:HG12	1:B:202:LEU:HG	1.89	0.55
1:A:274:LEU:O	1:A:278:ARG:HG3	2.05	0.55
1:A:384:LEU:HD23	1:A:384:LEU:O	2.07	0.54
1:B:162:PRO:HG3	1:B:167:VAL:HG12	1.89	0.54
1:B:285:ARG:HG2	1:B:285:ARG:O	2.11	0.51
1:A:315:ILE:HA	1:A:338:ILE:O	2.10	0.51
1:B:272:ARG:O	1:B:276:LYS:HG2	2.10	0.51
1:B:188:THR:HG1	4:B:702:GLU:N	2.09	0.50
1:B:379:ARG:NH1	1:B:381:PRO:HG3	2.27	0.50
1:A:62:VAL:HB	1:A:63:PRO:HD3	1.94	0.50
1:B:275:ARG:O	1:B:279:GLU:HG3	2.12	0.49
1:A:82:MET:HG2	1:A:105:ILE:HD11	1.95	0.48
1:B:231:HIS:O	1:B:260:LYS:HA	2.13	0.48
1:B:224:TRP:CD2	1:B:286:VAL:HG21	2.47	0.48
1:A:359:LEU:HD21	1:A:380:LEU:HD21	1.94	0.48
1:B:323:ARG:HH21	1:B:325:GLU:CD	2.16	0.48
1:A:224:TRP:CZ3	1:A:315:ILE:HD11	2.49	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:216:LEU:CD2	1:A:247:LEU:HB3	2.43	0.48
1:A:162:PRO:HG3	1:A:167:VAL:HG12	1.96	0.47
1:A:333:GLU:H	1:A:333:GLU:CD	2.17	0.47
1:A:216:LEU:HD22	1:A:247:LEU:HB3	1.96	0.47
1:A:423:GLY:HA3	1:A:454:LEU:HD23	1.96	0.47
1:A:486:TYR:OH	1:A:490:ASN:HA	2.15	0.47
1:B:331:GLU:HB3	1:B:492:TYR:CD2	2.50	0.46
1:B:62:VAL:HB	1:B:63:PRO:HD3	1.97	0.46
1:A:398:GLU:OE1	1:A:398:GLU:N	2.43	0.45
1:A:179:ASP:O	1:A:447:GLY:HA3	2.17	0.45
1:A:384:LEU:HD23	1:A:384:LEU:C	2.37	0.45
1:A:55:HIS:CE1	1:A:71:ARG:HG2	2.52	0.44
1:A:381:PRO:HD3	1:A:393:VAL:HG13	2.00	0.44
1:A:491:ARG:HG3	1:A:491:ARG:HH11	1.83	0.43
1:B:287:VAL:HB	1:B:314:LEU:HD23	2.00	0.43
1:A:490:ASN:C	1:A:490:ASN:HD22	2.22	0.43
1:B:96:LEU:N	1:B:97:PRO:HD3	2.34	0.43
1:B:264:ASN:HD22	5:B:903:EPE:H72	1.83	0.43
1:A:343:GLN:OE1	1:A:478:ARG:HD2	2.18	0.43
1:A:280:ARG:HH11	1:A:280:ARG:HA	1.83	0.43
1:B:183:ILE:HA	1:B:202:LEU:O	2.18	0.43
1:A:368:PHE:HB3	1:A:369:PRO:HD3	2.01	0.42
1:A:377:GLN:HA	1:A:390:PHE:CD1	2.54	0.42
1:A:275:ARG:O	1:A:279:GLU:HG3	2.19	0.42
1:A:486:TYR:OH	1:A:490:ASN:ND2	2.52	0.42
1:A:195:LYS:HA	1:A:195:LYS:HE2	2.02	0.42
1:A:280:ARG:HB3	1:A:280:ARG:NH1	2.35	0.42
1:A:276:LYS:HD2	6:A:1165:HOH:O	2.18	0.42
1:A:86:LEU:HD11	1:A:103:SER:HB3	2.02	0.41
1:B:222:TYR:OH	1:B:512:ILE:HD12	2.20	0.41
5:B:902:EPE:H102	5:B:902:EPE:H22	1.73	0.41
1:A:299:LEU:O	1:A:303:MET:HG3	2.20	0.41
1:B:500:TRP:CZ2	1:B:503:GLY:HA2	2.55	0.41
1:A:458:SER:HA	1:A:467:VAL:O	2.21	0.41
1:B:227:VAL:HG12	1:B:286:VAL:HB	2.03	0.41
1:A:268:LYS:HA	1:A:268:LYS:HD3	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/490 (91%)	423 (95%)	21 (5%)	1 (0%)	47	55
1	B	446/490 (91%)	431 (97%)	15 (3%)	0	100	100
All	All	891/980 (91%)	854 (96%)	36 (4%)	1 (0%)	51	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	97	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	385/422 (91%)	376 (98%)	9 (2%)	50	63
1	B	386/422 (92%)	375 (97%)	11 (3%)	43	56
All	All	771/844 (91%)	751 (97%)	20 (3%)	50	58

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	MET
1	A	202	LEU
1	A	289[A]	CYS
1	A	289[B]	CYS
1	A	290	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	313	SER
1	A	478	ARG
1	A	488	GLU
1	A	490	ASN
1	B	35	GLN
1	B	82	MET
1	B	202	LEU
1	B	259	ASP
1	B	289[A]	CYS
1	B	289[B]	CYS
1	B	368	PHE
1	B	465	GLU
1	B	488	GLU
1	B	490	ASN
1	B	508	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	170	GLN
1	A	173	ASN
1	A	250	GLN
1	A	257	HIS
1	A	422	HIS
1	A	490	ASN
1	A	501	HIS
1	B	35	GLN
1	B	170	GLN
1	B	173	ASN
1	B	250	GLN
1	B	343	GLN
1	B	490	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 2 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	804	1	14,14,15	0.52	0	17,19,21	0.62	0
5	EPE	B	903	-	15,15,15	1.68	3 (20%)	18,20,20	1.09	2 (11%)
2	NAG	A	801	1	14,14,15	0.58	0	17,19,21	0.69	1 (5%)
2	NAG	B	803	1	14,14,15	0.55	0	17,19,21	0.70	1 (5%)
5	EPE	A	901	-	15,15,15	1.79	3 (20%)	18,20,20	1.09	2 (11%)
2	NAG	A	802	1	14,14,15	0.52	0	17,19,21	0.66	0
5	EPE	B	902	-	15,15,15	1.58	3 (20%)	18,20,20	1.11	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	804	1	-	0/6/23/26	0/1/1/1
5	EPE	B	903	-	-	3/9/19/19	0/1/1/1
2	NAG	A	801	1	-	2/6/23/26	0/1/1/1
2	NAG	B	803	1	-	0/6/23/26	0/1/1/1
5	EPE	A	901	-	-	2/9/19/19	0/1/1/1
2	NAG	A	802	1	-	0/6/23/26	0/1/1/1
5	EPE	B	902	-	-	2/9/19/19	0/1/1/1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	901	EPE	O1S-S	4.34	1.57	1.45
5	A	901	EPE	O2S-S	4.27	1.57	1.45
5	B	903	EPE	O2S-S	4.16	1.57	1.45
5	B	903	EPE	O1S-S	4.09	1.57	1.45
5	B	902	EPE	O1S-S	3.97	1.56	1.45
5	B	902	EPE	O2S-S	3.78	1.56	1.45
5	A	901	EPE	O3S-S	2.95	1.58	1.47
5	B	903	EPE	O3S-S	2.60	1.56	1.47
5	B	902	EPE	O3S-S	2.41	1.56	1.47

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	902	EPE	O2S-S-C10	2.36	109.75	106.92
5	A	901	EPE	O2S-S-C10	2.31	109.70	106.92
5	B	903	EPE	O3S-S-C10	2.28	109.46	105.77
5	B	902	EPE	O3S-S-C10	2.26	109.42	105.77
5	A	901	EPE	O3S-S-C10	2.21	109.34	105.77
5	B	903	EPE	O1S-S-C10	2.20	109.56	106.92
2	B	803	NAG	C2-N2-C7	-2.09	119.92	122.90
2	A	801	NAG	C2-N2-C7	-2.03	120.01	122.90

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	801	NAG	O5-C5-C6-O6
2	A	801	NAG	C4-C5-C6-O6
5	B	903	EPE	C9-C10-S-O3S
5	A	901	EPE	C10-C9-N1-C2
5	B	903	EPE	C9-C10-S-O1S
5	B	903	EPE	C9-C10-S-O2S
5	A	901	EPE	C10-C9-N1-C6
5	B	902	EPE	C8-C7-N4-C5
5	B	902	EPE	C8-C7-N4-C3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	903	EPE	1	0
5	B	902	EPE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	448/490 (91%)	0.04	32 (7%) 16 14	21, 37, 69, 88	0
1	B	449/490 (91%)	-0.19	13 (2%) 51 49	17, 30, 51, 67	0
All	All	897/980 (91%)	-0.08	45 (5%) 28 27	17, 33, 59, 88	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	510	TYR	6.6
1	A	382	GLY	5.7
1	A	397	ASN	5.6
1	A	384	LEU	5.6
1	B	510	TYR	5.6
1	B	488	GLU	5.5
1	A	488	GLU	5.3
1	B	490	ASN	5.2
1	A	490	ASN	5.0
1	B	384	LEU	4.6
1	A	282	PRO	4.2
1	A	391	LYS	4.0
1	B	486	TYR	3.6
1	A	390	PHE	3.6
1	B	385	LEU	3.5
1	A	388	PRO	3.4
1	B	489	ALA	3.4
1	B	491	ARG	3.3
1	B	388	PRO	3.2
1	A	385	LEU	3.2
1	B	382	GLY	3.2
1	A	389	ASN	3.1
1	A	489	ALA	3.0
1	B	35	GLN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	381	PRO	3.0
1	A	383	HIS	2.8
1	A	226	TYR	2.8
1	A	348	ARG	2.8
1	A	277	LEU	2.7
1	A	491	ARG	2.7
1	A	93	PRO	2.7
1	A	395	THR	2.7
1	A	360	ASP	2.5
1	A	94	VAL	2.5
1	A	268	LYS	2.4
1	A	486	TYR	2.4
1	B	282	PRO	2.4
1	A	436	VAL	2.3
1	A	471	GLU	2.2
1	A	403	ASN	2.2
1	B	283	LYS	2.1
1	A	393	VAL	2.1
1	A	124	ARG	2.1
1	A	283	LYS	2.1
1	A	358	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	A	801	14/15	0.65	0.44	64,66,67,67	0
5	EPE	A	901	15/15	0.70	0.27	72,73,74,74	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	A	802	14/15	0.80	0.37	59,61,62,62	0
2	NAG	B	804	14/15	0.80	0.45	61,63,64,64	0
5	EPE	B	903	15/15	0.83	0.37	61,67,69,69	0
2	NAG	B	803	14/15	0.84	0.20	44,46,47,48	0
5	EPE	B	902	15/15	0.87	0.30	34,48,56,58	0
4	GLU	B	702	10/10	0.96	0.10	25,27,32,34	0
3	MG	B	1002	1/1	0.96	0.06	32,32,32,32	0
3	MG	A	1001	1/1	0.97	0.04	42,42,42,42	0
4	GLU	A	701	10/10	0.97	0.12	21,23,25,26	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.