



wwPDB X-ray Structure Validation Summary Report ⓘ

May 13, 2020 – 01:10 am BST

PDB ID : 6EWX
Title : Structure of Pragmin pseudo-kinase reveals a dimerization mechanism to regulate protein tyrosine phosphorylation and nuclear transcription
Authors : Gelin, M.; Allemand, F.; Fournet, A.; Labesse, G.
Deposited on : 2017-11-06
Resolution : 2.77 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.77 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	<p>2% 56% 17% 26%</p>
1	B	484	<p>% 59% 14% 26%</p>

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 5705 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEAK1-related kinase-activating pseudokinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	359	2853	1818	498	514	23	0	1	0
1	B	358	2793	1780	475	513	25	0	5	0

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	885	MET	-	initiating methionine	UNP D3ZMK9
A	886	GLY	-	expression tag	UNP D3ZMK9
A	887	SER	-	expression tag	UNP D3ZMK9
A	888	SER	-	expression tag	UNP D3ZMK9
A	889	HIS	-	expression tag	UNP D3ZMK9
A	890	HIS	-	expression tag	UNP D3ZMK9
A	891	HIS	-	expression tag	UNP D3ZMK9
A	892	HIS	-	expression tag	UNP D3ZMK9
A	893	HIS	-	expression tag	UNP D3ZMK9
A	894	HIS	-	expression tag	UNP D3ZMK9
A	895	SER	-	expression tag	UNP D3ZMK9
A	896	SER	-	expression tag	UNP D3ZMK9
A	897	GLY	-	expression tag	UNP D3ZMK9
A	898	LEU	-	expression tag	UNP D3ZMK9
A	899	VAL	-	expression tag	UNP D3ZMK9
A	900	PRO	-	expression tag	UNP D3ZMK9
A	901	ARG	-	expression tag	UNP D3ZMK9
A	902	GLY	-	expression tag	UNP D3ZMK9
A	903	SER	-	expression tag	UNP D3ZMK9
A	904	HIS	-	expression tag	UNP D3ZMK9
A	905	MET	-	expression tag	UNP D3ZMK9
B	885	MET	-	initiating methionine	UNP D3ZMK9
B	886	GLY	-	expression tag	UNP D3ZMK9
B	887	SER	-	expression tag	UNP D3ZMK9
B	888	SER	-	expression tag	UNP D3ZMK9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	889	HIS	-	expression tag	UNP D3ZMK9
B	890	HIS	-	expression tag	UNP D3ZMK9
B	891	HIS	-	expression tag	UNP D3ZMK9
B	892	HIS	-	expression tag	UNP D3ZMK9
B	893	HIS	-	expression tag	UNP D3ZMK9
B	894	HIS	-	expression tag	UNP D3ZMK9
B	895	SER	-	expression tag	UNP D3ZMK9
B	896	SER	-	expression tag	UNP D3ZMK9
B	897	GLY	-	expression tag	UNP D3ZMK9
B	898	LEU	-	expression tag	UNP D3ZMK9
B	899	VAL	-	expression tag	UNP D3ZMK9
B	900	PRO	-	expression tag	UNP D3ZMK9
B	901	ARG	-	expression tag	UNP D3ZMK9
B	902	GLY	-	expression tag	UNP D3ZMK9
B	903	SER	-	expression tag	UNP D3ZMK9
B	904	HIS	-	expression tag	UNP D3ZMK9
B	905	MET	-	expression tag	UNP D3ZMK9

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	19	Total 19	O 19	0	0
3	B	30	Total 30	O 30	0	0

GLN	GLY	GLY	PRO	GLY	GLU	K1169	L1170	L1171	P1172	I1175	M1178	F1179	L1180	K1181	Q1184	G1187	GLY	THR	THR	ASN	LEU	GLN	GLN	LYS	K1196	S1197	Q1198	E1204	I1205	S1209	Q1210	Y1211	E1216	E1225	E1234	VAL	ARG	ALA	GLN	LEU	ARG	GLU	ARG	ASP	TYR	ARG	R1246	E1247	S1256
I1257	Y1258	H1267	R1296	E1300	T1312	L1313	L1317	D1316	M1319	K1320	R1321	E1334	R1335	E1342	D1343	W1344	Q1348	E1354	P1355	L1362	L1368																												

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	193.43Å 58.92Å 110.64Å 90.00° 118.48° 90.00°	Depositor
Resolution (Å)	55.00 – 2.77 55.00 – 2.77	Depositor EDS
% Data completeness (in resolution range)	88.7 (55.00-2.77) 88.7 (55.00-2.77)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.69 (at 2.77Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.188 , 0.246 0.189 , 0.247	Depositor DCC
R_{free} test set	1236 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	56.8	Xtrriage
Anisotropy	0.068	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 59.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5705	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2915	0.63	1/3938 (0.0%)
1	B	0.45	0/2874	0.64	2/3892 (0.1%)
All	All	0.46	0/5789	0.64	3/7830 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	1362	LEU	CB-CG-CD2	-5.63	101.42	111.00
1	B	1257	LEU	CA-CB-CG	5.59	128.15	115.30
1	A	1273	ASP	CB-CG-OD1	5.35	123.12	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	967	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2853	0	2872	50	0
1	B	2793	0	2777	52	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	19	0	0	1	0
3	B	30	0	0	4	0
All	All	5705	0	5649	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 99 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1246:ARG:HA	1:B:1267:HIS:HB3	1.53	0.90
1:A:957:PHE:HB2	1:A:1065:ILE:HD11	1.52	0.90
1:A:1345:LEU:HD23	1:A:1348:GLN:HE21	1.36	0.88
1:B:918:ILE:HA	1:B:924:THR:HG21	1.65	0.78
1:B:1181:LYS:NZ	3:B:1502:HOH:O	2.16	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	344/484 (71%)	312 (91%)	25 (7%)	7 (2%)	7 22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	349/484 (72%)	324 (93%)	18 (5%)	7 (2%)	7	22
All	All	693/968 (72%)	636 (92%)	43 (6%)	14 (2%)	7	22

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	970	ASN
1	B	993	ILE
1	B	1335	ARG
1	B	1013	SER
1	B	1210	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/424 (75%)	310 (97%)	9 (3%)	43	74
1	B	312/424 (74%)	309 (99%)	3 (1%)	76	91
All	All	631/848 (74%)	619 (98%)	12 (2%)	57	83

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1130	CYS
1	A	1247	GLU
1	B	1198	GLN
1	A	1090	GLU
1	A	1337	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1059	GLN
1	A	1348	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	B	1401	-	4,4,4	0.19	0	6,6,6	0.38	0
2	SO4	A	1401	-	4,4,4	0.15	0	6,6,6	0.34	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	359/484 (74%)	0.06	11 (3%) 49 44	23, 59, 95, 137	26 (7%)
1	B	358/484 (73%)	-0.01	7 (1%) 65 61	23, 56, 106, 133	27 (7%)
All	All	717/968 (74%)	0.03	18 (2%) 57 52	23, 58, 103, 137	53 (7%)

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	915	LEU	3.8
1	B	1013	SER	3.7
1	A	969	CYS	3.4
1	A	1028	VAL	3.3
1	A	964	LEU	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	SO4	B	1401	5/5	0.98	0.11	45,53,59,71	0
2	SO4	A	1401	5/5	0.98	0.12	65,65,71,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.