



# Full wwPDB X-ray Structure Validation Report i

Oct 30, 2023 – 10:05 AM EDT

PDB ID : 8EXZ  
Title : Structure of GDAP1 containing CMT mutant T157P  
Authors : Googins, M.R.; VanDemark, A.P.  
Deposited on : 2022-10-26  
Resolution : 2.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3617 (2.84-2.80)
Clashscore	141614	4060 (2.84-2.80)
Ramachandran outliers	138981	3978 (2.84-2.80)
Sidechain outliers	138945	3980 (2.84-2.80)
RSRZ outliers	127900	3552 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 7332 atoms, of which 3634 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ganglioside-induced differentiation-associated protein 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	221	Total	C	H	N	O	S	0	0	0
			3624	1164	1796	317	338	9			
1	B	227	Total	C	H	N	O	S	0	0	0
			3708	1191	1838	328	342	9			



There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	157	PRO	THR	engineered mutation	UNP O88741
B	157	PRO	THR	engineered mutation	UNP O88741


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ganglioside-induced differentiation-associated protein 1



- Molecule 1: Ganglioside-induced differentiation-associated protein 1



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.59 Å    80.06 Å    85.81 Å 90.00°    90.00°    90.00°	Depositor
Resolution (Å)	55.86 – 2.82 58.54 – 2.82	Depositor EDS
% Data completeness (in resolution range)	98.1 (55.86-2.82) 87.7 (58.54-2.82)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.39 (at 2.81 Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
$R$ , $R_{free}$	0.255 , 0.279 0.260 , 0.271	Depositor DCC
$R_{free}$ test set	991 reflections (7.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	48.4	Xtriage
Anisotropy	0.821	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 57.9	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.52$ , $< L^2 > = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7332	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 55.73 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.0307e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.26	0/1872	0.48	0/2536
1	B	0.28	0/1916	0.48	0/2597
All	All	0.27	0/3788	0.48	0/5133

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1828	1796	1794	23	0
1	B	1870	1838	1836	24	0
All	All	3698	3634	3630	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:279:TYR:O	1:B:283:VAL:HG23	1.89	0.73
1:A:146:LEU:HD12	1:A:205:LEU:HD23	1.74	0.70

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:71:LEU:HD21	1:B:137:THR:OG1	1.95	0.67
1:B:80:LEU:HD22	1:B:93:ILE:HG23	1.79	0.65
1:A:42:LEU:HD23	1:A:249:VAL:HG13	1.83	0.59
1:B:143:HIS:ND1	1:B:202:VAL:HG22	2.18	0.58
1:A:71:LEU:CD1	1:A:137:THR:HG21	2.35	0.57
1:A:71:LEU:HD13	1:A:137:THR:HG21	1.86	0.57
1:A:43:VAL:HG13	1:A:109:LEU:HD12	1.85	0.56
1:A:26:LEU:CD2	1:A:82:HIS:CD2	2.89	0.55
1:B:43:VAL:HG13	1:B:109:LEU:HD12	1.89	0.55
1:B:133:MET:O	1:B:137:THR:HG23	2.08	0.54
1:B:146:LEU:HD12	1:B:205:LEU:HD23	1.88	0.54
1:A:26:LEU:HD21	1:A:82:HIS:CD2	2.44	0.52
1:A:217:ASP:O	1:A:221:THR:HG23	2.09	0.52
1:A:35:PHE:CE1	1:A:253:VAL:HG12	2.45	0.52
1:A:87:ILE:HD12	1:A:87:ILE:N	2.26	0.51
1:B:124:TYR:O	1:B:128:LEU:HD23	2.09	0.51
1:B:46:GLU:OE1	1:B:109:LEU:HD21	2.11	0.50
1:A:26:LEU:HD23	1:A:82:HIS:CD2	2.48	0.48
1:B:71:LEU:HD12	1:B:71:LEU:O	2.14	0.48
1:B:60:LEU:HD23	1:B:60:LEU:O	2.15	0.46
1:A:71:LEU:CD1	1:A:137:THR:CG2	2.93	0.46
1:A:77:VAL:O	1:A:77:VAL:HG23	2.16	0.46
1:B:87:ILE:HD12	1:B:87:ILE:N	2.30	0.46
1:A:279:TYR:O	1:A:283:VAL:HG23	2.16	0.46
1:B:68:PHE:CD1	1:B:141:ILE:HD11	2.51	0.45
1:A:251:LEU:HD23	1:A:279:TYR:CZ	2.53	0.44
1:B:137:THR:HG22	1:B:261:LEU:HD11	1.99	0.43
1:A:146:LEU:HD12	1:A:205:LEU:CD2	2.44	0.43
1:B:255:LEU:HD22	1:B:268:TRP:CZ3	2.53	0.43
1:B:35:PHE:CE1	1:B:253:VAL:HG12	2.54	0.43
1:A:111:PRO:HG2	1:A:118:TYR:HA	2.00	0.43
1:B:143:HIS:CE1	1:B:202:VAL:HG22	2.54	0.43
1:A:88:CYS:O	1:A:89:GLU:HB2	2.18	0.42
1:B:151:MET:HG3	1:B:260:PHE:HA	2.01	0.42
1:A:140:CYS:SG	1:A:147:THR:HG21	2.58	0.42
1:B:77:VAL:O	1:B:77:VAL:HG23	2.19	0.42
1:B:111:PRO:HG2	1:B:118:TYR:HA	2.03	0.41
1:B:287:LYS:HD3	1:B:287:LYS:H	1.86	0.41
1:A:124:TYR:O	1:A:128:LEU:HD23	2.21	0.41
1:A:70:ARG:O	1:A:70:ARG:CG	2.69	0.41
1:B:146:LEU:HD12	1:B:205:LEU:CD2	2.52	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:LEU:CD1	1:A:205:LEU:HD23	2.47	0.40
1:B:251:LEU:HD23	1:B:279:TYR:CZ	2.57	0.40
1:B:280:TYR:CE2	1:B:284:LEU:HD11	2.57	0.40
1:A:106:THR:O	1:A:108:ARG:NH1	2.55	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	215/358 (60%)	208 (97%)	6 (3%)	1 (0%)	29 59
1	B	221/358 (62%)	217 (98%)	4 (2%)	0	100 100
All	All	436/716 (61%)	425 (98%)	10 (2%)	1 (0%)	47 76

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	83	GLY

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	204/318 (64%)	200 (98%)	4 (2%)	55 83

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	B	208/318 (65%)	199 (96%)	9 (4%)	29 60
All	All	412/636 (65%)	399 (97%)	13 (3%)	39 71

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105	ARG
1	A	120	ARG
1	A	140	CYS
1	A	265	ARG
1	B	50	LYS
1	B	71	LEU
1	B	104	GLU
1	B	140	CYS
1	B	141	ILE
1	B	145	GLU
1	B	230	THR
1	B	281	GLU
1	B	287	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	221/358 (61%)	0.92	21 (9%) <span style="background-color: red; color: white; border: 1px solid black; padding: 2px;">8</span> <span style="background-color: red; color: white; border: 1px solid black; padding: 2px;">4</span>	36, 60, 91, 120	0
1	B	227/358 (63%)	1.05	27 (11%) <span style="background-color: red; color: white; border: 1px solid black; padding: 2px;">4</span> <span style="background-color: red; color: white; border: 1px solid black; padding: 2px;">2</span>	38, 61, 85, 96	0
All	All	448/716 (62%)	0.99	48 (10%) <span style="background-color: red; color: white; border: 1px solid black; padding: 2px;">6</span> <span style="background-color: red; color: white; border: 1px solid black; padding: 2px;">3</span>	36, 61, 88, 120	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	295	HIS	4.4
1	A	145	GLU	4.0
1	B	296	VAL	3.9
1	B	297	ASN	3.5
1	B	239	LEU	3.3
1	B	240	CYS	3.3
1	B	237	PRO	3.2
1	B	294	GLY	3.2
1	B	86	ILE	3.1
1	B	118	TYR	3.0
1	A	71	LEU	3.0
1	B	285	LYS	2.9
1	B	71	LEU	2.8
1	A	246	LEU	2.8
1	B	107	PRO	2.8
1	A	199	HIS	2.8
1	B	101	PHE	2.6
1	B	46	GLU	2.6
1	B	70	ARG	2.6
1	B	270	HIS	2.6
1	A	282	ARG	2.6
1	A	148	VAL	2.6
1	A	70	ARG	2.6
1	B	105	ARG	2.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	136	TYR	2.5
1	A	25	HIS	2.5
1	A	265	ARG	2.5
1	A	239	LEU	2.5
1	B	80	LEU	2.5
1	B	25	HIS	2.4
1	A	242	GLU	2.4
1	B	26	LEU	2.4
1	A	69	MET	2.4
1	B	226	ARG	2.4
1	A	73	SER	2.4
1	A	81	VAL	2.3
1	B	231	PRO	2.3
1	B	109	LEU	2.3
1	A	79	VAL	2.3
1	A	105	ARG	2.3
1	A	234	GLY	2.2
1	A	203	LYS	2.1
1	B	272	LYS	2.1
1	B	283	VAL	2.1
1	B	102	LEU	2.0
1	A	285	LYS	2.0
1	A	284	LEU	2.0
1	B	79	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.