



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 26, 2023 – 04:04 AM EDT

PDB ID : 3EYC  
Title : New crystal structure of human tear lipocalin in complex with 1,4-butanediol in space group P21  
Authors : Breustedt, D.A.; Keil, L.; Skerra, A.  
Deposited on : 2008-10-20  
Resolution : 2.60 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	162	 62% 23% 12%
1	B	162	 59% 24% 14%
1	C	162	 57% 27% 15%
1	D	162	 66% 17% 15%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4524 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lipocalin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	143	1106	694	190	216	6	41	0	0
1	B	140	1090	684	187	213	6	49	0	0
1	C	137	1058	660	182	210	6	47	0	0
1	D	137	1061	666	182	208	5	57	0	0

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	101	SER	CYS	engineered mutation	UNP P31025
A	158	ALA	ASP	engineered mutation	UNP P31025
A	159	TRP	-	expression tag	UNP P31025
A	160	SER	-	expression tag	UNP P31025
A	161	HIS	-	expression tag	UNP P31025
A	162	PRO	-	expression tag	UNP P31025
A	163	GLN	-	expression tag	UNP P31025
A	164	PHE	-	expression tag	UNP P31025
A	165	GLU	-	expression tag	UNP P31025
A	166	LYS	-	expression tag	UNP P31025
B	101	SER	CYS	engineered mutation	UNP P31025
B	158	ALA	ASP	engineered mutation	UNP P31025
B	159	TRP	-	expression tag	UNP P31025
B	160	SER	-	expression tag	UNP P31025
B	161	HIS	-	expression tag	UNP P31025
B	162	PRO	-	expression tag	UNP P31025
B	163	GLN	-	expression tag	UNP P31025
B	164	PHE	-	expression tag	UNP P31025
B	165	GLU	-	expression tag	UNP P31025
B	166	LYS	-	expression tag	UNP P31025
C	101	SER	CYS	engineered mutation	UNP P31025

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	158	ALA	ASP	engineered mutation	UNP P31025
C	159	TRP	-	expression tag	UNP P31025
C	160	SER	-	expression tag	UNP P31025
C	161	HIS	-	expression tag	UNP P31025
C	162	PRO	-	expression tag	UNP P31025
C	163	GLN	-	expression tag	UNP P31025
C	164	PHE	-	expression tag	UNP P31025
C	165	GLU	-	expression tag	UNP P31025
C	166	LYS	-	expression tag	UNP P31025
D	101	SER	CYS	engineered mutation	UNP P31025
D	158	ALA	ASP	engineered mutation	UNP P31025
D	159	TRP	-	expression tag	UNP P31025
D	160	SER	-	expression tag	UNP P31025
D	161	HIS	-	expression tag	UNP P31025
D	162	PRO	-	expression tag	UNP P31025
D	163	GLN	-	expression tag	UNP P31025
D	164	PHE	-	expression tag	UNP P31025
D	165	GLU	-	expression tag	UNP P31025
D	166	LYS	-	expression tag	UNP P31025

- Molecule 2 is 1,4-BUTANEDIOL (three-letter code: BU1) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	4	2		
2	B	1	Total	C	O	0	0
			6	4	2		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	C	O	0	0
			6	4	2		
2	D	1	Total	C	O	0	0
			6	4	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	42	Total	O	0	0
			42	42		
3	B	34	Total	O	0	0
			34	34		
3	C	55	Total	O	0	0
			55	55		
3	D	54	Total	O	0	0
			54	54		





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.91Å 34.17Å 143.48Å 90.00° 90.35° 90.00°	Depositor
Resolution (Å)	20.00 – 2.60 19.60 – 2.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.60) 99.5 (19.60-2.60)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.34 (at 2.59Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.228 , 0.274 0.222 , 0.269	Depositor DCC
$R_{free}$ test set	1003 reflections (5.11%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	44.8	Xtrriage
Anisotropy	0.224	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 53.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.032 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4524	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BU1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	1/1126 (0.1%)	0.86	7/1520 (0.5%)
1	B	0.51	0/1108	0.95	10/1494 (0.7%)
1	C	0.49	0/1072	0.71	0/1442
1	D	0.50	0/1076	0.72	0/1448
All	All	0.51	1/4382 (0.0%)	0.82	17/5904 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	14	SER	CA-CB	-6.64	1.43	1.52

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	125	GLU	CB-CA-C	8.66	127.73	110.40
1	B	49	LEU	CB-CA-C	-8.13	94.74	110.20
1	B	48	ASN	CB-CG-OD1	7.94	137.48	121.60
1	A	125	GLU	CG-CD-OE1	7.78	133.87	118.30
1	B	49	LEU	CB-CG-CD1	-7.66	97.97	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	111	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1106	0	1104	31	1
1	B	1090	0	1088	36	0
1	C	1058	0	1056	38	0
1	D	1061	0	1063	27	0
2	A	6	0	10	0	0
2	B	6	0	10	0	0
2	C	6	0	10	2	0
2	D	6	0	10	0	0
3	A	42	0	0	7	0
3	B	34	0	0	3	0
3	C	55	0	0	7	0
3	D	54	0	0	3	0
All	All	4524	0	4351	127	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 127 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:GLU:HB2	3:A:175:HOH:O	1.51	1.10
1:A:125:GLU:CB	3:A:175:HOH:O	1.99	1.08
1:D:44:LEU:HD12	1:D:44:LEU:O	1.56	1.02
1:B:24:VAL:HG22	1:B:143:SER:HB2	1.50	0.94
1:D:44:LEU:HD23	1:D:50:GLU:HB2	1.55	0.89

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:GLU:OE2	1:A:154:SER:OG[1_545]	2.19	0.01

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/162 (87%)	140 (99%)	1 (1%)	0	100	100
1	B	136/162 (84%)	132 (97%)	2 (2%)	2 (2%)	10	21
1	C	131/162 (81%)	128 (98%)	2 (2%)	1 (1%)	19	39
1	D	131/162 (81%)	128 (98%)	3 (2%)	0	100	100
All	All	539/648 (83%)	528 (98%)	8 (2%)	3 (1%)	25	47

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	13	VAL
1	B	30	GLU
1	C	14	SER

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/137 (88%)	113 (93%)	8 (7%)	16	33
1	B	120/137 (88%)	116 (97%)	4 (3%)	38	64

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	116/137 (85%)	110 (95%)	6 (5%)	23	46
1	D	116/137 (85%)	112 (97%)	4 (3%)	37	63
All	All	473/548 (86%)	451 (95%)	22 (5%)	26	50

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	89	ILE
1	C	132	LYS
1	C	95	ASP
1	D	41	LEU
1	A	137	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	84	HIS
1	C	92	HIS
1	D	62	GLN
1	C	96	HIS
1	A	106	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BU1	B	1	-	5,5,5	0.34	0	4,4,4	0.53	0
2	BU1	C	1	-	5,5,5	0.35	0	4,4,4	0.54	0
2	BU1	D	1	-	5,5,5	0.35	0	4,4,4	0.54	0
2	BU1	A	1	-	5,5,5	0.35	0	4,4,4	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BU1	B	1	-	-	0/3/3/3	-
2	BU1	C	1	-	-	0/3/3/3	-
2	BU1	D	1	-	-	0/3/3/3	-
2	BU1	A	1	-	-	0/3/3/3	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	BU1	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	143/162 (88%)	-0.33	1 (0%) 87 86	21, 39, 67, 89	11 (7%)
1	B	140/162 (86%)	-0.16	2 (1%) 75 71	18, 48, 76, 145	13 (9%)
1	C	137/162 (84%)	-0.25	1 (0%) 87 86	17, 42, 74, 98	11 (8%)
1	D	137/162 (84%)	-0.21	2 (1%) 73 70	21, 41, 71, 108	13 (9%)
All	All	557/648 (85%)	-0.24	6 (1%) 80 78	17, 42, 74, 145	48 (8%)

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	29	PRO	4.8
1	D	44	LEU	3.7
1	C	109	PRO	2.7
1	A	92	HIS	2.6
1	D	46	GLY	2.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BU1	A	1	6/6	0.92	0.35	34,36,38,40	0
2	BU1	B	1	6/6	0.92	0.32	56,57,58,58	0
2	BU1	C	1	6/6	0.92	0.36	40,40,41,43	0
2	BU1	D	1	6/6	0.93	0.39	48,49,50,51	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.