



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 21, 2023 – 04:13 AM JST

PDB ID : 7EYU
Title : Fe(II)/(alpha)ketoglutarate-dependent dioxygenase SptF-N65T mutant with andiconin D
Authors : Tao, H.; Mori, T.; Abe, I.
Deposited on : 2021-06-01
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1692 (2.04-2.04)
Clashscore	141614	1773 (2.04-2.04)
Ramachandran outliers	138981	1752 (2.04-2.04)
Sidechain outliers	138945	1752 (2.04-2.04)
RSRZ outliers	127900	1672 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	
1	B	296	

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4637 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-oxoglutarate/Fe(II)-dependent dioxygenase SptF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	265	2141	1358	376	392	15	0	0	0
1	A	270	2162	1367	378	402	15	0	0	0

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	3	MET	-	initiating methionine	UNP A0A6J4CX17
B	65	THR	ASN	engineered mutation	UNP A0A6J4CX17
B	286	LYS	-	expression tag	UNP A0A6J4CX17
B	287	LEU	-	expression tag	UNP A0A6J4CX17
B	288	ALA	-	expression tag	UNP A0A6J4CX17
B	289	ALA	-	expression tag	UNP A0A6J4CX17
B	290	ALA	-	expression tag	UNP A0A6J4CX17
B	291	LEU	-	expression tag	UNP A0A6J4CX17
B	292	GLU	-	expression tag	UNP A0A6J4CX17
B	293	HIS	-	expression tag	UNP A0A6J4CX17
B	294	HIS	-	expression tag	UNP A0A6J4CX17
B	295	HIS	-	expression tag	UNP A0A6J4CX17
B	296	HIS	-	expression tag	UNP A0A6J4CX17
B	297	HIS	-	expression tag	UNP A0A6J4CX17
B	298	HIS	-	expression tag	UNP A0A6J4CX17
A	3	MET	-	initiating methionine	UNP A0A6J4CX17
A	65	THR	ASN	engineered mutation	UNP A0A6J4CX17
A	286	LYS	-	expression tag	UNP A0A6J4CX17
A	287	LEU	-	expression tag	UNP A0A6J4CX17
A	288	ALA	-	expression tag	UNP A0A6J4CX17
A	289	ALA	-	expression tag	UNP A0A6J4CX17
A	290	ALA	-	expression tag	UNP A0A6J4CX17
A	291	LEU	-	expression tag	UNP A0A6J4CX17
A	292	GLU	-	expression tag	UNP A0A6J4CX17
A	293	HIS	-	expression tag	UNP A0A6J4CX17

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	294	HIS	-	expression tag	UNP A0A6J4CX17
A	295	HIS	-	expression tag	UNP A0A6J4CX17
A	296	HIS	-	expression tag	UNP A0A6J4CX17
A	297	HIS	-	expression tag	UNP A0A6J4CX17
A	298	HIS	-	expression tag	UNP A0A6J4CX17

- Molecule 2 is N-OXALYLGLYCINE (three-letter code: OGA) (formula: $C_4H_5NO_5$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	B	1	10	4	1	5	0	0

- Molecule 3 is Andiconin D (three-letter code: OCC) (formula: $C_{25}H_{30}O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	C O	0	0
			29	25 4		

- Molecule 4 is FE (II) ION (three-letter code: FE2) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Fe	0	0
			1	1		
4	A	1	Total	Fe	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	154	Total	O	0	0
			154	154		
5	A	139	Total	O	0	0
			139	139		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	42.81Å 50.58Å 71.98Å 109.11° 91.29° 112.86°	Depositor
Resolution (Å)	27.85 – 2.05 43.40 – 2.05	Depositor EDS
% Data completeness (in resolution range)	96.1 (27.85-2.05) 96.1 (43.40-2.05)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.39 (at 2.05Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.241 , 0.278 0.240 , 0.277	Depositor DCC
R_{free} test set	2017 reflections (6.46%)	wwPDB-VP
Wilson B-factor (Å ²)	12.1	Xtrriage
Anisotropy	1.565	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 25.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.085 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	4637	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OGA, OCC, FE2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/2229	0.44	0/3037
1	B	0.24	0/2210	0.41	0/3011
All	All	0.24	0/4439	0.43	0/6048

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2162	0	2043	34	0
1	B	2141	0	2041	21	0
2	B	10	0	3	1	0
3	B	29	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	139	0	0	2	0
5	B	154	0	0	0	0
All	All	4637	0	4087	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:269:ALA:HB2	1:A:274:LEU:HA	1.74	0.69
1:A:235:HIS:HE1	1:A:267:TRP:HZ2	1.40	0.69
1:A:192:GLN:HG3	1:A:193:PRO:HD2	1.75	0.67
1:A:155:LEU:HD11	1:A:219:PHE:HD1	1.64	0.62
1:A:150:ASN:HD21	1:A:220:ALA:HB1	1.65	0.60
1:B:258:LEU:HB2	1:B:266:VAL:HG23	1.83	0.59
1:B:231:ILE:HD12	1:A:266:VAL:HG12	1.84	0.59
1:A:127:ILE:HA	1:A:206:SER:HB3	1.84	0.58
1:A:265:LYS:O	1:A:268:LYS:NZ	2.33	0.58
1:A:77:SER:O	1:A:84:ARG:HD3	2.04	0.57
1:A:27:THR:HG21	1:A:167:VAL:HG21	1.86	0.55
1:A:262:ASN:ND2	5:A:415:HOH:O	2.40	0.54
1:A:265:LYS:HB2	1:A:268:LYS:HD3	1.90	0.53
1:A:78:LEU:HD23	1:A:237:LEU:HD22	1.91	0.53
1:B:77:SER:O	1:B:84:ARG:NH1	2.37	0.53
1:B:127:ILE:HA	1:B:206:SER:HB3	1.92	0.51
1:B:231:ILE:HG13	1:A:268:LYS:HZ3	1.76	0.51
1:B:139:TRP:HH2	1:A:105:TYR:HH	1.59	0.50
1:B:18:VAL:HG13	1:B:28:LEU:HD21	1.93	0.50
1:B:73:PRO:HD2	1:B:115:ILE:HB	1.95	0.49
1:A:83:TYR:HA	1:A:87:VAL:HB	1.93	0.49
1:A:116:LEU:N	1:A:218:GLY:O	2.45	0.49
1:B:76:LEU:HD11	1:B:115:ILE:HG12	1.95	0.49
1:A:116:LEU:HD11	1:A:152:THR:HG23	1.93	0.48
1:B:72:GLN:NE2	2:B:301:OGA:O2	2.44	0.48
1:A:76:LEU:CD1	1:A:115:ILE:HD11	2.43	0.48
1:B:72:GLN:HA	1:B:115:ILE:O	2.14	0.47
1:A:199:LEU:HD11	1:A:205:HIS:CG	2.50	0.47
1:A:189:VAL:HG12	1:A:191:MET:HG3	1.97	0.47
1:B:71:LYS:HB2	1:B:117:HIS:HB3	1.95	0.47
1:A:90:HIS:CE1	1:A:92:TRP:HB2	2.50	0.46
1:B:53:TRP:CD2	1:B:73:PRO:HB3	2.51	0.46
1:A:116:LEU:HD12	1:A:218:GLY:HA3	1.97	0.46
1:A:199:LEU:HD21	1:A:205:HIS:CE1	2.50	0.46
1:B:17:ASP:OD1	1:B:20:ARG:NH1	2.42	0.46
1:A:118:LEU:HB2	1:A:216:ARG:HB3	1.99	0.45
1:A:156:THR:HG23	1:A:217:ARG:HB2	1.98	0.45
1:A:118:LEU:HD11	1:A:207:ALA:HB1	2.00	0.44
1:A:155:LEU:HD13	1:A:218:GLY:HA2	1.99	0.44
1:B:140:ARG:O	1:B:176:ASN:ND2	2.51	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:90:HIS:CE1	1:B:92:TRP:HB2	2.53	0.43
1:B:245:MET:HB2	1:B:250:GLN:HG3	2.00	0.43
1:A:48:ASP:OD1	5:A:401:HOH:O	2.21	0.42
1:A:239:ARG:HD2	1:A:239:ARG:HA	1.78	0.42
1:B:239:ARG:N	1:B:239:ARG:HD2	2.34	0.42
1:A:76:LEU:HD13	1:A:115:ILE:HD11	2.02	0.42
1:B:76:LEU:HD23	1:B:83:TYR:CE1	2.56	0.41
1:B:235:HIS:CD2	1:A:235:HIS:ND1	2.89	0.41
1:A:160:VAL:HG23	1:A:190:LEU:HD21	2.01	0.41
1:B:233:SER:HB2	1:A:267:TRP:CD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/296 (90%)	263 (99%)	3 (1%)	0	100	100
1	B	263/296 (89%)	255 (97%)	8 (3%)	0	100	100
All	All	529/592 (89%)	518 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/261 (91%)	232 (98%)	6 (2%)	47	40
1	B	238/261 (91%)	229 (96%)	9 (4%)	33	26
All	All	476/522 (91%)	461 (97%)	15 (3%)	39	32

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	48	ASP
1	B	81	LYS
1	B	109	TRP
1	B	131	HIS
1	B	138	LYS
1	B	166	ARG
1	B	216	ARG
1	B	224	HIS
1	B	261	HIS
1	A	68	SER
1	A	69	LYS
1	A	109	TRP
1	A	131	HIS
1	A	166	ARG
1	A	224	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OCC	B	302	-	31,34,34	4.45	10 (32%)	39,63,63	2.24	7 (17%)
2	OGA	B	301	4	9,9,9	1.51	2 (22%)	10,11,11	1.35	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OGA	B	301	4	-	0/8/9/9	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	OCC	CAC-CAB	-13.43	1.39	1.53
3	B	302	OCC	CAT-CAU	-12.28	1.39	1.53
3	B	302	OCC	CAS-CAR	9.15	1.53	1.33
3	B	302	OCC	CAE-CAF	-8.81	1.39	1.51
3	B	302	OCC	CBB-CAR	-6.28	1.39	1.47
3	B	302	OCC	CAP-CAU	-4.94	1.40	1.53
3	B	302	OCC	CAA-CAF	4.22	1.39	1.33
3	B	302	OCC	CAA-CAB	-3.88	1.40	1.46
2	B	301	OGA	C2-C1	-3.15	1.50	1.54
3	B	302	OCC	CAT-CAS	2.46	1.55	1.50
2	B	301	OGA	O2-C1	-2.43	1.23	1.30
3	B	302	OCC	CAX-CAT	-2.30	1.52	1.55

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	OCC	OBC-CBB-CAR	-7.87	121.92	130.45
3	B	302	OCC	OBA-CAZ-CAO	-6.18	101.59	106.56
3	B	302	OCC	OBA-CBB-CAR	5.35	113.23	107.88
3	B	302	OCC	CAZ-OBA-CBB	-4.57	105.08	110.33
3	B	302	OCC	OBA-CBB-OBC	2.99	124.84	121.07
2	B	301	OGA	O1-C1-C2	-2.65	116.27	122.18
3	B	302	OCC	CAG-CAD-CAC	-2.51	110.54	113.61
3	B	302	OCC	CAC-CAB-CAA	2.44	121.38	118.85

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	OGA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/296 (91%)	0.13	1 (0%) 92 93	7, 16, 27, 38	0
1	B	265/296 (89%)	0.11	4 (1%) 73 76	6, 15, 29, 41	0
All	All	535/592 (90%)	0.12	5 (0%) 84 86	6, 15, 28, 41	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	261	HIS	2.8
1	B	262	ASN	2.5
1	B	177	GLY	2.4
1	A	115	ILE	2.4
1	B	263	ASN	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	OCC	B	302	29/29	0.89	0.15	10,14,21,25	0
4	FE2	A	301	1/1	0.89	0.07	48,48,48,48	0
2	OGA	B	301	10/10	0.93	0.14	11,16,19,20	10
4	FE2	B	303	1/1	0.97	0.05	32,32,32,32	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FE2 A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers [i](#)

There are no such residues in this entry.