



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 02:00 pm GMT

PDB ID : 2EZ9  
Title : Pyruvate oxidase variant F479W in complex with reaction intermediate analogue 2-phosphonolactyl-thiamin diphosphate  
Authors : Wille, G.; Meyer, D.; Steinmetz, A.; Hinze, E.; Golbik, R.; Tittmann, K.  
Deposited on : 2005-11-10  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	2957 (1.60-1.60)
Clashscore	122126	3202 (1.60-1.60)
Ramachandran outliers	120053	3117 (1.60-1.60)
Sidechain outliers	120020	3116 (1.60-1.60)
RSRZ outliers	108989	2883 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	603	 % 94%
1	B	603	 % 93%

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 10254 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	585	4523	2868	780	861	14	0	0	0
1	B	585	4523	2868	780	861	14	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	479	TRP	PHE	ENGINEERED	UNP P37063
A	561	MET	THR	see remark 999	UNP P37063
B	479	TRP	PHE	ENGINEERED	UNP P37063
B	561	MET	THR	see remark 999	UNP P37063

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Mg	0	0
			1	1		
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Na	0	0
			1	1		
3	A	1	Total	Na	0	0
			1	1		


- Molecule 4 is 3-[(4-AMINO-2-METHYLPYRIMIDIN-5-YL)METHYL]-2-{(1S)-1-HYDROXY-1-[(R)-HYDROXY(METHOXY)PHOSPHORYL]ETHYL}-5-(2-[(S)-HYDROXY(PHOSPHONOXY)PHOSPHORYL]OXY)ETHYL)-4-METHYL-1,3-THIAZOL-3-IUM

(three-letter code: TDK) (formula:  $C_{15}H_{26}N_4O_{11}P_3S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
			Total	C	N	O	P			S	
4	A	1	Total	34	15	4	11	3	1	0	0
4	B	1	Total	34	15	4	11	3	1	0	0

- Molecule 5 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula:  $C_{27}H_{33}N_9O_{15}P_2$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
5	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	510	Total	O	0	0
			510	510		
6	B	520	Total	O	0	0
			520	520		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyruvate oxidase



- Molecule 1: Pyruvate oxidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.66Å 154.52Å 165.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.62 – 1.60 19.62 – 1.60	Depositor EDS
% Data completeness (in resolution range)	82.9 (19.62-1.60) 83.0 (19.62-1.60)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.98 (at 1.60Å)	Xtrriage
Refinement program	REFMAC refmac_5.2.0005	Depositor
R, $R_{free}$	0.158 , 0.180 0.159 , 0.180	Depositor DCC
$R_{free}$ test set	3328 reflections (2.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	11.8	Xtrriage
Anisotropy	0.024	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 57.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	10254	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TDK, NA, MG, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.44	0/4619	0.58	0/6291
1	B	0.45	0/4619	0.58	0/6291
All	All	0.44	0/9238	0.58	0/12582

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4523	0	4468	15	0
1	B	4523	0	4468	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	34	0	22	3	0
4	B	34	0	22	3	0
5	A	53	0	31	0	0
5	B	53	0	31	0	0
6	A	510	0	0	7	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	520	0	0	6	0
All	All	10254	0	9042	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:318:THR:HG23	6:A:2105:HOH:O	1.72	0.87
1:B:328:LYS:HD3	6:B:2123:HOH:O	1.85	0.76
1:B:232:GLN:HG3	6:B:2119:HOH:O	1.97	0.63
1:A:318:THR:HG21	1:A:321:ALA:HB2	1.81	0.62
1:A:224:ARG:HG2	1:A:250:ILE:HD11	1.84	0.60
4:B:1613:TDK:OM1	4:B:1613:TDK:N4'	2.33	0.60
1:B:57:ARG:HD2	6:B:1858:HOH:O	2.03	0.58
1:A:316:HIS:HE1	6:A:2050:HOH:O	1.86	0.57
1:A:422:MET:SD	4:A:1611:TDK:HM43	2.46	0.56
1:B:422:MET:SD	4:B:1613:TDK:HM43	2.46	0.56
4:A:1611:TDK:OM1	4:A:1611:TDK:N4'	2.38	0.55
1:B:316:HIS:HD2	6:B:1723:HOH:O	1.90	0.53
1:B:568:GLU:HG2	6:B:2098:HOH:O	2.09	0.52
1:A:318:THR:HG22	1:B:188:SER:OG	2.11	0.51
1:A:50:ARG:HH11	1:A:50:ARG:HB3	1.77	0.50
4:B:1613:TDK:H4'1	4:B:1613:TDK:CMA	2.25	0.50
1:B:232:GLN:HG2	1:B:334:LEU:HD11	1.95	0.49
1:A:232:GLN:HG3	6:A:2118:HOH:O	2.13	0.48
1:A:57:ARG:HD2	6:A:2108:HOH:O	2.13	0.48
6:A:1773:HOH:O	1:B:148:HIS:HE1	1.96	0.48
1:A:318:THR:CG2	1:B:188:SER:OG	2.62	0.47
1:A:568:GLU:HG2	6:A:1990:HOH:O	2.14	0.47
1:B:406:LYS:HE3	6:B:1790:HOH:O	2.14	0.46
4:A:1611:TDK:CMA	4:A:1611:TDK:H4'1	2.28	0.45
1:A:515:ARG:HD2	1:A:541:ASP:OD2	2.17	0.45
1:A:39:SER:HB3	1:A:174:TRP:CD2	2.52	0.45
1:A:316:HIS:HD2	6:A:1673:HOH:O	2.02	0.43
1:B:399:LEU:HD23	1:B:399:LEU:C	2.39	0.43
1:B:354:LYS:HB3	1:B:354:LYS:HE2	1.92	0.41
1:A:214:ARG:NH1	1:A:276:GLN:HE21	2.18	0.41
1:B:377:LEU:CD1	1:B:400:ASN:HB3	2.51	0.41
1:B:263:ASN:HB2	1:B:287:TYR:OH	2.20	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:387:ASP:O	1:B:439:ARG:HD2	2.20	0.41
1:B:228:LYS:HB2	1:B:228:LYS:HE3	1.79	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	583/603 (97%)	571 (98%)	11 (2%)	1 (0%)	49 27
1	B	583/603 (97%)	572 (98%)	10 (2%)	1 (0%)	49 27
All	All	1166/1206 (97%)	1143 (98%)	21 (2%)	2 (0%)	49 27

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	263	ASN
1	A	263	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	470/485 (97%)	467 (99%)	3 (1%)	87 79
1	B	470/485 (97%)	467 (99%)	3 (1%)	87 79

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	940/970 (97%)	934 (99%)	6 (1%)	87	79

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	438	GLU
1	A	515	ARG
1	A	592	ASP
1	B	193	LEU
1	B	515	ARG
1	B	524	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	148	HIS
1	A	232	GLN
1	A	263	ASN
1	A	276	GLN
1	A	316	HIS
1	A	464	HIS
1	A	476	GLN
1	B	10	ASN
1	B	52	HIS
1	B	148	HIS
1	B	263	ASN
1	B	276	GLN
1	B	316	HIS
1	B	464	HIS
1	B	476	GLN
1	B	489	GLN
1	B	572	GLN
1	B	588	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	TDK	A	1611	2	26,35,35	1.29	4 (15%)	32,55,55	2.05	9 (28%)
5	FAD	A	1612	-	51,58,58	1.00	5 (9%)	57,89,89	1.88	8 (14%)
4	TDK	B	1613	2	26,35,35	1.29	3 (11%)	32,55,55	2.17	10 (31%)
5	FAD	B	1614	-	51,58,58	1.00	3 (5%)	57,89,89	2.05	9 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TDK	A	1611	2	-	0/26/35/35	0/2/2/2
5	FAD	A	1612	-	-	0/28/50/50	0/6/6/6
4	TDK	B	1613	2	-	0/26/35/35	0/2/2/2
5	FAD	B	1614	-	-	0/28/50/50	0/6/6/6

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1614	FAD	C1'-N10	-2.39	1.45	1.48
5	A	1612	FAD	C8A-N9A	-2.25	1.34	1.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1611	TDK	PC-OM3	-2.23	1.51	1.56
5	A	1612	FAD	C1'-N10	-2.03	1.46	1.48
4	A	1611	TDK	C4'-N3'	2.05	1.38	1.35
5	A	1612	FAD	C4-N3	2.08	1.36	1.33
4	A	1611	TDK	C7'-C5'	2.17	1.55	1.51
4	B	1613	TDK	C7'-C5'	2.23	1.55	1.51
5	A	1612	FAD	C4X-N5	2.26	1.36	1.33
5	A	1612	FAD	C10-N1	2.62	1.36	1.33
4	B	1613	TDK	PB-O3A	2.63	1.64	1.60
5	B	1614	FAD	C4X-N5	2.82	1.37	1.33
5	B	1614	FAD	C10-N1	2.91	1.37	1.33
4	B	1613	TDK	PC-OM2	3.55	1.62	1.57
4	A	1611	TDK	PC-OM2	3.71	1.63	1.57

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1614	FAD	N3A-C2A-N1A	-9.00	121.16	128.86
5	A	1612	FAD	N3A-C2A-N1A	-8.31	121.75	128.86
4	B	1613	TDK	OM2-PC-OM4	-4.00	105.56	114.66
4	A	1611	TDK	CM4-C4-C5	-3.52	119.89	127.60
4	B	1613	TDK	CM4-C4-C5	-3.49	119.97	127.60
4	B	1613	TDK	C5'-C7'-N3	-3.42	107.63	113.31
5	B	1614	FAD	C4X-C4-N3	-3.42	118.60	123.47
5	A	1612	FAD	C4X-C4-N3	-3.17	118.96	123.47
4	A	1611	TDK	OM2-PC-OM4	-3.14	107.53	114.66
4	B	1613	TDK	N1'-C2'-N3'	-3.13	120.03	125.55
4	A	1611	TDK	C5'-C7'-N3	-2.94	108.43	113.31
5	B	1614	FAD	C9A-N10-C10	-2.92	117.88	121.77
4	A	1611	TDK	N1'-C2'-N3'	-2.72	120.76	125.55
5	B	1614	FAD	C1B-N9A-C4A	-2.70	121.98	126.64
5	A	1612	FAD	C4A-C5A-N7A	-2.40	107.09	109.41
5	A	1612	FAD	C9A-C5X-N5	-2.39	118.92	122.32
5	A	1612	FAD	C9A-N10-C10	-2.32	118.68	121.77
5	B	1614	FAD	C4A-C5A-N7A	-2.08	107.40	109.41
5	B	1614	FAD	C1'-C2'-C3'	2.01	115.58	109.82
4	B	1613	TDK	CM4-C4-N3	2.11	125.34	122.69
4	A	1611	TDK	CM2-C2'-N1'	2.23	119.55	117.07
5	A	1612	FAD	C1'-C2'-C3'	2.29	116.36	109.82
4	B	1613	TDK	CM2-C2'-N3'	2.42	121.25	117.20
4	A	1611	TDK	OM3-PC-CMA	2.69	110.93	106.27
4	A	1611	TDK	C6'-N1'-C2'	2.79	120.81	115.95

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1614	FAD	C4-C4X-N5	2.92	121.98	118.70
4	B	1613	TDK	OM3-PC-CMA	3.10	111.64	106.27
5	A	1612	FAD	C4-C4X-N5	3.16	122.26	118.70
4	B	1613	TDK	C6'-N1'-C2'	3.35	121.78	115.95
4	B	1613	TDK	C5-C4-N3	3.35	114.67	107.66
4	A	1611	TDK	C5-C4-N3	3.59	115.17	107.66
5	B	1614	FAD	C1'-N10-C9A	4.16	122.01	118.31
4	B	1613	TDK	C6-C5-C4	5.89	132.16	127.43
4	A	1611	TDK	C6-C5-C4	6.35	132.53	127.43
5	A	1612	FAD	C4-N3-C2	7.12	121.20	115.14
5	B	1614	FAD	C4-N3-C2	7.66	121.66	115.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1611	TDK	3	0
4	B	1613	TDK	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	585/603 (97%)	-0.59	6 (1%) 82 82	7, 11, 20, 35	0
1	B	585/603 (97%)	-0.63	4 (0%) 87 88	7, 10, 19, 34	0
All	All	1170/1206 (97%)	-0.61	10 (0%) 84 84	7, 11, 20, 35	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	560	ALA	4.3
1	B	9	THR	4.2
1	A	592	ASP	4.2
1	B	592	ASP	3.2
1	A	9	THR	3.0
1	A	561	MET	2.8
1	A	593	ASP	2.5
1	B	180	GLU	2.4
1	A	180	GLU	2.4
1	B	368	GLU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	TDK	A	1611	34/34	0.97	0.07	8,12,21,21	0
4	TDK	B	1613	34/34	0.97	0.07	7,10,20,21	0
5	FAD	A	1612	53/53	0.98	0.05	6,8,10,10	0
5	FAD	B	1614	53/53	0.99	0.04	6,7,9,9	0
2	MG	B	1512	1/1	1.00	0.04	9,9,9,9	0
3	NA	A	1513	1/1	1.00	0.09	6,6,6,6	1
2	MG	A	1510	1/1	1.00	0.04	9,9,9,9	0
3	NA	B	1511	1/1	1.00	0.09	6,6,6,6	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.