

Full wwPDB X-ray Structure Validation Report (i)

Jun 22, 2024 – 01:22 PM EDT

:	6EZ1
:	Structure of h. salinarum RosR (vng0258) grown from NaBr
:	Shaanan, B.; Kutnowski, N.
	2017-11-13
:	1.75 Å(reported)
	::

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	127	83%	•	13%
1	В	127	80%	6%	13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BR	А	217	-	-	Х	-



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3792 atoms, of which 1708 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	1	
1	٨	110	Total	С	Н	Ν	Ο	0	1	
1	А	110	1740	553	853	150	184	0	1	
1	В	110	Total	С	Н	Ν	Ο	0	1	
1	D	110	1745	554	855	151	185	0	1	

• Molecule 1 is a protein called DNA binding protein.

Chain	Residue	Modelled	Actual	Comment	Reference
А	117	ALA	-	expression tag	UNP Q9HSF4
А	118	ALA	-	expression tag	UNP Q9HSF4
А	119	ALA	-	expression tag	UNP Q9HSF4
А	120	LEU	-	expression tag	UNP Q9HSF4
А	121	GLU	-	expression tag	UNP Q9HSF4
A	122	HIS	-	expression tag	UNP Q9HSF4
A	123	HIS	-	expression tag	UNP Q9HSF4
A	124	HIS	-	expression tag	UNP Q9HSF4
А	125	HIS	-	expression tag	UNP Q9HSF4
А	126	HIS	-	expression tag	UNP Q9HSF4
A	127	HIS	-	expression tag	UNP Q9HSF4
В	117	ALA	-	expression tag	UNP Q9HSF4
В	118	ALA	-	expression tag	UNP Q9HSF4
В	119	ALA	-	expression tag	UNP Q9HSF4
В	120	LEU	-	expression tag	UNP Q9HSF4
В	121	GLU	-	expression tag	UNP Q9HSF4
В	122	HIS	-	expression tag	UNP Q9HSF4
В	123	HIS	-	expression tag	UNP Q9HSF4
В	124	HIS	-	expression tag	UNP Q9HSF4
В	125	HIS	-	expression tag	UNP Q9HSF4
В	126	HIS	-	expression tag	UNP Q9HSF4
В	127	HIS	-	expression tag	UNP Q9HSF4

There are 22 discrepancies between the modelled and reference sequences:

• Molecule 2 is BROMIDE ION (three-letter code: BR) (formula: Br) (labeled as "Ligand of Interest" by depositor).


Trace

0

0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	20	TotalBr2121	0	1
2	В	13	Total Br 14 14	0	1



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	А	1	Total 5	0 4	S 1	0	0


• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	127	Total O 127 127	0	0
4	В	140	Total O 140 140	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: DNA binding protein



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	40.88Å 71.14Å 41.72Å	Depositor
a, b, c, α , β , γ	90.00° 110.33° 90.00°	Depositor
Resolution (Å)	34.28 - 1.75	Depositor
Tresolution (A)	34.28 - 1.75	EDS
% Data completeness	96.9 (34.28-1.75)	Depositor
(in resolution range)	96.9 (34.28-1.75)	EDS
R _{merge}	0.11	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.00 (at 1.75 \text{\AA})$	Xtriage
Refinement program	REFMAC refmac5 version 5.8.0158, PHENIX dev-2947	Depositor
R, R_{free}	0.170 , 0.209	Depositor
It, It _{free}	0.180 , 0.217	DCC
R_{free} test set	1111 reflections (5.07%)	wwPDB-VP
Wilson B-factor $(Å^2)$	17.5	Xtriage
Anisotropy	0.366	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.43 , 53.4	EDS
L-test for $twinning^2$	$< L > = 0.48, < L^2 > = 0.30$	Xtriage
Estimated twinning fraction	0.041 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3792	wwPDB-VP
Average B, all atoms $(Å^2)$	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.11% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BR, $\mathrm{SO4}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	А	0.26	0/901	0.44	0/1219	
1	В	0.27	0/901	0.44	0/1219	
All	All	0.27	0/1802	0.44	0/2438	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	887	853	855	4	0
1	В	890	855	854	8	0
2	А	21	0	0	5	0
2	В	14	0	0	6	0
3	А	5	0	0	0	0
4	А	127	0	0	2	1
4	В	140	0	0	2	1
All	All	2084	1708	1709	16	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:106:ARG:NH1	2:B:210:BR:BR	2.54	0.94
2:B:211:BR:BR	2:B:212:BR:BR	2.96	0.94
2:A:201:BR:BR	4:A:383:HOH:O	2.53	0.80
1:B:76:ASN:HB2	2:B:205[B]:BR:BR	2.49	0.68
1:B:32:TYR:HE1	1:B:71:LEU:HD12	1.64	0.62
2:A:205:BR:BR	2:A:217:BR:BR	3.27	0.62
1:B:75:THR:HB	2:B:205[A]:BR:BR	2.58	0.58
1:B:32:TYR:CE1	1:B:71:LEU:HD12	2.40	0.56
2:A:217:BR:BR	2:B:203:BR:BR	3.35	0.55
2:A:213:BR:BR	2:B:207:BR:BR	3.37	0.53
1:A:95:THR:HG23	4:B:322:HOH:O	2.13	0.48
4:A:322:HOH:O	1:B:95:THR:HG23	2.14	0.47
1:A:72:ASP:OD2	2:A:220:BR:BR	2.88	0.46
1:A:20:LYS:NZ	1:A:91[B]:ASP:OD2	2.47	0.44
1:B:74:ARG:NH1	4:B:312:HOH:O	2.51	0.43
1:A:43:TYR:OH	1:B:20:LYS:NZ	2.53	0.41

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:310:HOH:O	4:B:313:HOH:O[2_555]	2.19	0.01

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	109/127~(86%)	105~(96%)	4 (4%)	0	100	100
1	В	109/127~(86%)	104 (95%)	5 (5%)	0	100	100

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles	
All	All	218/254~(86%)	209~(96%)	9~(4%)	0	100 100	

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	93/107~(87%)	93 (100%)	0	100 100
1	В	93/107~(87%)	93 (100%)	0	100 100
All	All	186/214~(87%)	186 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 36 ligands modelled in this entry, 35 are monoatomic - leaving 1 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Type	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Chain	Res	Res Link	Bond lengths			Bond angles		
	туре		i nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2							
3	SO4	А	221	2	4,4,4	0.26	0	$6,\!6,\!6$	0.08	0							

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< RSRZ >	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	110/127~(86%)	0.08	4 (3%) 42 49	10, 17, 46, 84	0
1	В	110/127~(86%)	0.25	9 (8%) 11 15	8, 16, 66, 131	0
All	All	220/254~(86%)	0.16	13 (5%) 22 27	8, 17, 51, 131	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	73	LYS	9.3
1	В	74	ARG	8.1
1	В	72	ASP	7.7
1	А	74	ARG	5.6
1	В	71	LEU	5.6
1	В	70	GLU	4.9
1	В	75	THR	4.1
1	А	75	THR	3.6
1	А	71	LEU	3.3
1	В	76	ASN	3.0
1	В	11	ASP	2.9
1	А	73	LYS	2.3
1	В	69	SER	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BR	А	212	1/1	0.68	0.09	45,45,45,45	1
2	BR	А	214	1/1	0.82	0.10	47,47,47,47	1
2	BR	А	216	1/1	0.85	0.07	51,51,51,51	1
2	BR	А	219	1/1	0.86	0.07	38,38,38,38	1
3	SO4	А	221	5/5	0.86	0.17	16,24,40,41	5
2	BR	В	204	1/1	0.87	0.09	34,34,34,34	1
2	BR	В	210	1/1	0.87	0.07	$50,\!50,\!50,\!50$	1
2	BR	А	210	1/1	0.87	0.12	41,41,41,41	1
2	BR	В	201	1/1	0.90	0.06	27,27,27,27	1
2	BR	А	203	1/1	0.91	0.06	36,36,36,36	0
2	BR	А	220	1/1	0.91	0.09	105,105,105,105	0
2	BR	В	207	1/1	0.92	0.08	28,28,28,28	1
2	BR	А	209	1/1	0.93	0.06	38,38,38,38	1
2	BR	В	208	1/1	0.93	0.07	37,37,37,37	1
2	BR	В	209	1/1	0.93	0.11	22,22,22,22	1
2	BR	А	206	1/1	0.93	0.05	26,26,26,26	1
2	BR	В	211	1/1	0.93	0.08	35,35,35,35	1
2	BR	А	208	1/1	0.93	0.07	36,36,36,36	1
2	BR	В	213	1/1	0.95	0.09	26,26,26,26	1
2	BR	В	206	1/1	0.95	0.10	27,27,27,27	1
2	BR	В	202	1/1	0.96	0.07	24,24,24,24	1
2	BR	А	202	1/1	0.96	0.08	19,19,19,19	1
2	BR	А	211	1/1	0.96	0.09	33,33,33,33	1
2	BR	А	204[A]	1/1	0.96	0.08	27,27,27,27	1
2	BR	А	204[B]	1/1	0.96	0.08	47,47,47,47	1
2	BR	А	201	1/1	0.97	0.05	35,35,35,35	0
2	BR	А	218	1/1	0.97	0.15	29,29,29,29	1
2	BR	В	203	1/1	0.97	0.04	23,23,23,23	1
2	BR	А	213	1/1	0.97	0.14	21,21,21,21	1
2	BR	В	205[A]	1/1	0.97	0.07	25,25,25,25	1
2	BR	В	205[B]	1/1	0.97	0.07	50,50,50,50	1
2	BR	А	207	1/1	0.97	0.06	30,30,30,30	1
2	BR	В	212	1/1	0.98	0.07	37,37,37,37	1
2	BR	А	205	1/1	0.98	0.05	24,24,24,24	1
2	BR	А	215	1/1	0.98	0.18	24,24,24,24	1
2	BR	А	217	1/1	0.99	0.07	22,22,22,22	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.











































































































































































6.5 Other polymers (i)

There are no such residues in this entry.

