

# Full wwPDB X-ray Structure Validation Report (i)

#### Jun 24, 2024 – 08:18 PM EDT

PDB ID	:	6F3G
Title	:	IRAK4 IN COMPLEX WITH inhibitor
Authors	:	Xue, Y.; Degorce, S.L.; Robb, G.R.; Ferguson, A.D.
Deposited on	:	2017-11-28
Resolution	:	2.37  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R <sub>free</sub>	130704	5509(2.40-2.36)
Clashscore	141614	6082 (2.40-2.36)
Ramachandran outliers	138981	5973(2.40-2.36)
Sidechain outliers	138945	5975(2.40-2.36)
RSRZ outliers	127900	5397(2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
			30%		
1	А	295	83%	11%	6%
			27%		
2	В	295	83%	11%	5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	SEP	А	346	-	-	-	Х
2	TPO	В	345	-	-	-	Х
2	SEP	В	346	-	-	-	Х



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 4544 atoms, of which 4 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Interleukin-1 receptor-associated kinase 4.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace			
1	А	277	Total 2209	C 1385	N 371	O 436	Р 3	S 14	0	0	0

• Molecule 2 is a protein called Interleukin-1 receptor-associated kinase 4.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
2	В	279	Total 2211	C 1390	N 373	0 432	Р 2	S 14	0	0	0

• Molecule 3 is {N}4, {N}4-dimethyl- {N}1-(5-propan-2-ylpyrrolo[3,2-d]pyrimidin-4-yl)cyclo hexane-1,4-diamine (three-letter code: CJN) (formula:  $C_{17}H_{27}N_5$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	А	1	Total 24	C 17	Н 2	N 5	0	0
3	В	1	Total 24	C 17	Н 2	N 5	0	0



• Molecule 4 is SULFATE ION (three-letter code: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	В	1	$\begin{array}{ccc} \text{Total} & \text{O} & \text{S} \\ 5 & 4 & \text{I} \end{array}$	5 1	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	37	$\begin{array}{cc} \text{Total} & \text{O} \\ 37 & 37 \end{array}$	0	0
5	В	34	$\begin{array}{cc} \text{Total} & \text{O} \\ 34 & 34 \end{array}$	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Interleukin-1 receptor-associated kinase 4



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants	87.51Å 110.28Å 142.13Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Bosolution (Å)	29.13 - 2.37	Depositor
	29.04 - 2.37	EDS
% Data completeness	99.9 (29.13-2.37)	Depositor
(in resolution range)	99.9(29.04-2.37)	EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.30 (at 2.36 \text{\AA})$	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
B B.	0.290 , $0.314$	Depositor
$n, n_{free}$	0.286 , $0.308$	DCC
$R_{free}$ test set	1384 reflections $(4.90\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	49.9	Xtriage
Anisotropy	0.535	Xtriage
Bulk solvent $k_{sol}(e/A^3), B_{sol}(A^2)$	0.32 , $44.7$	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.47, < L^2>=0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4544	wwPDB-VP
Average B, all atoms $(Å^2)$	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 20.98 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.7309e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, TPO, SEP, CJN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.45	0/2209	0.64	0/2972	
2	В	0.48	0/2224	0.66	0/2991	
All	All	0.46	0/4433	0.65	0/5963	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2209	0	2180	13	0
2	В	2211	0	2182	19	4
3	А	22	2	0	0	0
3	В	22	2	0	0	0
4	В	5	0	0	0	0
5	А	37	0	0	0	0
5	В	34	0	0	1	0
All	All	4540	4	4362	31	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.



A + a 1	A 4 a ma 2	Interatomic	Clash
Atom-1	Atom-2	distance $(\text{\AA})$	overlap (Å)
2:B:178:ASN:C	2:B:179:ASN:N	2.17	0.99
1:A:169:SER:HB3	1:A:172:GLU:HG3	1.59	0.83
1:A:185:ILE:HD11	1:A:192:MET:HG2	1.65	0.78
2:B:210:VAL:CG1	2:B:261:VAL:HG13	2.20	0.71
2:B:192:MET:SD	2:B:264:TYR:HE1	2.19	0.66
2:B:210:VAL:HG11	2:B:261:VAL:HG13	1.80	0.62
1:A:237:MET:SD	1:A:330:PHE:CD2	2.98	0.57
1:A:237:MET:SD	1:A:330:PHE:HD2	2.28	0.56
1:A:390:HIS:O	2:B:390:HIS:O	2.24	0.56
2:B:210:VAL:HG13	2:B:261:VAL:HG13	1.86	0.56
1:A:409:THR:HG23	1:A:412:ASP:H	1.70	0.56
2:B:210:VAL:HG11	2:B:261:VAL:CG1	2.36	0.56
2:B:266:PRO:CG	2:B:321:GLU:HG3	2.37	0.54
1:A:222:THR:HG22	1:A:225:GLU:HG3	1.89	0.53
1:A:246:VAL:HG11	1:A:318:LEU:HD12	1.91	0.53
1:A:233:GLU:O	1:A:237:MET:HG2	2.10	0.52
2:B:174:LYS:HE2	2:B:179:ASN:OD1	2.10	0.51
2:B:266:PRO:HG3	2:B:321:GLU:HG3	1.93	0.50
2:B:210:VAL:CG1	2:B:261:VAL:CG1	2.92	0.47
1:A:345:TPO:HB	1:A:364:ILE:HD11	1.99	0.45
2:B:210:VAL:HG22	2:B:249:LEU:HD12	1.98	0.44
2:B:265:MET:HE1	5:B:624:HOH:O	2.17	0.44
2:B:237:MET:SD	2:B:248:LEU:HB2	2.59	0.43
2:B:246:VAL:HG21	2:B:318:LEU:HD12	2.00	0.42
2:B:192:MET:SD	2:B:264:TYR:CE1	3.08	0.42
1:A:222:THR:HG23	1:A:225:GLU:H	1.85	0.41
2:B:345:TPO:HB	2:B:364:ILE:HD11	2.00	0.41
2:B:414:ILE:HG12	2:B:426:VAL:HG11	2.01	0.41
2:B:311:ASP:HB2	2:B:332:LEU:HD12	2.02	0.41
1:A:168:PHE:HE2	1:A:250:GLY:HA3	1.86	0.41
1:A:414:ILE:HG12	1:A:426:VAL:HG11	2.02	0.40

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:172:GLU:OE1	2:B:172:GLU:OE1[4_545]	0.41	1.79
2:B:172:GLU:CD	2:B:172:GLU:OE1[4_545]	1.55	0.65
2:B:172:GLU:CG	2:B:172:GLU:CG[4_545]	1.81	0.39



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:172:GLU:CD	2:B:172:GLU:CD[4_545]	2.19	0.01

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	263/295~(89%)	256~(97%)	7 (3%)	0	100 100
2	В	267/295~(90%)	258~(97%)	8~(3%)	1 (0%)	34 46
All	All	530/590~(90%)	514 (97%)	15 (3%)	1 (0%)	47 61

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	В	181	ASP

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric Outlie		Percentiles	
1	А	243/254~(96%)	233~(96%)	10 (4%)	30 45	
2	В	243/255~(95%)	235~(97%)	8 (3%)	38 55	
All	All	486/509~(96%)	468 (96%)	18 (4%)	34 50	

All (18) residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	А	164	ARG
1	А	194	GLU
1	А	206	ASN
1	А	290	LYS
1	А	328	SER
1	А	344	MET
1	А	352	THR
1	А	389	GLU
1	А	401	GLU
1	А	408	LYS
2	В	169	SER
2	В	185	ILE
2	В	206	ASN
2	В	214	LYS
2	В	290	LYS
2	В	389	GLU
2	В	401	GLU
2	В	408	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	206	ASN
2	В	452	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Turne	Chain	Chain	Chain	Chain	Chain	Chain	Dec	Ros Link	B	Bond lengths			Bond angles		
	Moi Type		I nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2						
1	TPO	А	342	1	8,10,11	1.04	0	10,14,16	1.02	1 (10%)						
2	TPO	В	345	2	8,10,11	0.99	0	10,14,16	1.23	1 (10%)						
1	SEP	А	346	1	8,9,10	0.84	0	8,12,14	4.63	3 (37%)						
2	SEP	В	346	2	8,9,10	0.77	0	8,12,14	2.30	4 (50%)						
1	TPO	А	345	1	8,10,11	0.93	0	10,14,16	2.04	3 (30%)						

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	А	342	1	-	1/9/11/13	-
2	TPO	В	345	2	-	4/9/11/13	-
1	SEP	А	346	1	-	2/5/8/10	-
2	SEP	В	346	2	-	0/5/8/10	-
1	TPO	А	345	1	-	3/9/11/13	-

There are no bond length outliers.

All (	12)	bond	angle	outliers	are	listed	below:
-------	-----	------	-------	----------	-----	--------	--------

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
1	А	346	SEP	OG-CB-CA	12.31	120.12	108.14
2	В	346	SEP	OG-CB-CA	4.86	112.88	108.14
1	А	345	TPO	O3P-P-OG1	4.18	124.74	105.99
1	А	346	SEP	OG-P-O1P	3.24	115.57	106.47
2	В	346	SEP	O3P-P-OG	2.87	114.36	106.73
2	В	345	TPO	O3P-P-OG1	2.50	117.20	105.99
1	А	345	TPO	O3P-P-O2P	-2.44	98.33	107.64
1	А	345	TPO	O3P-P-O1P	-2.35	101.49	110.68
1	А	342	TPO	CG2-CB-CA	-2.20	108.83	113.16
2	В	346	SEP	OG-P-O1P	2.17	112.56	106.47
2	В	346	SEP	P-OG-CB	-2.08	112.58	118.30
1	А	346	SEP	P-OG-CB	-2.05	112.64	118.30

There are no chirality outliers.

All (10) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
1	А	345	TPO	N-CA-CB-OG1
1	А	345	TPO	O-C-CA-CB
1	А	346	SEP	N-CA-CB-OG
2	В	345	TPO	N-CA-CB-OG1
2	В	345	TPO	O-C-CA-CB
1	А	346	SEP	CB-OG-P-O2P
1	А	342	TPO	CA-CB-OG1-P
2	В	345	TPO	CB-OG1-P-O1P
1	А	345	TPO	CB-OG1-P-O2P
2	В	345	TPO	CB-OG1-P-O2P

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	345	TPO	1	0
1	А	345	TPO	1	0

#### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry (i)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Turne	Chain	Dec	Tinle	Bo	ond leng	$\mathbf{ths}$	B	ond ang	les
IVIOI	туре	Chain	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	SO4	В	501	-	4,4,4	0.14	0	6,6,6	0.20	0
3	CJN	В	502	-	19,24,24	0.87	0	22,34,34	2.80	9 (40%)
3	CJN	А	501	-	19,24,24	0.89	0	22,34,34	2.95	7 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



- mea	115 110 Ou		nat Kn	iu were	identified.		
Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CJN	В	502	-	-	2/12/22/22	0/3/3/3
3	CJN	А	501	_	-	0/12/22/22	0/3/3/3

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. ', means no outliers of that kind were identified

| 0/3/3/3 |

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
3	В	502	CJN	C9-N10-C11	8.70	124.05	116.59
3	А	501	CJN	C9-N10-C11	8.49	123.87	116.59
3	А	501	CJN	N8-C9-N10	-6.24	118.93	128.68
3	В	502	CJN	N8-C9-N10	-5.35	120.31	128.68
3	А	501	CJN	C9-N8-C7	5.24	122.63	115.40
3	В	502	CJN	C22-N21-C17	4.50	120.13	112.39
3	А	501	CJN	C5-N4-C2	-3.81	121.63	125.40
3	В	502	CJN	C9-N8-C7	3.67	120.45	115.40
3	А	501	CJN	C22-N21-C17	3.62	118.61	112.39
3	В	502	CJN	C16-C15-C13	-3.09	108.07	111.48
3	А	501	CJN	C16-C15-C13	-2.98	108.19	111.48
3	А	501	CJN	C3-C2-C1	-2.11	106.78	113.28
3	В	502	CJN	C11-N12-C13	-2.10	120.18	124.16
3	В	502	CJN	C3-C2-C1	-2.04	107.00	113.28
3	В	502	CJN	C23-N21-C17	2.03	115.88	112.39
3	В	502	CJN	C1-C2-N4	2.01	113.59	110.30

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	502	CJN	C16-C17-N21-C23
3	В	502	CJN	C19-C17-N21-C23

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	В	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	В	178:ASN	С	179:ASN	N	2.17



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q < 0.9
1	А	274/295~(92%)	1.87	89 (32%) 0 0	32, 64, 161, 194	0
2	В	277/295~(93%)	1.53	79 (28%) 0 0	29, 62, 109, 145	0
All	All	551/590~(93%)	1.70	168 (30%) 0 0	29, 63, 139, 194	0

All (168) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	165	PHE	13.2
1	А	171	TYR	11.6
1	А	199	VAL	11.6
1	А	216	ALA	10.1
1	А	187	VAL	10.0
1	А	205	VAL	9.3
1	А	198	GLY	7.9
2	В	196	GLY	7.9
2	В	258	LEU	7.3
1	А	239	LYS	7.2
1	А	222	THR	7.2
1	А	179	ASN	7.1
2	В	347	ARG	7.1
1	А	170	PHE	6.9
1	А	258	LEU	6.8
2	В	199	VAL	6.5
1	А	185	ILE	6.5
1	А	184	PRO	6.5
1	А	347	ARG	6.4
2	В	349	VAL	6.2
2	В	195	GLY	6.1
1	А	251	PHE	6.1
2	В	348	ILE	5.9
1	А	349	VAL	5.8



6F3G
------

Continued from previous page							
Mol	Chain	Res	Type	RSRZ			
1	А	183	ARG	5.8			
1	А	166	HIS	5.7			
2	В	226	LEU	5.6			
1	А	390	390 HIS				
1	А	241	241 GLN				
1	А	228	GLN	5.4			
2	В	201	TYR	5.4			
2	В	343	VAL	5.3			
1	А	227	LYS	5.2			
2	В	230	PHE	5.1			
2	В	185	ILE	5.1			
1	A	224	GLU	5.0			
1	А	206	ASN	5.0			
2	В	335	ALA	5.0			
2	В	187	VAL	4.9			
2	В	198	GLY	4.8			
2	В	189	GLY	4.8			
1	А	344	MET	4.8			
1	А	327	ILE	4.8			
1	А	226	LEU	4.7			
1	А	232	GLN	4.7			
1	А	215	LEU	4.7			
1	А	167	SER	4.6			
1	А	257	ASP	4.6			
1	А	348	ILE	4.6			
2	В	252	SER	4.6			
2	В	327	ILE	4.5			
1	А	254	ASP	4.5			
1	А	178	ASN	4.5			
2	В	227	LYS	4.4			
1	А	230	PHE	4.1			
1	А	175	ASN	4.1			
2	В	374	GLY	4.1			
1	А	190	ASN	4.0			
1	А	235	LYS	4.0			
1	А	253	SER	4.0			
1	А	377	LEU	3.9			
2	В	170	PHE	3.9			
2	В	223	THR	3.9			
2	В	375	VAL	3.9			
2	В	190	ASN	3.8			
2	В	257	ASP	3.8			



6F3G	
------	--

		- r · · · · ·	r ng r	
Mol	Chain	Res	Type	RSRZ
2	В	344	MET	3.8
2	В	376	VAL	3.7
1	А	201	TYR	3.7
1	А	207	ASN	3.7
1	А	238	ALA	3.7
1	А	407	GLU	3.6
2	В	172	GLU	3.6
2	В	373	PHE	3.6
2	В	336	SER	3.6
2	В	371	TYR	3.6
1	А	231	ASP	3.6
2	В	214	LYS	3.6
1	А	312	ILE	3.5
1	А	376	VAL	3.5
1	А	194	GLU	3.5
1	А	248	LEU	3.5
1	А	174	LYS	3.5
1	А	168	PHE	3.4
2	В	259	CYS	3.4
1	А	334 ARG		3.4
1	А	263	VAL	3.3
2	В	225	GLU	3.3
1	А	180	PHE	3.3
2	В	173	LEU	3.3
1	А	343	VAL	3.3
2	В	179	ASN	3.2
2	В	224	GLU	3.2
1	А	225	GLU	3.2
2	В	420	ASP	3.2
1	А	330	PHE	3.2
2	В	168	PHE	3.1
2	В	180	PHE	3.1
2	В	457	MET	3.1
1	А	259	CYS	3.1
2	В	204	TYR	3.1
1	А	212	VAL	3.0
1	А	260	LEU	3.0
2	В	231	ASP	3.0
2	В	286	HIS	2.9
1	А	237	MET	2.9
2	В	253	SER	2.8
2	В	176	VAL	2.8



6F3G
------

Mol	Chain	Res Type		RSRZ
2	В	194	GLU	2.8
1	А	375	VAL	2.8
2	В	235	LYS	2.8
1	А	453	LEU	2.8
2	В	171	TYR	2.8
2	В	312	ILE	2.8
2	В	228	GLN	2.7
2	В	377	LEU	2.7
2	В	317	ILE	2.7
1	А	450	VAL	2.7
1	А	208	THR	2.7
1	А	245	LEU	2.7
1	A	240	CYS	2.6
1	А	252	SER	2.6
2	В	191	LYS	2.6
2	В	229	GLN	2.6
1	А	176	VAL	2.5
2	В	455	GLN	2.5
1	А	182	GLU	2.5
1	А	204	TYR	2.5
2	В	390	HIS	2.5
2	В	372	SER	2.4
2	В	443	LYS	2.4
1	А	373	PHE	2.4
1	А	362	GLY	2.4
2	В	197	PHE	2.4
1	А	408	LYS	2.4
1	А	200	VAL	2.4
1	А	270	LEU	2.3
2	В	238	ALA	2.3
2	В	295	ALA	2.3
2	В	167	SER	2.3
1	A	439	GLU	2.3
1	A	389	GLU	2.2
1	А	455	GLN	2.2
1	A	262	TYR	2.2
2	В	169	SER	2.2
2	В	440	LYS	2.2
2	В	245	LEU	2.2
2	В	410	ILE	2.1
1	А	192	MET	2.1
2	В	203	GLY	2.1



Mol	Chain	Res Type		RSRZ	
2	В	438	HIS	2.1	
1	А	210	VAL	2.1	
1	А	310	ARG	2.1	
2	В	192	MET	2.1	
1	А	329	ASP	2.1	
2	В	251	PHE	2.1	
2	В	406	GLU	2.1	
2	В	270	LEU	2.1	
2	В	437	LEU	2.1	
2	В	314	SER	2.1	
2	В	240	CYS	2.0	
2	В	306	HIS	2.0	
2	В	378	LEU	2.0	
2	В	411	GLU	2.0	
1	А	374	GLY	2.0	
1	А	214	LYS	2.0	
1	А	223	THR	2.0	
1	А	335	ALA	2.0	

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
1	SEP	А	346	10/11	0.45	0.78	128,133,140,141	4
2	TPO	В	345	11/12	0.55	0.46	129,138,143,144	0
1	TPO	А	345	11/12	0.58	0.36	121,127,133,135	0
1	TPO	А	342	11/12	0.74	0.24	116,120,125,128	4
2	SEP	В	346	10/11	0.76	0.47	141,150,162,162	0

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
3	CJN	А	501	22/22	0.81	0.22	$34,\!45,\!47,\!51$	0
4	SO4	В	501	5/5	0.88	0.18	92,94,95,96	0
3	CJN	В	502	22/22	0.92	0.17	28,35,44,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers (i)

There are no such residues in this entry.

