



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 29, 2024 – 03:19 am BST

PDB ID : 6F3N
Title : Crystal structure of S-adenosyl-L-homocysteine hydrolase from *Pseudomonas aeruginosa* cocrystallized with SAH in the presence of K⁺ and Zn²⁺ cations
Authors : Czyrko, J.; Brzezinski, K.
Deposited on : 2017-11-28
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36.2
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	472	 94%
1	B	472	 94%
1	C	472	 94%
1	D	472	 95%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 16438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Adenosylhomocysteinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	460	Total 3613	C 2282	N 625	O 684	S 22	0	13	0
1	B	460	Total 3593	C 2269	N 622	O 680	S 22	0	8	0
1	C	460	Total 3605	C 2278	N 622	O 682	S 23	0	10	0
1	D	461	Total 3641	C 2299	N 629	O 690	S 23	0	15	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q9I685
A	-1	ASN	-	expression tag	UNP Q9I685
A	0	ALA	-	expression tag	UNP Q9I685
B	-2	SER	-	expression tag	UNP Q9I685
B	-1	ASN	-	expression tag	UNP Q9I685
B	0	ALA	-	expression tag	UNP Q9I685
C	-2	SER	-	expression tag	UNP Q9I685
C	-1	ASN	-	expression tag	UNP Q9I685
C	0	ALA	-	expression tag	UNP Q9I685
D	-2	SER	-	expression tag	UNP Q9I685
D	-1	ASN	-	expression tag	UNP Q9I685
D	0	ALA	-	expression tag	UNP Q9I685

- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	K 1	0	0
2	B	1	Total 1	K 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total K 1 1	0	0
2	D	1	Total K 1 1	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Zn 1 1	0	1
3	B	1	Total Zn 1 1	0	1
3	C	1	Total Zn 1 1	0	1
3	D	1	Total Zn 1 1	0	1

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	B	1	Total O P 5 4 1	0	0
4	C	1	Total O P 5 4 1	0	0

Continued on next page...



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
6	A	1	38	20	10	8	0	1
6	B	1	38	20	10	8	0	1
6	C	1	38	20	10	8	0	1
6	D	1	38	20	10	8	0	1

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	400	Total	O	0	13
			403	403		
8	B	468	Total	O	0	9
			473	473		
8	C	319	Total	O	0	6
			323	323		
8	D	424	Total	O	0	9
			425	425		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Adenosylhomocysteinase




- Molecule 1: Adenosylhomocysteinase



- Molecule 1: Adenosylhomocysteinase



- Molecule 1: Adenosylhomocysteinase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	170.59Å 99.28Å 111.72Å 90.00° 101.90° 90.00°	Depositor
Resolution (Å)	24.82 – 1.85 24.82 – 1.85	Depositor EDS
% Data completeness (in resolution range)	97.9 (24.82-1.85) 98.0 (24.82-1.85)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 1.85Å)	Xtrriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.151 , 0.176 0.161 , 0.187	Depositor DCC
R_{free} test set	1485 reflections (0.98%)	wwPDB-VP
Wilson B-factor (Å ²)	18.8	Xtrriage
Anisotropy	0.104	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 53.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16438	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.56 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.2122e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ADN, ZN, GOL, K, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.71	0/3701	0.91	13/5002 (0.3%)
1	B	0.72	0/3672	0.85	12/4964 (0.2%)
1	C	0.69	1/3693 (0.0%)	0.83	11/4990 (0.2%)
1	D	0.72	0/3729	0.84	7/5037 (0.1%)
All	All	0.71	1/14795 (0.0%)	0.86	43/19993 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	434	GLU	CD-OE2	6.05	1.32	1.25

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	351[A]	ARG	NE-CZ-NH2	-14.92	112.84	120.30
1	A	351[B]	ARG	NE-CZ-NH2	-14.92	112.84	120.30
1	A	351[A]	ARG	NE-CZ-NH1	13.21	126.90	120.30
1	A	351[B]	ARG	NE-CZ-NH1	13.21	126.90	120.30
1	D	315	ARG	NE-CZ-NH1	8.65	124.62	120.30
1	D	241	ARG	NE-CZ-NH1	7.51	124.06	120.30
1	B	428	ASP	CB-CG-OD1	7.37	124.93	118.30
1	D	315	ARG	NE-CZ-NH2	-7.30	116.65	120.30
1	B	372	ARG	NE-CZ-NH2	-7.28	116.66	120.30
1	B	241	ARG	NE-CZ-NH2	7.07	123.83	120.30
1	A	241	ARG	NE-CZ-NH1	6.89	123.74	120.30
1	D	373	LEU	CA-CB-CG	6.83	131.01	115.30
1	C	428	ASP	CB-CG-OD1	6.80	124.42	118.30
1	A	428	ASP	CB-CG-OD1	6.77	124.40	118.30
1	C	171	ARG	NE-CZ-NH1	6.76	123.68	120.30
1	A	315	ARG	NE-CZ-NH2	-6.75	116.92	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	373	LEU	CA-CB-CG	6.62	130.52	115.30
1	A	315	ARG	NE-CZ-NH1	6.49	123.55	120.30
1	D	171	ARG	NE-CZ-NH2	-6.36	117.12	120.30
1	D	428	ASP	CB-CG-OD1	6.25	123.93	118.30
1	C	43	ARG	NE-CZ-NH1	5.93	123.27	120.30
1	B	241	ARG	NE-CZ-NH1	-5.86	117.37	120.30
1	B	428	ASP	CB-CG-OD2	-5.86	113.03	118.30
1	C	43	ARG	NE-CZ-NH2	-5.75	117.42	120.30
1	C	171	ARG	NE-CZ-NH2	-5.59	117.50	120.30
1	B	465	ASP	CB-CG-OD1	5.59	123.33	118.30
1	D	372	ARG	NE-CZ-NH1	5.49	123.05	120.30
1	A	157	ARG	NE-CZ-NH1	5.38	122.99	120.30
1	A	428	ASP	CB-CG-OD2	-5.37	113.47	118.30
1	C	372	ARG	NE-CZ-NH2	-5.31	117.64	120.30
1	C	428	ASP	CB-CG-OD2	-5.31	113.52	118.30
1	C	385	ARG	NE-CZ-NH1	5.25	122.93	120.30
1	A	175	MET	CG-SD-CE	-5.21	91.87	100.20
1	A	372	ARG	NE-CZ-NH2	-5.18	117.71	120.30
1	C	465	ASP	CB-CG-OD1	5.15	122.94	118.30
1	B	132	ASP	CB-CG-OD1	5.14	122.93	118.30
1	B	17	ASP	CB-CG-OD2	-5.11	113.70	118.30
1	B	385	ARG	NE-CZ-NH1	5.09	122.85	120.30
1	C	258[A]	MET	CG-SD-CE	5.08	108.32	100.20
1	C	258[B]	MET	CG-SD-CE	5.08	108.32	100.20
1	B	43	ARG	NE-CZ-NH2	-5.07	117.76	120.30
1	B	373[A]	LEU	CB-CA-C	-5.04	100.63	110.20
1	B	373[B]	LEU	CB-CA-C	-5.04	100.63	110.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3613	0	3632	5	0
1	B	3593	0	3611	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3605	0	3632	4	0
1	D	3641	0	3662	5	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
4	C	5	0	0	0	0
4	D	5	0	0	0	0
5	A	44	0	26	2	0
5	B	44	0	26	3	0
5	C	44	0	26	0	0
5	D	44	0	26	2	0
6	A	38	0	26	2	0
6	B	38	0	26	2	0
6	C	38	0	26	0	0
6	D	38	0	26	2	0
7	D	6	0	8	0	0
8	A	403	0	0	1	0
8	B	473	0	0	0	0
8	C	323	0	0	0	0
8	D	425	0	0	3	0
All	All	16438	0	14753	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:287:LEU:HD12	8:D:878:HOH:O	1.96	0.64
1:B:140:GLY:HA3	1:B:323[B]:HIS:CE1	2.38	0.57
1:D:172:LEU:HD23	1:D:175[B]:MET:CE	2.33	0.57
1:B:373[A]:LEU:HD22	5:B:504:NAD:N7N	2.24	0.53
1:A:43[A]:ARG:NH1	8:A:604:HOH:O	2.42	0.52
5:A:504:NAD:C4N	6:A:505[A]:ADN:H3'	2.41	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:126:LYS:NZ	8:D:607:HOH:O	2.44	0.49
1:A:351[A]:ARG:HD3	1:A:363:ASP:OD1	2.13	0.49
1:C:140:GLY:HA3	1:C:323[B]:HIS:CE1	2.50	0.47
5:D:505:NAD:C4N	6:D:506[A]:ADN:H3'	2.45	0.47
5:A:504:NAD:C4N	6:A:505[B]:ADN:H3'	2.45	0.47
5:B:504:NAD:C4N	6:B:505[B]:ADN:H3'	2.45	0.47
1:B:11:THR:HB	1:B:13:TYR:CE2	2.52	0.45
5:D:505:NAD:C4N	6:D:506[B]:ADN:H3'	2.47	0.44
1:D:43:ARG:NH1	8:D:608:HOH:O	2.44	0.44
1:C:241[B]:ARG:NH2	1:C:263:ASP:O	2.35	0.44
5:B:504:NAD:C4N	6:B:505[A]:ADN:H3'	2.48	0.43
1:A:413:GLU:OE2	1:A:417:ARG:CZ	2.67	0.43
1:C:81:ARG:NH1	1:C:126:LYS:HD3	2.34	0.43
1:A:468:ARG:HB2	1:B:196:LYS:HE3	2.01	0.42
1:C:404:GLN:NE2	1:C:417:ARG:HD2	2.35	0.41
1:A:42:ARG:HG3	1:A:72:THR:HG23	2.01	0.41
1:D:140:GLY:HA3	1:D:323[B]:HIS:CE1	2.54	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	468/472 (99%)	459 (98%)	8 (2%)	1 (0%)	47 33
1	B	465/472 (98%)	453 (97%)	11 (2%)	1 (0%)	47 33
1	C	467/472 (99%)	459 (98%)	7 (2%)	1 (0%)	47 33
1	D	472/472 (100%)	462 (98%)	9 (2%)	1 (0%)	47 33
All	All	1872/1888 (99%)	1833 (98%)	35 (2%)	4 (0%)	47 33

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	61	HIS
1	B	61	HIS
1	C	61	HIS
1	D	61	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	387/385 (100%)	386 (100%)	1 (0%)	92	91
1	B	384/385 (100%)	384 (100%)	0	100	100
1	C	386/385 (100%)	386 (100%)	0	100	100
1	D	390/385 (101%)	389 (100%)	1 (0%)	92	91
All	All	1547/1540 (100%)	1545 (100%)	2 (0%)	93	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	373	LEU
1	D	373	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 25 ligands modelled in this entry, 8 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	ADN	B	505[B]	-	18,21,21	0.64	0	18,31,31	1.03	1 (5%)
6	ADN	D	506[B]	-	18,21,21	0.60	0	18,31,31	0.97	1 (5%)
5	NAD	D	505	-	42,48,48	0.56	0	50,73,73	0.86	2 (4%)
6	ADN	A	505[A]	-	18,21,21	0.62	0	18,31,31	0.93	1 (5%)
6	ADN	C	505[A]	-	18,21,21	0.62	0	18,31,31	0.90	1 (5%)
4	PO4	D	503	-	4,4,4	0.77	0	6,6,6	0.95	0
5	NAD	B	504	-	42,48,48	0.60	0	50,73,73	0.81	1 (2%)
4	PO4	A	503	-	4,4,4	0.77	0	6,6,6	0.68	0
6	ADN	B	505[A]	-	18,21,21	0.67	0	18,31,31	1.01	1 (5%)
5	NAD	A	504	-	42,48,48	0.57	0	50,73,73	0.88	3 (6%)
5	NAD	C	504	-	42,48,48	0.67	0	50,73,73	0.78	1 (2%)
6	ADN	A	505[B]	-	18,21,21	0.63	0	18,31,31	1.00	1 (5%)
4	PO4	B	503	-	4,4,4	0.81	0	6,6,6	0.70	0
6	ADN	C	505[B]	-	18,21,21	0.63	0	18,31,31	0.96	1 (5%)
4	PO4	C	503	-	4,4,4	1.05	0	6,6,6	1.02	0
6	ADN	D	506[A]	-	18,21,21	0.63	0	18,31,31	0.94	1 (5%)
7	GOL	D	504	-	5,5,5	0.51	0	5,5,5	0.92	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ADN	B	505[B]	-	-	0/2/22/22	0/3/3/3
6	ADN	D	506[B]	-	-	0/2/22/22	0/3/3/3
5	NAD	D	505	-	-	5/26/62/62	0/5/5/5
6	ADN	A	505[A]	-	-	2/2/22/22	0/3/3/3
6	ADN	C	505[A]	-	-	2/2/22/22	0/3/3/3
5	NAD	B	504	-	-	5/26/62/62	0/5/5/5
6	ADN	B	505[A]	-	-	1/2/22/22	0/3/3/3
6	ADN	D	506[A]	-	-	0/2/22/22	0/3/3/3
5	NAD	A	504	-	-	5/26/62/62	0/5/5/5
5	NAD	C	504	-	-	5/26/62/62	0/5/5/5
6	ADN	A	505[B]	-	-	0/2/22/22	0/3/3/3
7	GOL	D	504	-	-	0/4/4/4	-
6	ADN	C	505[B]	-	-	0/2/22/22	0/3/3/3

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	D	505	NAD	C2N-N1N-C1D	-2.78	112.95	119.14
5	A	504	NAD	PN-O3-PA	-2.75	123.38	132.83
5	D	505	NAD	PN-O3-PA	-2.61	123.88	132.83
6	D	506[B]	ADN	C5-C6-N6	2.47	124.11	120.35
6	B	505[A]	ADN	C5-C6-N6	2.46	124.09	120.35
6	D	506[A]	ADN	C5-C6-N6	2.45	124.08	120.35
6	B	505[B]	ADN	C5-C6-N6	2.42	124.03	120.35
5	B	504	NAD	C2N-N1N-C1D	-2.42	113.75	119.14
6	C	505[B]	ADN	C5-C6-N6	2.39	123.98	120.35
6	C	505[A]	ADN	C5-C6-N6	2.34	123.91	120.35
6	A	505[B]	ADN	C5-C6-N6	2.29	123.84	120.35
5	A	504	NAD	C2N-N1N-C1D	-2.29	114.03	119.14
6	A	505[A]	ADN	C5-C6-N6	2.20	123.69	120.35
5	C	504	NAD	C2N-N1N-C1D	-2.14	114.36	119.14
5	A	504	NAD	C5A-C6A-N6A	2.14	123.60	120.35

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	504	NAD	O4D-C1D-N1N-C2N
5	A	504	NAD	O4D-C1D-N1N-C6N
5	A	504	NAD	C2D-C1D-N1N-C2N

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	A	504	NAD	C2D-C1D-N1N-C6N
5	B	504	NAD	O4D-C1D-N1N-C2N
5	B	504	NAD	O4D-C1D-N1N-C6N
5	B	504	NAD	C2D-C1D-N1N-C2N
5	B	504	NAD	C2D-C1D-N1N-C6N
5	C	504	NAD	O4D-C1D-N1N-C2N
5	C	504	NAD	O4D-C1D-N1N-C6N
5	C	504	NAD	C2D-C1D-N1N-C2N
5	C	504	NAD	C2D-C1D-N1N-C6N
5	D	505	NAD	O4D-C1D-N1N-C2N
5	D	505	NAD	O4D-C1D-N1N-C6N
5	D	505	NAD	C2D-C1D-N1N-C2N
5	D	505	NAD	C2D-C1D-N1N-C6N
6	A	505[A]	ADN	C3'-C4'-C5'-O5'
6	C	505[A]	ADN	C3'-C4'-C5'-O5'
6	A	505[A]	ADN	O4'-C4'-C5'-O5'
6	C	505[A]	ADN	O4'-C4'-C5'-O5'
6	B	505[A]	ADN	C3'-C4'-C5'-O5'
5	A	504	NAD	O4B-C4B-C5B-O5B
5	C	504	NAD	O4B-C4B-C5B-O5B
5	B	504	NAD	O4B-C4B-C5B-O5B
5	D	505	NAD	O4B-C4B-C5B-O5B


There are no ring outliers.



9 monomers are involved in 7 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	505[B]	ADN	1	0
6	D	506[B]	ADN	1	0
5	D	505	NAD	2	0
6	A	505[A]	ADN	1	0
5	B	504	NAD	3	0
6	B	505[A]	ADN	1	0
5	A	504	NAD	2	0
6	A	505[B]	ADN	1	0
6	D	506[A]	ADN	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.



Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	460/472 (97%)	-0.29	6 (1%) 77 78	13, 19, 32, 45	14 (3%)
1	B	460/472 (97%)	-0.41	4 (0%) 84 84	12, 17, 29, 50	13 (2%)
1	C	460/472 (97%)	-0.20	8 (1%) 70 70	14, 21, 39, 54	18 (3%)
1	D	461/472 (97%)	-0.36	2 (0%) 92 92	13, 18, 29, 44	13 (2%)
All	All	1841/1888 (97%)	-0.31	20 (1%) 80 81	12, 19, 33, 54	58 (3%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	411	ALA	4.7
1	C	412	ALA	4.3
1	B	355	ASP	4.2
1	C	409	LEU	3.4
1	C	355	ASP	3.3
1	A	412	ALA	3.2
1	B	415	ALA	3.2
1	A	355	ASP	3.1
1	C	416	LYS	2.9
1	A	46	ALA	2.9
1	C	410	PRO	2.7
1	C	10	PHE	2.6
1	A	411	ALA	2.5
1	D	9	GLY	2.4
1	B	410	PRO	2.4
1	C	411	ALA	2.3
1	D	293	LEU	2.2
1	A	10	PHE	2.2
1	A	47	GLY	2.1
1	C	177	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	ADN	A	505[A]	19/19	0.92	0.12	17,18,26,27	19
6	ADN	A	505[B]	19/19	0.92	0.12	16,17,20,20	19
6	ADN	B	505[A]	19/19	0.92	0.13	14,16,25,25	19
6	ADN	B	505[B]	19/19	0.92	0.13	14,16,19,19	19
7	GOL	D	504	6/6	0.92	0.11	25,28,28,30	0
6	ADN	C	505[B]	19/19	0.93	0.10	17,17,19,20	19
6	ADN	D	506[A]	19/19	0.93	0.12	17,18,25,25	19
6	ADN	D	506[B]	19/19	0.93	0.12	17,18,21,21	19
6	ADN	C	505[A]	19/19	0.93	0.10	17,19,29,30	19
3	ZN	D	502[B]	1/1	0.96	0.08	27,27,27,27	1
5	NAD	B	504	44/44	0.97	0.07	14,15,17,18	0
5	NAD	C	504	44/44	0.97	0.07	14,19,21,21	0
5	NAD	D	505	44/44	0.97	0.07	14,16,19,21	0
5	NAD	A	504	44/44	0.97	0.07	14,17,19,19	0
4	PO4	A	503	5/5	0.98	0.08	22,22,24,25	0
4	PO4	C	503	5/5	0.98	0.15	24,25,26,28	0
3	ZN	A	502[B]	1/1	0.98	0.15	33,33,33,33	1
3	ZN	C	502[B]	1/1	0.99	0.21	27,27,27,27	1
4	PO4	D	503	5/5	0.99	0.11	21,22,22,24	0
2	K	D	501	1/1	0.99	0.04	15,15,15,15	0
3	ZN	B	502[B]	1/1	0.99	0.13	29,29,29,29	1
4	PO4	B	503	5/5	0.99	0.06	19,19,21,21	0
2	K	B	501	1/1	1.00	0.03	13,13,13,13	0
2	K	C	501	1/1	1.00	0.03	17,17,17,17	0
2	K	A	501	1/1	1.00	0.03	14,14,14,14	0




The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around ADN A 505 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ADN B 505 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADN B 505 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ADN C 505 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADN D 506 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ADN D 506 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ADN C 505 (A):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD B 504:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAD C 504:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAD D 505:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 504:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.