



# Full wwPDB X-ray Structure Validation Report ⓘ

May 16, 2020 – 04:00 am BST

PDB ID : 1F60  
Title : CRYSTAL STRUCTURE OF THE YEAST ELONGATION FACTOR COM-  
PLEX EEF1A:EEF1BA  
Authors : Andersen, G.R.; Pedersen, L.; Valente, L.; Kinzy, T.G.; Nyborg, J.  
Deposited on : 2000-06-19  
Resolution : 1.67 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	458	 5% <span style="margin-left: 150px;">83%</span> <span style="margin-left: 100px;">10%</span> . .
2	B	94	 % <span style="margin-left: 150px;">82%</span> <span style="margin-left: 100px;">13%</span> . .

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4824 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ELONGATION FACTOR EEF1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	440	3388	2152	592	630	14	0	0	0

- Molecule 2 is a protein called ELONGATION FACTOR EEF1BA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	90	703	441	112	146	4	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	567	Total	O	0	0
			567	567		
3	B	166	Total	O	0	0
			166	166		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ELONGATION FACTOR EEF1A



- Molecule 2: ELONGATION FACTOR EEF1BA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	63.85Å 91.81Å 92.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.67 19.72 – 1.67	Depositor EDS
% Data completeness (in resolution range)	99.3 (20.00-1.67) 99.2 (19.72-1.67)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.79 (at 1.67Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.187 , 0.221 0.179 , 0.212	Depositor DCC
$R_{free}$ test set	3217 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	14.4	Xtrriage
Anisotropy	0.334	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 53.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.004 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4824	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.47% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	0/3458	1.31	24/4674 (0.5%)
2	B	0.58	0/713	1.14	2/964 (0.2%)
All	All	0.59	0/4171	1.29	26/5638 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	29	TYR	CB-CG-CD2	-11.39	114.16	121.00
1	A	69	ARG	NE-CZ-NH1	10.63	125.61	120.30
1	A	69	ARG	NE-CZ-NH2	-9.95	115.32	120.30
1	A	146	ARG	NE-CZ-NH2	-9.94	115.33	120.30
1	A	29	TYR	CB-CG-CD1	9.02	126.41	121.00
1	A	279	ALA	CA-C-O	-8.94	101.32	120.10
1	A	37	ARG	CD-NE-CZ	8.63	135.68	123.60
1	A	96	ARG	NE-CZ-NH2	-8.02	116.29	120.30
1	A	280	PRO	CA-N-CD	-7.98	100.33	111.50
1	A	280	PRO	N-CA-CB	7.83	112.69	103.30
1	A	69	ARG	CD-NE-CZ	6.99	133.39	123.60
1	A	5	LYS	CB-CG-CD	6.82	129.33	111.60
1	A	280	PRO	N-CD-CG	6.51	112.97	103.20
2	B	1182	ASP	CB-CG-OD2	-6.41	112.53	118.30
1	A	134	ARG	NE-CZ-NH2	-6.28	117.16	120.30
1	A	98	PHE	CB-CG-CD1	-6.15	116.50	120.80

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	264	ARG	NE-CZ-NH2	6.09	123.35	120.30
1	A	146	ARG	NH1-CZ-NH2	5.65	125.61	119.40
1	A	428	ARG	NE-CZ-NH2	-5.65	117.48	120.30
1	A	37	ARG	NE-CZ-NH1	5.54	123.07	120.30
1	A	130	ASP	CB-CG-OD1	5.48	123.23	118.30
1	A	141	PHE	CB-CG-CD1	-5.27	117.11	120.80
1	A	68	GLU	OE1-CD-OE2	-5.24	117.01	123.30
1	A	156	ASP	CB-CG-OD1	-5.19	113.63	118.30
2	B	1148	MET	CA-CB-CG	5.02	121.83	113.30
1	A	375	GLU	CB-CA-C	-5.01	100.37	110.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	279	ALA	Mainchain,Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3388	0	3440	41	0
2	B	703	0	688	8	0
3	A	567	0	0	11	1
3	B	166	0	0	4	0
All	All	4824	0	4128	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance ( $\text{\AA}$ )	Clash overlap ( $\text{\AA}$ )
2:B:1148:MET:HE2	2:B:1151:LEU:HD22	1.28	1.10
1:A:2:GLY:HA3	3:A:2698:HOH:O	1.71	0.90

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:425:ARG:NH2	1:A:428:ARG:H	1.76	0.83
1:A:108:GLN:HG3	3:A:2474:HOH:O	1.82	0.80
1:A:294:HIS:CE1	2:B:1160:PRO:HG2	2.17	0.79
1:A:261:PRO:HD2	1:A:308:PHE:O	1.96	0.66
1:A:288:LYS:HE2	1:A:311:LYS:HG2	1.79	0.64
1:A:362:HIS:CD2	1:A:363:THR:H	2.16	0.64
1:A:251:VAL:HG12	1:A:315:VAL:HG12	1.81	0.61
1:A:333:LYS:HE3	1:A:441:LYS:HD3	1.82	0.61
1:A:37:ARG:HG2	1:A:37:ARG:HH11	1.66	0.60
2:B:1148:MET:CE	2:B:1151:LEU:HD22	2.18	0.60
1:A:36:LYS:HE2	3:A:2545:HOH:O	2.01	0.59
1:A:106:THR:HB	1:A:264:ARG:NH1	2.18	0.58
1:A:107:SER:H	1:A:264:ARG:HH12	1.52	0.57
2:B:1156:HIS:HD2	3:B:2171:HOH:O	1.87	0.56
1:A:365:HIS:HE1	3:A:2333:HOH:O	1.87	0.56
1:A:360:ASP:OD1	1:A:365:HIS:HD2	1.89	0.55
1:A:328:LYS:HD2	3:A:2329:HOH:O	2.06	0.54
1:A:26:HIS:HD2	3:A:2060:HOH:O	1.89	0.54
1:A:26:HIS:HE1	3:A:2042:HOH:O	1.91	0.52
1:A:2:GLY:CA	3:A:2698:HOH:O	2.42	0.52
1:A:156:ASP:OD2	1:A:195:ASN:ND2	2.41	0.52
1:A:362:HIS:HD2	1:A:363:THR:H	1.56	0.51
1:A:293:HIS:HD2	3:B:2515:HOH:O	1.94	0.50
1:A:358:VAL:HA	1:A:367:ALA:HA	1.94	0.50
1:A:362:HIS:HE1	1:A:409:CYS:O	1.95	0.50
1:A:208:TRP:O	1:A:210:LYS:HD2	2.12	0.49
1:A:108:GLN:OE1	1:A:143:LEU:HD13	2.15	0.47
1:A:5:LYS:HE2	1:A:81:GLU:OE1	2.14	0.46
2:B:1136:LEU:HB3	2:B:1158:PHE:CE1	2.52	0.45
2:B:1117:PRO:N	3:B:2582:HOH:O	2.48	0.45
2:B:1180:SER:HB3	2:B:1183:ASP:HB2	1.98	0.45
2:B:1148:MET:CE	2:B:1179:VAL:HG13	2.47	0.45
1:A:26:HIS:CE1	3:A:2042:HOH:O	2.69	0.44
1:A:425:ARG:NH2	1:A:428:ARG:N	2.56	0.43
1:A:254:ILE:CG2	1:A:257:ILE:HD12	2.49	0.43
1:A:84:LYS:HE3	3:A:2306:HOH:O	2.18	0.43
1:A:110:ASP:OD2	1:A:236:PRO:HG3	2.19	0.42
1:A:422:PHE:CZ	1:A:434:GLY:HA3	2.54	0.42
1:A:425:ARG:HG3	3:B:2621:HOH:O	2.19	0.42
1:A:7:HIS:CE1	1:A:86:GLN:OE1	2.73	0.42
1:A:107:SER:H	1:A:264:ARG:NH1	2.17	0.41

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:THR:HB	1:A:83:PRO:HD2	2.01	0.41
1:A:177:LYS:HE2	3:A:2561:HOH:O	2.19	0.41
1:A:5:LYS:HE3	1:A:82:THR:O	2.21	0.41
1:A:37:ARG:HD3	1:A:37:ARG:HA	1.73	0.41
1:A:82:THR:HB	1:A:83:PRO:CD	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:2165:HOH:O	3:A:2307:HOH:O[1_655]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/458 (96%)	426 (97%)	11 (2%)	1 (0%)	47	29
2	B	88/94 (94%)	88 (100%)	0	0	100	100
All	All	526/552 (95%)	514 (98%)	11 (2%)	1 (0%)	47	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	280	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	367/379 (97%)	361 (98%)	6 (2%)	62	46
2	B	79/82 (96%)	77 (98%)	2 (2%)	47	26
All	All	446/461 (97%)	438 (98%)	8 (2%)	59	40

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	TYR
1	A	37	ARG
1	A	108	GLN
1	A	293	HIS
1	A	328	LYS
1	A	425	ARG
2	B	1128	LYS
2	B	1192	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	HIS
1	A	26	HIS
1	A	86	GLN
1	A	205	ASN
1	A	293	HIS
1	A	294	HIS
1	A	296	GLN
1	A	362	HIS
1	A	365	HIS
1	A	377	ASN
1	A	388	HIS
2	B	1142	ASN
2	B	1156	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	440/458 (96%)	0.13	25 (5%) 23 25	8, 15, 35, 45	0
2	B	90/94 (95%)	-0.18	1 (1%) 80 83	9, 14, 23, 29	0
All	All	530/552 (96%)	0.08	26 (4%) 29 31	8, 15, 33, 45	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	106	THR	7.0
1	A	107	SER	5.6
1	A	294	HIS	5.5
1	A	315	VAL	5.0
1	A	194	TRP	4.2
1	A	316	LYS	4.0
1	A	348	PRO	3.9
1	A	237	SER	3.6
1	A	312	ASN	3.6
1	A	256	GLY	3.5
1	A	314	SER	3.4
1	A	349	GLY	3.3
2	B	1117	PRO	3.0
1	A	328	LYS	3.0
1	A	298	GLU	2.9
1	A	2	GLY	2.7
1	A	239	PRO	2.6
1	A	3	LYS	2.6
1	A	313	VAL	2.5
1	A	295	GLU	2.5
1	A	296	GLN	2.3
1	A	238	ARG	2.2
1	A	282	GLY	2.2
1	A	241	ASP	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	264	ARG	2.0
1	A	281	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.