

Full wwPDB X-ray Structure Validation Report (i)

Nov 21, 2023 – 07:27 AM JST

PDB ID : 7F8G

Title : Structure-activity relationship studies of allosteric inhibitors of EYA2 tyrosine

phosphatase

Authors: Anantharajan, J.; Baburajendran, N.

Deposited on : 2021-07-02

Resolution : 3.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS: 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 3.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	130704	1659 (3.60-3.40)
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)
RSRZ outliers	127900	1559 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	294	71%	16%	• 13%
1	В	294	71%	17%	12%



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 3995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Eyes absent homolog 2.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	В	259	Total 1992	C 1271	N 343	O 368	S 10	3	1	0
1	A	257	Total 1943	C 1247	N 325	O 360	S 11	8	1	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	245	GLY	-	expression tag	UNP O00167
В	246	PRO	-	expression tag	UNP O00167
В	247	LEU	-	expression tag	UNP O00167
В	248	GLY	-	expression tag	UNP O00167
В	249	SER	-	expression tag	UNP O00167
В	250	PRO	-	expression tag	UNP O00167
В	251	GLU	-	expression tag	UNP O00167
В	252	PHE	-	expression tag	UNP O00167
A	245	GLY	-	expression tag	UNP O00167
A	246	PRO	-	expression tag	UNP O00167
A	247	LEU	-	expression tag	UNP O00167
A	248	GLY	-	expression tag	UNP O00167
A	249	SER	-	expression tag	UNP O00167
A	250	PRO	-	expression tag	UNP O00167
A	251	GLU	-	expression tag	UNP O00167
A	252	PHE	-	expression tag	UNP O00167

• Molecule 2 is 3-phenoxy- $\{N\}-[(\{E\})-(5-pyrimidin-2-ylsulfanylfuran-2-yl)methylidene amino]benzamide (three-letter code: 1RI) (formula: <math>C_{22}H_{16}N_4O_3S$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms			ZeroOcc	AltConf			
2	D	1	Total	С	N	О	S	0	0	
2	Ъ	1	30	22	4	3	1	0		
9	Λ	1	Total	С	N	О	S	0	0	
2	A	1	30	22	4	3	1	U	U	



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Eyes absent homolog 2





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants	89.92Å 89.92Å 96.85Å	Donositon
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.96 - 3.49	Depositor
Resolution (A)	44.96 - 3.49	EDS
% Data completeness	96.7 (44.96-3.49)	Depositor
(in resolution range)	96.7 (44.96-3.49)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.20 (at 3.48Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
P.P.	0.213 , 0.267	Depositor
R, R_{free}	0.213 , 0.268	DCC
R_{free} test set	1080 reflections (10.04%)	wwPDB-VP
Wilson B-factor (Å ²)	82.7	Xtriage
Anisotropy	0.490	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.34, 43.0	EDS
L-test for twinning ²	$< L > = 0.44, < L^2> = 0.27$	Xtriage
	0.064 for -h,-k,l	
Estimated twinning fraction	0.069 for h,-h-k,-l	Xtriage
	0.052 for -k,-h,-l	
F_o, F_c correlation	0.91	EDS
Total number of atoms	3995	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.35% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 1RI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.25	0/1986	0.43	0/2705	
1	В	0.25	0/2035	0.44	0/2766	
All	All	0.25	0/4021	0.44	0/5471	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1943	0	1813	27	0
1	В	1992	0	1897	31	0
2	A	30	0	0	0	0
2	В	30	0	0	0	0
All	All	3995	0	3710	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



A	A	Interatomic	Clash
Atom-1	Atom-2	$\operatorname{distance}\ (\text{\AA})$	overlap (Å)
1:B:438:ARG:NH2	1:B:533:LEU:O	2.21	0.71
1:A:438:ARG:NH2	1:A:533:LEU:O	2.27	0.68
1:B:312:ILE:HD13	1:B:457:LYS:HG2	1.76	0.67
1:B:274:ASP:OD1	1:B:480:LYS:NZ	2.23	0.65
1:B:375:ASP:OD1	1:B:378:ARG:NH1	2.32	0.62
1:A:312:ILE:HD13	1:A:457:LYS:HG2	1.86	0.58
1:A:315:LEU:HD13	1:A:460:LEU:HD21	1.85	0.56
1:B:432:LEU:HD13	1:B:466:VAL:HG21	1.87	0.56
1:B:444:VAL:HG12	1:B:471:ASN:HB3	1.90	0.54
1:B:444:VAL:HG21	1:B:487:ILE:HD13	1.91	0.53
1:A:444:VAL:HG21	1:A:487:ILE:HD13	1.91	0.53
1:A:481:GLU:HG3	1:A:513:HIS:HE2	1.72	0.52
1:A:434:LEU:HD21	1:A:534:GLU:HG2	1.92	0.52
1:A:500:ILE:HD13	1:A:529:LEU:HD21	1.93	0.51
1:A:432:LEU:HD13	1:A:466:VAL:HG21	1.93	0.50
1:B:334:ASP:OD1	1:B:388:LYS:NZ	2.31	0.50
1:A:438:ARG:HB2	1:A:441:CYS:HB2	1.92	0.50
1:A:333:VAL:HG11	1:A:469:ILE:HG12	1.92	0.50
1:B:492:GLY:O	1:B:497:TYR:OH	2.25	0.50
1:A:444:VAL:HG12	1:A:471:ASN:HB3	1.93	0.49
1:B:458:VAL:HG13	1:B:463:LEU:HB2	1.94	0.49
1:B:533:LEU:HD23	1:B:538:LEU:HB2	1.95	0.48
1:B:300:THR:O	1:B:304:ILE:HG13	2.13	0.48
1:A:458:VAL:HG13	1:A:463:LEU:HB2	1.94	0.48
1:B:448:THR:HG23	1:B:480:LYS:HE2	1.96	0.47
1:B:410:TRP:CH2	1:B:460:LEU:HB3	2.50	0.47
1:B:480:LYS:HE3	1:B:505:GLU:HG2	1.97	0.46
1:A:452:ILE:HB	1:A:453:PRO:HD3	1.97	0.46
1:B:326:GLU:O	1:B:450:GLN:NE2	2.47	0.46
1:A:429:LEU:HD21	1:A:463:LEU:HD23	1.98	0.45
1:B:497:TYR:O	1:B:516:PRO:HD2	2.17	0.45
1:A:424:TRP:HH2	1:A:520:ILE:O	2.00	0.44
1:A:335:ASP:OD1	1:A:486:ARG:NH2	2.51	0.44
1:B:448:THR:CG2	1:B:480:LYS:HE2	2.47	0.44
1:B:281:ILE:CG1	1:B:449:THR:HG23	2.48	0.44
1:B:386:ARG:O	1:B:390:MET:HG2	2.18	0.44
1:A:316:ALA:HB1	1:A:322:PHE:HB2	2.00	0.44
1:A:350:PHE:O	1:A:383:ARG:NH2	2.51	0.43
1:A:410:TRP:CH2	1:A:460:LEU:HB3	2.53	0.43
1:B:424:TRP:HH2	1:B:520:ILE:O	2.02	0.43
1:A:274:ASP:HB3	1:A:501:GLY:HA3	2.00	0.43
1:B:281:ILE:HG12	1:B:449:THR:HG23	1.98	0.43

Continued on next page...



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance} ({ m \AA})$	overlap (Å)
1:A:391:TYR:HA	1:A:459:LEU:HD11	2.01	0.43
1:A:344:ASP:OD2	1:A:346:SER:OG	2.26	0.43
1:B:281:ILE:HD11	1:B:449:THR:HG23	2.01	0.42
1:B:331:ILE:HD12	1:B:478:THR:HG21	2.00	0.42
1:A:500:ILE:HG12	1:A:518:TRP:HB2	2.02	0.42
1:B:285:LEU:HD23	1:B:285:LEU:HA	1.91	0.42
1:B:445:LEU:HB2	1:B:467:PHE:CE1	2.55	0.42
1:A:331:ILE:HG13	1:A:474:SER:O	2.20	0.42
1:B:457:LYS:HB3	1:B:461:TYR:CE1	2.55	0.41
1:B:488:MET:HA	1:B:497:TYR:CE1	2.55	0.41
1:A:522:CYS:SG	1:A:523:HIS:N	2.93	0.41
1:B:269:ARG:HG2	1:B:496:VAL:HB	2.02	0.41
1:A:418:GLU:OE2	1:A:425:LEU:HB3	2.21	0.41
1:A:502:ASP:OD1	1:A:502:ASP:N	2.54	0.41
1:B:410:TRP:CZ2	1:B:460:LEU:HB3	2.55	0.40
1:B:373:GLY:O	1:B:377:MET:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	252/294 (86%)	235 (93%)	17 (7%)	0	100	100
1	В	256/294 (87%)	239 (93%)	17 (7%)	0	100	100
All	All	508/588 (86%)	474 (93%)	34 (7%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	193/253 (76%)	190 (98%)	3 (2%)	62 83
1	В	203/253 (80%)	198 (98%)	5 (2%)	47 75
All	All	396/506 (78%)	388 (98%)	8 (2%)	55 79

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	282	PHE
1	В	317	ASP
1	В	443	ASN
1	В	459	LEU
1	В	486	ARG
1	A	282	PHE
1	A	459	LEU
1	A	486	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	es Link	Bond lengths			Bond angles			
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
	2	1RI	В	601	-	28,33,33	0.27	0	33,43,43	0.60	1 (3%)
	2	1RI	A	601	-	28,33,33	0.27	0	33,43,43	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1RI	В	601	-	-	0/14/18/18	0/4/4/4
2	1RI	A	601	-	-	0/14/18/18	0/4/4/4

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^{o})$	$\operatorname{Ideal}(^{o})$
2	В	601	1RI	C5-O2-C4	2.05	123.61	118.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring



in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

5.7 Other polymers (i)

There are no such residues in this entry.

Torsions



Rings

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	257/294 (87%)	0.68	19 (7%) 14 14	53, 92, 131, 145	6 (2%)
1	В	259/294~(88%)	0.39	3 (1%) 79 73	52, 73, 99, 119	3 (1%)
All	All	516/588 (87%)	0.53	22 (4%) 35 31	52, 79, 123, 145	9 (1%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	357	SER	5.0
1	A	384	TYR	3.5
1	A	492	GLY	3.1
1	A	469	ILE	2.8
1	A	312	ILE	2.8
1	A	325	LEU	2.7
1	A	497	TYR	2.6
1	A	328	CYS	2.6
1	В	518	TRP	2.5
1	A	431	ALA	2.5
1	A	452	ILE	2.5
1	A	443	ASN	2.5
1	A	311	MET	2.5
1	В	357	SER	2.4
1	A	442	VAL	2.4
1	A	472	ILE	2.3
1	A	270	VAL	2.2
1	A	451	LEU	2.2
1	A	529	LEU	2.1
1	A	456	ALA	2.1
1	A	320	LEU	2.1
1	В	320	LEU	2.0



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	1RI	В	601	30/30	0.98	0.39	52,62,76,83	0
2	1RI	A	601	30/30	0.98	0.38	53,64,80,85	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 1RI B 601:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around 1RI A 601:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

