



Full wwPDB EM Validation Report ⓘ

Nov 23, 2022 – 02:12 AM JST

PDB ID : 7F9W
EMDB ID : EMD-31499
Title : CD25 in complex with Fab
Authors : Liu, C.
Deposited on : 2021-07-05
Resolution : 3.20 Å (reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	161	
2	C	220	
3	B	452	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4217 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-2 receptor subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	106	854	537	150	156	11	1	0

- Molecule 2 is a protein called Light chain of Fab.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	C	218	1694	1062	282	345	5	0	0


- Molecule 3 is a protein called Heavy chain of Fab.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	B	223	1669	1055	281	327	6	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Interleukin-2 receptor subunit alpha



- Molecule 2: Light chain of Fab



- Molecule 3: Heavy chain of Fab



GLY PRO
 SER THR
 VAL PHE
 LEU PHE
 PRO PRO
 LYS LYS
 PRO THR
 LYS ASP
 THR THR
 LEU MET
 ILE ILE
 SER SER
 ARG ARG
 THR THR
 PRO PRO
 VAL VAL
 ASP ASP
 VAL VAL
 HIS HIS
 GLU GLU
 VAL VAL
 THR THR
 PRO PRO
 GLU GLU
 VAL VAL
 THR THR
 ASP ASP
 TYR TYR

ASN SER
 THR TYR
 VAL ARG
 VAL VAL
 SER SER
 VAL VAL
 THR THR
 LYS LYS
 THR THR
 VAL VAL
 LEU LEU
 HIS HIS
 GLN GLN
 ASP ASP
 TRP TRP
 LEU LEU
 ASN ASN
 GLY GLY
 LYS LYS
 THR THR
 VAL VAL
 VAL VAL
 ASN ASN
 LYS LYS
 VAL VAL
 ALA ALA
 LEU LEU
 PRO PRO
 GLU GLU
 ASN ASN
 ALA ALA
 ASP ASP
 PRO PRO
 ILE ILE
 GLU GLU
 THR THR
 LYS LYS
 THR THR
 THR THR
 PRO PRO
 PHE PHE
 THR THR
 VAL VAL
 LYS LYS
 THR THR
 VAL VAL
 ASP ASP
 LYS LYS
 SER SER
 ARG ARG
 ASN ASN

GLU LEU
 THR THR
 LYS LYS
 ASN ASN
 VAL VAL
 VAL VAL
 SER SER
 LEU LEU
 THR THR
 LYS LYS
 THR THR
 CYS CYS
 LEU LEU
 VAL VAL
 HIS HIS
 LYS LYS
 GLY GLY
 PHE PHE
 TYR TYR
 SER SER
 ASP ASP
 THR THR
 ILE ILE
 ALA ALA
 VAL VAL
 THR THR
 LYS LYS
 THR THR
 THR THR
 PRO PRO
 PHE PHE
 THR THR
 VAL VAL
 LYS LYS
 THR THR
 VAL VAL
 ASP ASP
 SER SER
 ASP ASP
 GLY GLY
 SER SER
 THR THR
 PHE PHE
 LEU LEU
 TYR TYR
 SER SER
 LYS LYS
 THR THR
 LEU LEU
 THR THR
 VAL VAL
 ASP ASP
 LYS LYS
 SER SER
 ARG ARG

TRP GLN
 GLN GLY
 VAL VAL
 PHE PHE
 SER SER
 CYS CYS
 SER SER
 VAL VAL
 MET MET
 HIS HIS
 GLU GLU
 ALA ALA
 LEU LEU
 HIS HIS
 ASN ASN
 HIS HIS
 TYR TYR
 THR THR
 GLN GLN
 LYS LYS
 SER SER
 LEU LEU
 SER SER
 LEU LEU
 SER SER
 PRO PRO
 GLY GLY
 LYS LYS

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	149373	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	1.196	Depositor
Minimum map value	-0.649	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.023	Depositor
Recommended contour level	0.189	Depositor
Map size (\AA)	194.40001, 194.40001, 194.40001	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.54, 0.54, 0.54	Depositor

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/880	0.49	0/1193
2	C	0.37	0/1732	0.50	0/2354
3	B	0.35	0/1709	0.51	0/2325
All	All	0.34	0/4321	0.50	0/5872

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	854	0	799	16	0
2	C	1694	0	1636	34	0
3	B	1669	0	1642	27	0
All	All	4217	0	4077	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:111:GLU:OE2	2:C:172:GLN:NE2	2.11	0.83

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:34:ASN:HB3	2:C:36:LYS:HG3	1.66	0.77
2:C:83:SER:O	2:C:85:GLN:NE2	2.22	0.73
3:B:212:SER:HG	3:B:214:THR:HG1	1.36	0.71
3:B:100:ARG:NH2	3:B:102:ASP:OD2	2.27	0.68
3:B:202:THR:OG1	3:B:219:LYS:NZ	2.26	0.68
1:A:141:GLY:N	3:B:104:SER:OG	2.25	0.66
1:A:125:GLN:HE21	1:A:126:MET:H	1.45	0.64
3:B:71:THR:HB	3:B:80:TYR:HB2	1.82	0.61
3:B:39:GLN:HB2	3:B:45:LEU:HD23	1.83	0.60
3:B:140:THR:HB	3:B:144:THR:H	1.65	0.60
2:C:4:MET:HE3	2:C:96:GLN:HB2	1.85	0.59
3:B:177:ALA:HB2	3:B:187:LEU:HD23	1.85	0.57
2:C:119:PRO:HB3	2:C:145:PHE:HB3	1.86	0.56
1:A:141:GLY:H	3:B:104:SER:HG	1.54	0.56
2:C:142:LEU:HB2	2:C:181:LEU:HB3	1.88	0.56
2:C:55:TYR:HD2	2:C:56:TRP:CD1	2.25	0.55
2:C:141:LEU:HD11	3:B:190:VAL:HG11	1.88	0.55
3:B:135:PRO:HB3	3:B:147:LEU:HB3	1.89	0.54
1:A:47:THR:OG1	1:A:54:SER:O	2.21	0.53
2:C:6:GLN:HE22	2:C:93:TYR:HA	1.74	0.52
3:B:202:THR:HG1	3:B:219:LYS:HZ2	1.56	0.52
2:C:154:TRP:CG	2:C:185:LEU:HD23	2.44	0.52
3:B:50:ARG:NH2	3:B:99:GLU:OE2	2.40	0.52
1:A:48:GLY:HA2	1:A:53:SER:HA	1.92	0.52
2:C:116:VAL:HG12	2:C:147:PRO:HD3	1.92	0.52
3:B:102:ASP:OD1	3:B:103:TYR:N	2.42	0.52
3:B:36:TRP:HB2	3:B:48:MET:HB2	1.91	0.51
2:C:12:ALA:HA	2:C:111:GLU:O	2.11	0.51
2:C:29:VAL:HG11	2:C:96:GLN:HE22	1.77	0.49
3:B:6:GLN:NE2	3:B:96:CYS:SG	2.86	0.49
2:C:2:ILE:H	2:C:2:ILE:HD12	1.78	0.48
3:B:68:VAL:HG22	3:B:83:LEU:HD23	1.95	0.48
1:A:43:TYR:HD2	1:A:44:MET:O	1.98	0.47
2:C:71:SER:OG	2:C:72:GLY:N	2.48	0.47
2:C:143:ASN:HD21	3:B:192:THR:HG21	1.80	0.47
2:C:203:THR:HG22	2:C:210:PRO:HG3	1.95	0.47
3:B:133:LEU:HD21	3:B:150:LEU:HD22	1.96	0.47
2:C:146:TYR:CG	2:C:147:PRO:HA	2.50	0.46
2:C:155:LYS:HE3	2:C:158:ASN:C	2.36	0.46
3:B:153:ASP:HA	3:B:184:LEU:HB3	1.98	0.46
1:A:26:LEU:HD12	1:A:119:TYR:CE1	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:VAL:O	1:A:157:THR:HG23	2.15	0.46
1:A:26:LEU:HD11	1:A:117:ARG:HD2	1.97	0.45
2:C:55:TYR:CD2	2:C:56:TRP:HD1	2.34	0.45
2:C:191:ASP:HA	2:C:194:LYS:NZ	2.32	0.45
3:B:12:LYS:HD2	3:B:18:VAL:HB	1.97	0.45
1:A:125:GLN:NE2	1:A:126:MET:H	2.13	0.45
3:B:133:LEU:HD21	3:B:150:LEU:HB2	1.99	0.44
1:A:121:PHE:CD1	1:A:125:GLN:HG2	2.52	0.44
3:B:162:SER:O	3:B:206:ASN:N	2.50	0.44
2:C:4:MET:CE	2:C:96:GLN:HE21	2.31	0.44
2:C:119:PRO:HB2	2:C:142:LEU:HD12	1.98	0.43
2:C:154:TRP:CD2	2:C:185:LEU:HD23	2.53	0.43
2:C:142:LEU:HD11	2:C:202:VAL:HG11	2.00	0.43
2:C:172:GLN:HG2	2:C:177:SER:HA	1.99	0.43
3:B:128:PRO:HA	3:B:154:TYR:HB3	2.00	0.43
1:A:12:HIS:C	1:A:132:VAL:HG23	2.39	0.43
2:C:4:MET:HE1	2:C:96:GLN:HE21	1.84	0.42
1:A:21:LYS:O	1:A:24:THR:HG23	2.19	0.42
2:C:55:TYR:CD2	2:C:56:TRP:CD1	3.06	0.42
2:C:164:ASN:OD1	2:C:185:LEU:HB2	2.20	0.42
1:A:15:PHE:HB3	1:A:129:TYR:CE1	2.55	0.42
2:C:52:LEU:HD22	2:C:61:GLU:HG3	2.00	0.42
2:C:114:ARG:HD3	2:C:177:SER:O	2.20	0.42
3:B:2:VAL:HG22	3:B:26:GLY:HA3	2.02	0.41
3:B:221:GLU:HG2	3:B:222:PRO:HD2	2.01	0.41
1:A:138:LEU:HD22	1:A:162:ILE:HD11	2.02	0.41
3:B:70:LEU:HD23	3:B:70:LEU:HA	1.86	0.41
1:A:138:LEU:HD13	1:A:162:ILE:HD11	2.02	0.41
2:C:39:LEU:HD13	2:C:77:PHE:CD2	2.55	0.41
2:C:6:GLN:NE2	2:C:93:TYR:HA	2.36	0.41
2:C:36:LYS:HB3	2:C:56:TRP:CE3	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	101/161 (63%)	97 (96%)	4 (4%)	0	100	100
2	C	216/220 (98%)	205 (95%)	11 (5%)	0	100	100
3	B	221/452 (49%)	212 (96%)	9 (4%)	0	100	100
All	All	538/833 (65%)	514 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	93/143 (65%)	91 (98%)	2 (2%)	52	79
2	C	193/195 (99%)	193 (100%)	0	100	100
3	B	187/399 (47%)	186 (100%)	1 (0%)	88	95
All	All	473/737 (64%)	470 (99%)	3 (1%)	91	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	25[A]	MET
1	A	25[B]	MET
3	B	201	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	96	GLN
2	C	172	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-31499. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 177



Y Index: 190



Z Index: 189


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.189. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 22 nm³; this corresponds to an approximate mass of 20 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.312\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-31499 and PDB model 7F9W. Per-residue inclusion information can be found in section 3 on page 4.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.189 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.189).









9.4 Atom inclusion [i](#)



At the recommended contour level, 83% of all backbone atoms, 70% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.189) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7030	 0.4920
A	 0.5550	 0.4430
B	 0.7296	 0.5000
C	 0.7503	 0.5080

