



Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 02:28 am BST

PDB ID : 2FA0
Title : HMG-CoA synthase from Brassica juncea in complex with HMG-CoA and covalently bound to HMG-CoA
Authors : Pojer, F.; Ferrer, J.L.; Richard, S.B.; Noel, J.P.
Deposited on : 2005-12-06
Resolution : 2.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	450	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3725 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HMG-CoA synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	450	3510	2237	575	675	23	0	0	0

- Molecule 2 is 3-HYDROXY-3-METHYLGLUTARYL-COENZYME A (three-letter code: HMG) (formula: $C_{27}H_{39}N_7O_{20}P_3S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
2	A	1	116	54	14	40	6	2	0	1


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	99	Total	O	0	0
			99	99		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HMG-CoA synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	61.33Å 61.33Å 411.56Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	68.52 – 2.49 47.20 – 2.49	Depositor EDS
% Data completeness (in resolution range)	97.7 (68.52-2.49) 97.8 (47.20-2.49)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 2.48Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.188 , 0.272 0.191 , 0.197	Depositor DCC
R_{free} test set	859 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtrriage
Anisotropy	0.228	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3725	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HMG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.08	6/3586 (0.2%)	1.09	14/4844 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	116	ALA	C-N	16.94	1.73	1.34
1	A	117	CYS	CB-SG	-7.27	1.69	1.82
1	A	103	CYS	CB-SG	6.10	1.92	1.82
1	A	127	CYS	CB-SG	-5.69	1.72	1.81
1	A	101	GLU	CB-CG	5.17	1.61	1.52
1	A	95	PHE	CE2-CZ	5.09	1.47	1.37

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	116	ALA	C-N-CA	13.94	156.54	121.70
1	A	116	ALA	O-C-N	-13.18	101.61	122.70
1	A	116	ALA	CA-C-N	9.71	138.56	117.20
1	A	413	ARG	NE-CZ-NH1	8.51	124.55	120.30
1	A	218	ASP	CB-CG-OD1	8.17	125.65	118.30
1	A	218	ASP	CB-CG-OD2	-6.84	112.15	118.30
1	A	114	THR	CB-CA-C	-6.38	94.37	111.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	305	LEU	CA-CB-CG	6.12	129.38	115.30
1	A	103	CYS	CA-CB-SG	-5.85	103.47	114.00
1	A	365	MET	CG-SD-CE	5.45	108.92	100.20
1	A	413	ARG	NE-CZ-NH2	-5.37	117.61	120.30
1	A	153	GLU	N-CA-C	5.35	125.44	111.00
1	A	214	LEU	CA-CB-CG	5.28	127.44	115.30
1	A	139	ARG	NE-CZ-NH2	-5.19	117.70	120.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	THR	Peptide
1	A	117	CYS	Mainchain
1	A	149	ALA	Peptide
1	A	450	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3510	0	3456	114	0
2	A	116	0	77	23	0
3	A	99	0	0	12	0
All	All	3725	0	3533	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:116:ALA:C	1:A:117:CYS:N	1.73	1.41
1:A:117:CYS:SG	2:A:500[B]:HMG:C5	2.69	0.80
1:A:249:PRO:HG3	2:A:500[A]:HMG:C2P	2.17	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:PRO:HG3	2:A:500[A]:HMG:H2P2	1.70	0.72
1:A:288:SER:O	1:A:291:GLU:HB2	1.88	0.72
1:A:249:PRO:CG	2:A:500[A]:HMG:H2P2	2.19	0.72
1:A:288:SER:HB3	1:A:291:GLU:OE1	1.90	0.71
1:A:116:ALA:HB1	2:A:500[A]:HMG:O3	1.91	0.70
1:A:33:GLY:HA3	2:A:500[A]:HMG:N1A	2.06	0.70
1:A:185:HIS:HD2	3:A:597:HOH:O	1.75	0.70
1:A:343:LYS:O	1:A:347:LEU:HG	1.91	0.69
1:A:372:GLU:HA	1:A:372:GLU:OE1	1.94	0.67
1:A:287:LEU:HD22	1:A:291:GLU:HB3	1.76	0.67
1:A:114:THR:HG21	3:A:585:HOH:O	1.93	0.67
1:A:247:HIS:CD2	2:A:500[A]:HMG:O2	2.48	0.66
1:A:311:ASP:HB3	3:A:564:HOH:O	1.94	0.66
1:A:49:GLU:OE2	1:A:413:ARG:HD2	1.96	0.66
1:A:358:GLY:HA2	2:A:500[A]:HMG:O3	1.97	0.64
1:A:8:LEU:HD11	1:A:170:GLY:HA3	1.80	0.63
1:A:252:LYS:CE	2:A:500[A]:HMG:O8A	2.47	0.63
1:A:252:LYS:HE2	2:A:500[A]:HMG:O8A	1.99	0.63
1:A:450:LYS:O	1:A:451:LYS:HB3	1.99	0.62
1:A:200:GLU:HG3	1:A:414:TYR:CD1	2.34	0.62
1:A:205:ASP:OD2	1:A:208:LEU:HD23	1.99	0.61
1:A:396:HIS:HB3	3:A:589:HOH:O	1.98	0.61
1:A:258:PHE:CD2	1:A:314:VAL:CG2	2.84	0.61
1:A:307:LYS:O	1:A:311:ASP:OD1	2.20	0.60
1:A:340:VAL:O	1:A:344:HIS:HB3	2.02	0.60
1:A:424:GLU:H	1:A:424:GLU:CD	2.05	0.59
1:A:317:THR:HG22	1:A:339:LEU:HB2	1.83	0.59
1:A:116:ALA:O	1:A:117:CYS:N	2.32	0.58
1:A:117:CYS:SG	2:A:500[B]:HMG:O4	2.59	0.58
1:A:93:LYS:HD3	1:A:93:LYS:C	2.23	0.58
1:A:289:LEU:HA	1:A:292:SER:OG	2.03	0.57
1:A:237:SER:OG	1:A:239:ASN:HB2	2.05	0.57
1:A:74:GLN:O	1:A:139:ARG:HB3	2.06	0.56
1:A:49:GLU:HG2	1:A:421:THR:HG22	1.87	0.56
1:A:339:LEU:HD12	1:A:339:LEU:O	2.05	0.56
1:A:402:LYS:HE2	3:A:598:HOH:O	2.05	0.56
1:A:247:HIS:NE2	2:A:500[A]:HMG:O2	2.39	0.56
1:A:193:TYR:CE1	1:A:195:PRO:HG3	2.42	0.55
1:A:114:THR:CG2	3:A:585:HOH:O	2.53	0.55
1:A:297:ASP:HA	1:A:300:LYS:HD3	1.87	0.55
1:A:323:GLN:OE1	1:A:395:ARG:NH1	2.40	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:451:LYS:O	1:A:451:LYS:HG2	2.07	0.54
1:A:209:SER:O	2:A:500[B]:HMG:H2P1	2.09	0.53
1:A:373:ASN:CG	1:A:374:GLN:H	2.12	0.53
1:A:413:ARG:HD3	1:A:419:PHE:CE1	2.43	0.53
1:A:116:ALA:CB	2:A:500[A]:HMG:O3	2.56	0.52
1:A:266:PHE:HD1	1:A:267:LEU:HD23	1.75	0.52
1:A:183:GLY:HA2	3:A:555:HOH:O	2.10	0.52
1:A:117:CYS:SG	2:A:500[B]:HMG:O3	2.67	0.52
1:A:238:ILE:CD1	1:A:258:PHE:CE2	2.93	0.51
1:A:320:VAL:HG23	1:A:338:SER:HB3	1.92	0.51
1:A:289:LEU:HG	1:A:290:ASP:H	1.75	0.51
1:A:401:GLU:HB2	3:A:535:HOH:O	2.09	0.51
1:A:182:ARG:NH2	3:A:501:HOH:O	2.44	0.51
1:A:392:LEU:HB3	1:A:395:ARG:HH21	1.76	0.51
1:A:156:ALA:O	1:A:159:THR:OG1	2.30	0.50
1:A:137:ASP:OD2	1:A:139:ARG:HD3	2.11	0.50
1:A:78:LEU:C	1:A:78:LEU:HD23	2.33	0.49
1:A:401:GLU:OE1	1:A:401:GLU:HA	2.12	0.48
1:A:178:GLU:CG	1:A:369:ARG:HG2	2.44	0.48
1:A:443:LEU:HD12	1:A:445:ARG:HH11	1.78	0.47
1:A:34:LYS:NZ	2:A:500[B]:HMG:O2A	2.34	0.47
1:A:100:PHE:CD1	1:A:100:PHE:N	2.82	0.47
1:A:69:LYS:HB2	1:A:69:LYS:HE2	1.44	0.47
1:A:249:PRO:CG	2:A:500[A]:HMG:C2P	2.87	0.47
1:A:235:GLU:HG3	3:A:539:HOH:O	2.15	0.46
1:A:266:PHE:CD1	1:A:267:LEU:HD23	2.51	0.46
1:A:443:LEU:HD12	1:A:445:ARG:NH1	2.31	0.46
1:A:258:PHE:CD2	1:A:314:VAL:HG21	2.50	0.46
1:A:443:LEU:HB2	1:A:445:ARG:HD3	1.98	0.46
1:A:117:CYS:N	2:A:500[A]:HMG:O3	2.48	0.46
1:A:192:PHE:C	1:A:192:PHE:CD2	2.89	0.46
1:A:117:CYS:N	2:A:500[A]:HMG:O4	2.49	0.46
1:A:238:ILE:HD12	1:A:258:PHE:CZ	2.51	0.45
1:A:418:GLU:HA	1:A:440:VAL:O	2.16	0.45
1:A:98:GLN:O	1:A:101:GLU:HB2	2.16	0.45
1:A:200:GLU:HG3	1:A:414:TYR:CE1	2.50	0.45
1:A:225:CYS:SG	1:A:268:ARG:NH1	2.90	0.45
1:A:225:CYS:HB3	3:A:539:HOH:O	2.16	0.44
1:A:153:GLU:O	1:A:154:GLY:O	2.35	0.44
1:A:320:VAL:O	1:A:321:PRO:C	2.55	0.43
1:A:33:GLY:HA3	2:A:500[B]:HMG:N1A	2.33	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:78:LEU:C	1:A:78:LEU:CD2	2.86	0.43
1:A:77:ARG:HD3	1:A:79:GLU:OE2	2.19	0.43
1:A:249:PRO:HG3	2:A:500[A]:HMG:H2P1	1.97	0.43
1:A:310:TYR:CD1	1:A:314:VAL:HB	2.53	0.43
1:A:428:ASP:OD1	1:A:428:ASP:N	2.50	0.43
1:A:247:HIS:CD2	1:A:249:PRO:HD3	2.54	0.43
1:A:49:GLU:OE2	1:A:413:ARG:CD	2.66	0.43
1:A:249:PRO:CB	2:A:500[A]:HMG:H2P2	2.49	0.43
1:A:258:PHE:O	1:A:261:LEU:HB2	2.19	0.43
1:A:241:ALA:HB3	1:A:244:PHE:CE1	2.55	0.42
1:A:151:TYR:HB2	1:A:157:ARG:HA	2.01	0.42
1:A:346:ASP:O	1:A:350:LYS:NZ	2.44	0.42
1:A:6:GLY:HA3	1:A:175:ILE:O	2.20	0.42
1:A:181:LEU:HD22	1:A:224:LEU:HG	1.99	0.42
1:A:44:LEU:HD23	1:A:44:LEU:C	2.41	0.42
1:A:192:PHE:CD1	1:A:204:VAL:HG22	2.55	0.41
1:A:245:VAL:HG12	1:A:332:LEU:HD12	2.02	0.41
1:A:12:ILE:HG12	1:A:13:TYR:N	2.35	0.41
1:A:289:LEU:HD23	1:A:289:LEU:H	1.85	0.41
1:A:83:GLU:OE2	2:A:500[A]:HMG:O4	2.39	0.41
1:A:322:LYS:NZ	3:A:580:HOH:O	2.50	0.41
1:A:175:ILE:HD13	1:A:336:PHE:CZ	2.56	0.40
1:A:193:TYR:CD1	1:A:195:PRO:HG3	2.56	0.40
1:A:207:LYS:HZ3	1:A:208:LEU:HD13	1.86	0.40
1:A:238:ILE:CD1	1:A:258:PHE:CZ	3.04	0.40
1:A:252:LYS:HG3	1:A:256:LYS:HE3	2.03	0.40
1:A:296:ARG:O	1:A:297:ASP:C	2.58	0.40
1:A:75:ILE:HD13	1:A:140:TYR:HB2	2.04	0.40
1:A:280:LYS:HD3	1:A:305:LEU:HD11	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	448/450 (100%)	407 (91%)	34 (8%)	7 (2%)	9 17

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	153	GLU
1	A	444	TYR
1	A	154	GLY
1	A	206	GLY
1	A	32	LYS
1	A	101	GLU
1	A	111	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	381/381 (100%)	333 (87%)	48 (13%)	4 8

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	LEU
1	A	69	LYS
1	A	78	LEU
1	A	86	ILE
1	A	93	LYS
1	A	100	PHE
1	A	101	GLU
1	A	103	CYS
1	A	114	THR
1	A	150	VAL
1	A	181	LEU
1	A	182	ARG
1	A	186	MET
1	A	189	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	192	PHE
1	A	196	ASN
1	A	197	LEU
1	A	207	LYS
1	A	208	LEU
1	A	214	LEU
1	A	215	MET
1	A	230	LYS
1	A	231	LEU
1	A	234	LYS
1	A	238	ILE
1	A	252	LYS
1	A	262	LEU
1	A	269	ASN
1	A	271	SER
1	A	273	ILE
1	A	278	LYS
1	A	282	THR
1	A	289	LEU
1	A	290	ASP
1	A	298	LEU
1	A	300	LYS
1	A	304	GLN
1	A	319	LEU
1	A	338	SER
1	A	346	ASP
1	A	372	GLU
1	A	375	SER
1	A	380	SER
1	A	405	GLU
1	A	413	ARG
1	A	423	LYS
1	A	444	TYR
1	A	451	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	185	HIS
1	A	304	GLN
1	A	344	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HMG	A	500[B]	-	49,60,60	1.01	3 (6%)	59,90,90	2.45	8 (13%)
2	HMG	A	500[A]	1	49,60,60	1.00	2 (4%)	59,90,90	1.93	9 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HMG	A	500[B]	-	-	24/54/77/77	0/3/3/3
2	HMG	A	500[A]	1	-	15/54/77/77	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500[A]	HMG	C1-S1P	-3.56	1.67	1.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	500[B]	HMG	C1-S1P	-3.00	1.68	1.76
2	A	500[A]	HMG	C2-C3	-2.54	1.52	1.54
2	A	500[B]	HMG	C6-C3	2.16	1.55	1.52
2	A	500[B]	HMG	O7-C3	2.05	1.47	1.44

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500[B]	HMG	C2-C1-S1P	12.20	128.73	113.63
2	A	500[B]	HMG	O2-C1-S1P	-6.83	113.75	122.61
2	A	500[A]	HMG	C3P-N4P-C5P	6.80	135.46	122.84
2	A	500[B]	HMG	C7P-N8P-C9P	6.21	133.66	122.59
2	A	500[A]	HMG	C7P-N8P-C9P	5.89	133.09	122.59
2	A	500[A]	HMG	N3A-C2A-N1A	-5.87	119.50	128.68
2	A	500[B]	HMG	O6A-CCP-CBP	5.07	118.70	110.55
2	A	500[B]	HMG	N3A-C2A-N1A	-5.06	120.76	128.68
2	A	500[A]	HMG	C2-C1-S1P	4.54	119.25	113.63
2	A	500[A]	HMG	C2P-S1P-C1	4.21	114.97	101.87
2	A	500[B]	HMG	O2-C1-C2	-3.85	117.48	123.64
2	A	500[B]	HMG	CEP-CBP-CAP	3.01	114.05	108.82
2	A	500[A]	HMG	O2-C1-C2	-3.01	118.82	123.64
2	A	500[A]	HMG	C1B-N9A-C4A	-2.69	121.92	126.64
2	A	500[B]	HMG	C2P-S1P-C1	2.13	108.51	101.87
2	A	500[A]	HMG	O3B-C3B-C4B	-2.12	102.41	110.08
2	A	500[A]	HMG	C2A-N1A-C6A	2.05	122.26	118.75

There are no chirality outliers.

All (39) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500[B]	HMG	CCP-O6A-P2A-O3A
2	A	500[B]	HMG	C5B-O5B-P1A-O1A
2	A	500[B]	HMG	C5B-O5B-P1A-O2A
2	A	500[B]	HMG	C9P-CAP-CBP-CEP
2	A	500[B]	HMG	N8P-C9P-CAP-OAP
2	A	500[B]	HMG	O2-C1-S1P-C2P
2	A	500[B]	HMG	C2-C1-S1P-C2P
2	A	500[B]	HMG	C1-C2-C3-O7
2	A	500[B]	HMG	C1-C2-C3-C4
2	A	500[B]	HMG	C1-C2-C3-C6
2	A	500[A]	HMG	CCP-O6A-P2A-O3A
2	A	500[A]	HMG	C5B-O5B-P1A-O3A

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	500[A]	HMG	C5B-O5B-P1A-O1A
2	A	500[A]	HMG	C5B-O5B-P1A-O2A
2	A	500[A]	HMG	O4B-C4B-C5B-O5B
2	A	500[A]	HMG	C2P-C3P-N4P-C5P
2	A	500[A]	HMG	C3P-C2P-S1P-C1
2	A	500[A]	HMG	C2-C3-C4-C5
2	A	500[A]	HMG	O7-C3-C4-C5
2	A	500[A]	HMG	C6-C3-C4-C5
2	A	500[B]	HMG	O4B-C4B-C5B-O5B
2	A	500[A]	HMG	C3B-C4B-C5B-O5B
2	A	500[A]	HMG	C6P-C7P-N8P-C9P
2	A	500[B]	HMG	O9P-C9P-CAP-OAP
2	A	500[B]	HMG	OAP-CAP-CBP-CDP
2	A	500[B]	HMG	OAP-CAP-CBP-CEP
2	A	500[B]	HMG	C4B-C5B-O5B-P1A
2	A	500[B]	HMG	C9P-CAP-CBP-CDP
2	A	500[B]	HMG	C5B-O5B-P1A-O3A
2	A	500[B]	HMG	CCP-O6A-P2A-O5A
2	A	500[B]	HMG	CCP-O6A-P2A-O4A
2	A	500[A]	HMG	CCP-O6A-P2A-O5A
2	A	500[A]	HMG	CCP-O6A-P2A-O4A
2	A	500[B]	HMG	C9P-CAP-CBP-CCP
2	A	500[B]	HMG	C2-C3-C4-C5
2	A	500[B]	HMG	CDP-CBP-CCP-O6A
2	A	500[B]	HMG	CEP-CBP-CCP-O6A
2	A	500[B]	HMG	C3B-O3B-P3B-O8A
2	A	500[A]	HMG	CBP-CCP-O6A-P2A


There are no ring outliers.

2 monomers are involved in 23 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500[B]	HMG	6	0
2	A	500[A]	HMG	17	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	116:ALA	C	117:CYS	N	1.73

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	450/450 (100%)	-0.03	8 (1%) 68 71	5, 27, 55, 68	1 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	117	CYS	6.0
1	A	273	ILE	3.2
1	A	32	LYS	2.6
1	A	284	TYR	2.5
1	A	294	GLN	2.3
1	A	289	LEU	2.2
1	A	286	SER	2.2
1	A	266	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	HMG	A	500[B]	58/58	0.92	0.22	2,40,59,60	58
2	HMG	A	500[A]	58/58	0.92	0.22	16,28,37,38	58

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HMG A 500 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.