



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 26, 2023 – 01:16 PM EDT

PDB ID : 3FAP
Title : ATOMIC STRUCTURES OF THE RAPAMYCIN ANALOGS IN COMPLEX WITH BOTH HUMAN FKBP12 AND FRB DOMAIN OF FRAP
Authors : Liang, J.; Clardy, J.
Deposited on : 1999-05-06
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	107	 3% 64% 26% 7%
2	B	94	 2% 60% 29% 11%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1890 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called FK506-BINDING PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	107	832	527	146	155	4	0	0	0

- Molecule 2 is a protein called FKBP12-RAPAMYCIN ASSOCIATED PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	94	796	508	139	142	7	0	0	0

- Molecule 3 is C15-(R)-METHYLTHIENYL RAPAMYCIN (three-letter code: ARD) (formula: $C_{55}H_{81}NO_{12}S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	69	55	1	12	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	105	Total 105	O 105	0	0
4	B	88	Total 88	O 88	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: FK506-BINDING PROTEIN



- Molecule 2: FKBP12-RAPAMYCIN ASSOCIATED PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	43.40Å 51.72Å 101.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.85 25.06 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.4 (20.00-1.85) 99.5 (25.06-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	4.70	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.68 (at 1.85Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.207 , 0.273 0.202 , 0.272	Depositor DCC
R_{free} test set	1980 reflections (9.85%)	wwPDB-VP
Wilson B-factor (Å ²)	28.1	Xtrriage
Anisotropy	0.553	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	1890	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.84% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ARD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.44	8/851 (0.9%)	2.31	38/1146 (3.3%)
2	B	1.50	3/818 (0.4%)	2.21	33/1099 (3.0%)
All	All	1.47	11/1669 (0.7%)	2.26	71/2245 (3.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
2	B	0	1
All	All	0	2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	39	SER	C-N	9.50	1.55	1.34
2	B	135	GLY	C-O	-8.03	1.10	1.23
1	A	42	ARG	NE-CZ	6.97	1.42	1.33
2	B	153	THR	CB-OG1	5.98	1.55	1.43
1	A	59	TRP	N-CA	5.71	1.57	1.46
1	A	60	GLU	CD-OE2	-5.57	1.19	1.25
2	B	115	MET	N-CA	5.52	1.57	1.46
1	A	29	MET	CA-CB	5.48	1.66	1.53
1	A	8	SER	C-O	5.23	1.33	1.23
1	A	26	TYR	CG-CD1	5.23	1.46	1.39
1	A	40	ARG	CA-CB	5.12	1.65	1.53

All (71) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	175	ARG	CD-NE-CZ	17.89	148.64	123.60
2	B	198	ARG	NE-CZ-NH1	16.33	128.47	120.30
1	A	40	ARG	NE-CZ-NH2	15.02	127.81	120.30
1	A	13	ARG	NE-CZ-NH2	-12.63	113.99	120.30
1	A	57	ARG	NE-CZ-NH2	-11.78	114.41	120.30
1	A	13	ARG	CD-NE-CZ	11.67	139.94	123.60
2	B	198	ARG	NE-CZ-NH2	-11.65	114.48	120.30
1	A	40	ARG	NE-CZ-NH1	-10.49	115.06	120.30
1	A	100	ASP	CB-CG-OD1	9.89	127.20	118.30
1	A	100	ASP	CB-CG-OD2	-9.51	109.74	118.30
1	A	11	ASP	CB-CG-OD2	9.39	126.75	118.30
2	B	199	ARG	NE-CZ-NH1	8.82	124.71	120.30
1	A	42	ARG	CD-NE-CZ	-8.71	111.41	123.60
1	A	60	GLU	CG-CD-OE2	-8.49	101.32	118.30
2	B	118	GLU	OE1-CD-OE2	8.24	133.19	123.30
1	A	60	GLU	CG-CD-OE1	8.12	134.53	118.30
2	B	125	ARG	NE-CZ-NH2	-8.04	116.28	120.30
2	B	108	VAL	CA-CB-CG1	8.00	122.90	110.90
2	B	138	GLU	OE1-CD-OE2	7.89	132.77	123.30
2	B	131	ARG	NE-CZ-NH1	-7.82	116.39	120.30
2	B	122	GLU	OE1-CD-OE2	-7.69	114.08	123.30
2	B	115	MET	CA-CB-CG	7.65	126.31	113.30
2	B	183	VAL	CG1-CB-CG2	-7.59	98.76	110.90
2	B	114	GLU	OE1-CD-OE2	7.54	132.35	123.30
1	A	87	HIS	CA-C-O	7.47	135.78	120.10
2	B	165	ARG	NE-CZ-NH2	7.45	124.02	120.30
1	A	13	ARG	NE-CZ-NH1	7.42	124.01	120.30
1	A	41	ASP	CB-CG-OD2	7.32	124.88	118.30
2	B	199	ARG	NE-CZ-NH2	-7.04	116.78	120.30
2	B	112	TRP	CE3-CZ3-CH2	-6.77	113.75	121.20
2	B	173	TRP	CZ3-CH2-CZ2	6.72	129.66	121.60
2	B	146	MET	N-CA-CB	-6.71	98.53	110.60
2	B	173	TRP	CH2-CZ2-CE2	-6.66	110.75	117.40
2	B	146	MET	CB-CA-C	6.61	123.62	110.40
1	A	50	LEU	CA-CB-CG	6.57	130.40	115.30
2	B	149	ARG	NE-CZ-NH1	-6.45	117.08	120.30
1	A	75	THR	CA-CB-CG2	-6.36	103.49	112.40
1	A	8	SER	N-CA-CB	6.35	120.02	110.50
1	A	88	PRO	O-C-N	-6.18	112.70	123.20
1	A	71	ARG	NE-CZ-NH1	6.17	123.38	120.30
1	A	101	VAL	CA-CB-CG1	6.12	120.08	110.90
1	A	87	HIS	CB-CA-C	6.05	122.50	110.40
1	A	41	ASP	OD1-CG-OD2	-5.94	112.01	123.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	171	GLN	CG-CD-OE1	-5.78	110.04	121.60
2	B	188	GLN	CG-CD-OE1	-5.78	110.04	121.60
2	B	152	GLN	OE1-CD-NE2	5.75	135.12	121.90
1	A	43	ASN	OD1-CG-ND2	-5.72	108.74	121.90
2	B	183	VAL	CA-CB-CG1	5.65	119.37	110.90
2	B	152	GLN	CG-CD-NE2	-5.64	103.17	116.70
1	A	36	PHE	CB-CG-CD2	5.61	124.73	120.80
1	A	73	LYS	CG-CD-CE	-5.60	95.09	111.90
1	A	87	HIS	O-C-N	-5.58	110.50	121.10
1	A	40	ARG	CG-CD-NE	-5.54	100.16	111.80
2	B	171	GLN	OE1-CD-NE2	5.50	134.55	121.90
1	A	95	ALA	N-CA-CB	5.43	117.71	110.10
1	A	93	PRO	N-CD-CG	-5.42	95.07	103.20
2	B	180	SER	O-C-N	-5.33	114.14	123.20
2	B	166	ASP	CB-CG-OD1	5.33	123.09	118.30
1	A	101	VAL	CA-CB-CG2	5.27	118.80	110.90
2	B	122	GLU	CG-CD-OE1	5.22	128.75	118.30
1	A	88	PRO	CA-C-N	5.22	126.64	116.20
1	A	11	ASP	CA-CB-CG	5.18	124.80	113.40
1	A	61	GLU	OE1-CD-OE2	5.18	129.51	123.30
1	A	30	LEU	CB-CG-CD2	-5.13	102.28	111.00
1	A	11	ASP	CA-C-N	5.09	126.38	116.20
1	A	71	ARG	NH1-CZ-NH2	-5.08	113.81	119.40
1	A	72	ALA	N-CA-CB	5.06	117.18	110.10
2	B	172	GLU	OE1-CD-OE2	-5.04	117.25	123.30
1	A	39	SER	C-N-CA	-5.03	109.13	121.70
2	B	190	TRP	CA-CB-CG	-5.01	104.17	113.70
2	B	118	GLU	CB-CG-CD	-5.00	100.69	114.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	88	PRO	Mainchain
2	B	187	THR	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	832	0	831	17	0
2	B	796	0	760	22	0
3	A	69	0	81	2	0
4	A	105	0	0	3	0
4	B	88	0	0	5	0
All	All	1890	0	1672	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:115:MET:SD	2:B:146:MET:SD	2.64	0.95
1:A:3:GLN:HA	1:A:3:GLN:HE21	1.53	0.74
1:A:13:ARG:HB3	1:A:13:ARG:HH11	1.56	0.70
2:B:187:THR:HG22	4:B:363:HOH:O	1.92	0.70
2:B:166:ASP:OD2	2:B:199:ARG:NH2	2.29	0.66
2:B:115:MET:HG2	2:B:143:LEU:CD2	2.26	0.66
2:B:108:VAL:HG22	4:B:282:HOH:O	1.97	0.64
1:A:44:LYS:HD2	4:A:373:HOH:O	1.97	0.64
2:B:134:LYS:O	2:B:138:GLU:HG3	1.98	0.61
2:B:195:HIS:CD2	2:B:199:ARG:HH21	2.18	0.61
2:B:108:VAL:HA	4:B:346:HOH:O	2.00	0.59
2:B:115:MET:SD	2:B:146:MET:CE	2.91	0.59
1:A:88:PRO:O	2:B:183:VAL:HG11	2.04	0.57
1:A:13:ARG:HH11	1:A:13:ARG:CB	2.17	0.56
1:A:60:GLU:HG3	4:A:244:HOH:O	2.06	0.54
2:B:184:LYS:NZ	2:B:187:THR:HG21	2.24	0.53
1:A:90:ILE:HG12	2:B:184:LYS:HE2	1.91	0.52
2:B:174:CYS:O	2:B:178:MET:HG3	2.11	0.50
1:A:92:PRO:HG2	4:A:328:HOH:O	2.11	0.50
1:A:89:GLY:HA3	2:B:183:VAL:HG12	1.92	0.50
1:A:13:ARG:HB3	1:A:13:ARG:NH1	2.24	0.49
2:B:115:MET:HG2	2:B:143:LEU:HD22	1.93	0.49
1:A:90:ILE:HG13	2:B:183:VAL:HG13	1.93	0.49
3:A:402:ARD:C18	3:A:402:ARD:H30	2.42	0.48
1:A:90:ILE:HG12	2:B:184:LYS:CE	2.44	0.48
1:A:87:HIS:CD2	1:A:90:ILE:HD12	2.49	0.47
1:A:51:GLY:N	1:A:60:GLU:OE2	2.30	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:17:LYS:N	1:A:20:GLN:OE1	2.43	0.45
2:B:149:ARG:NH1	4:B:347:HOH:O	2.47	0.45
2:B:187:THR:CG2	4:B:363:HOH:O	2.58	0.45
3:A:402:ARD:H2O	2:B:194:TYR:CD1	2.52	0.45
1:A:48:PHE:HE2	1:A:50:LEU:HD13	1.83	0.43
2:B:149:ARG:HH11	2:B:149:ARG:HB3	1.85	0.41
2:B:141:GLU:N	2:B:142:PRO:HD2	2.35	0.41
1:A:48:PHE:CE2	1:A:50:LEU:HD13	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	105/107 (98%)	101 (96%)	4 (4%)	0	100	100
2	B	92/94 (98%)	91 (99%)	1 (1%)	0	100	100
All	All	197/201 (98%)	192 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	89/89 (100%)	80 (90%)	9 (10%)	7	1

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	B	82/82 (100%)	77 (94%)	5 (6%)	18 5
All	All	171/171 (100%)	157 (92%)	14 (8%)	11 2

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	8	SER
1	A	13	ARG
1	A	17	LYS
1	A	25	HIS
1	A	44	LYS
1	A	50	LEU
1	A	73	LYS
1	A	79	ASP
2	B	113	HIS
2	B	149	ARG
2	B	165	ARG
2	B	182	ASN
2	B	184	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	GLN
1	A	43	ASN
1	A	87	HIS
2	B	160	ASN
2	B	182	ASN
2	B	188	GLN
2	B	195	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ARD	A	402	-	68,73,73	2.63	33 (48%)	72,104,104	2.74	31 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ARD	A	402	-	-	1/79/126/126	0/4/5/5

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	ARD	C11-C10	6.68	1.63	1.53
3	A	402	ARD	O1-C1	5.21	1.46	1.34
3	A	402	ARD	C14-C13	5.21	1.64	1.52
3	A	402	ARD	C52-S1	-5.06	1.64	1.74
3	A	402	ARD	C32-C33	4.68	1.63	1.52
3	A	402	ARD	C7-C8	-4.63	1.47	1.53
3	A	402	ARD	C37-C36	4.62	1.64	1.52
3	A	402	ARD	C30-C29	4.32	1.57	1.51
3	A	402	ARD	O4-C8	4.10	1.29	1.22
3	A	402	ARD	C18-C17	3.91	1.55	1.43
3	A	402	ARD	C2-N1	-3.78	1.42	1.47
3	A	402	ARD	C17-C16	-3.63	1.30	1.33
3	A	402	ARD	C34-C33	3.58	1.56	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	ARD	O13-C39	3.48	1.50	1.43
3	A	402	ARD	C6-N1	3.41	1.53	1.47
3	A	402	ARD	C42-C10	3.20	1.60	1.53
3	A	402	ARD	O1-C33	-3.07	1.40	1.46
3	A	402	ARD	C39-C38	3.00	1.57	1.52
3	A	402	ARD	O6-C9	2.89	1.44	1.39
3	A	402	ARD	C53-C52	2.88	1.46	1.37
3	A	402	ARD	O12-C51	2.84	1.52	1.42
3	A	402	ARD	C14-C15	2.82	1.57	1.54
3	A	402	ARD	C48-C34	2.75	1.59	1.53
3	A	402	ARD	C55-C52	-2.59	1.42	1.50
3	A	402	ARD	O3-C7	2.58	1.28	1.23
3	A	402	ARD	O8-C25	2.55	1.25	1.21
3	A	402	ARD	O9-C26	2.52	1.50	1.41
3	A	402	ARD	C7-N1	-2.40	1.31	1.34
3	A	402	ARD	C26-C25	2.37	1.60	1.51
3	A	402	ARD	C32-C31	2.22	1.56	1.51
3	A	402	ARD	C43-C16	-2.17	1.46	1.50
3	A	402	ARD	C5-C4	2.16	1.60	1.51
3	A	402	ARD	O5-C13	2.15	1.49	1.44

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	ARD	C20-C19-C18	-7.29	108.60	124.81
3	A	402	ARD	C8-C7-N1	7.10	127.37	119.25
3	A	402	ARD	O1-C33-C34	-6.74	95.64	107.88
3	A	402	ARD	C18-C17-C16	-5.62	119.07	127.32
3	A	402	ARD	C55-C52-S1	5.42	125.86	119.88
3	A	402	ARD	C5-C6-N1	-5.32	102.33	110.67
3	A	402	ARD	C48-C34-C35	-4.75	103.57	110.69
3	A	402	ARD	C47-C30-C29	-4.19	106.83	110.75
3	A	402	ARD	O4-C8-C7	4.07	123.67	119.08
3	A	402	ARD	C32-C31-C30	-3.85	112.89	117.72
3	A	402	ARD	C43-C16-C17	3.81	128.74	124.03
3	A	402	ARD	C40-C41-C36	3.77	118.65	112.42
3	A	402	ARD	C12-C11-C10	-3.75	106.67	112.65
3	A	402	ARD	C37-C36-C41	-3.70	102.56	109.17
3	A	402	ARD	C1-C2-N1	3.63	118.81	112.05
3	A	402	ARD	C14-C13-C12	-3.18	106.49	113.83
3	A	402	ARD	C4-C3-C2	-3.04	105.56	111.23
3	A	402	ARD	O2-C1-C2	-3.02	117.68	124.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	ARD	O11-C31-C30	2.96	124.97	120.93
3	A	402	ARD	C44-C22-C23	-2.93	105.83	110.88
3	A	402	ARD	O13-C39-C38	-2.85	103.97	110.01
3	A	402	ARD	O3-C7-N1	-2.78	116.28	122.15
3	A	402	ARD	C24-C23-C22	2.72	120.88	114.67
3	A	402	ARD	C37-C38-C39	-2.66	108.18	111.00
3	A	402	ARD	C40-C39-C38	-2.65	106.54	110.97
3	A	402	ARD	C33-O1-C1	-2.47	114.18	117.89
3	A	402	ARD	C19-C18-C17	-2.31	118.75	123.47
3	A	402	ARD	C46-C28-C29	2.17	129.17	123.45
3	A	402	ARD	C35-C36-C37	-2.13	107.11	111.75
3	A	402	ARD	C45-C24-C23	-2.12	107.10	111.55
3	A	402	ARD	C44-C22-C21	2.04	115.36	110.86

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	402	ARD	C14-C15-C16-C17

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	ARD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	107/107 (100%)	0.26	3 (2%) 53 52	20, 33, 52, 84	0
2	B	94/94 (100%)	0.13	2 (2%) 63 63	21, 32, 52, 70	1 (1%)
All	All	201/201 (100%)	0.20	5 (2%) 57 56	20, 32, 52, 84	1 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	18	ARG	3.5
1	A	99	PHE	2.5
1	A	13	ARG	2.4
2	B	165	ARG	2.4
2	B	149	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	ARD	A	402	69/69	0.93	0.15	16,23,31,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.