



# Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2023 – 12:21 PM EDT

PDB ID : 8FBT  
Title : Crystal structure of *Cryptosporidium parvum* N-myristoyltransferase with bound myristoyl-CoA  
Authors : Staker, B.L.; Fenwick, M.; Seattle Structural Genomics Center for Infectious Disease (SSGCID)  
Deposited on : 2022-11-30  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.33  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.33


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	432	
1	B	432	

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 14061 atoms, of which 6781 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycylpeptide N-tetradecanoyltransferase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	B	411	6711	2186	3328	567	609	21	0	8	0
1	A	418	6721	2201	3307	573	618	22	0	11	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	35	GLY	-	expression tag	UNP Q5CV46
B	36	PRO	-	expression tag	UNP Q5CV46
B	37	GLY	-	expression tag	UNP Q5CV46
B	38	SER	-	expression tag	UNP Q5CV46
B	310	ALA	LYS	engineered mutation	UNP Q5CV46
B	311	ALA	GLU	engineered mutation	UNP Q5CV46
A	35	GLY	-	expression tag	UNP Q5CV46
A	36	PRO	-	expression tag	UNP Q5CV46
A	37	GLY	-	expression tag	UNP Q5CV46
A	38	SER	-	expression tag	UNP Q5CV46
A	310	ALA	LYS	engineered mutation	UNP Q5CV46
A	311	ALA	GLU	engineered mutation	UNP Q5CV46

- Molecule 2 is TETRADECANOYL-COA (three-letter code: MYA) (formula: C<sub>35</sub>H<sub>62</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
			Total	C	H	N	O	P			S
2	B	1	Total	C	H	N	O	P	S	0	0
			121	35	58	7	17	3	1		
2	A	1	Total	C	H	N	O	P	S	0	0
			121	35	58	7	17	3	1		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	H			O
3	B	1	Total	C	H	O	0	0
			10	2	6	2		
3	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
4	B	1	17	4	10	3	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
5	A	1	14	3	8	3	0	1


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	185	Total 185	O 185	0	13
6	A	151	Total 151	O 151	0	19


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycylpeptide N-tetradecanoyltransferase



- Molecule 1: Glycylpeptide N-tetradecanoyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	65.43Å 105.34Å 71.12Å 90.00° 112.48° 90.00°	Depositor
Resolution (Å)	49.83 – 2.20 49.83 – 2.19	Depositor EDS
% Data completeness (in resolution range)	99.0 (49.83-2.20) 99.0 (49.83-2.19)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.98 (at 2.18Å)	Xtrriage
Refinement program	PHENIX 1.20.1-4487	Depositor
R, $R_{free}$	0.164 , 0.217 0.164 , 0.215	Depositor DCC
$R_{free}$ test set	2251 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.8	Xtrriage
Anisotropy	0.150	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 74.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	14061	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.24% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MYA, EDO, PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.61	0/3491	0.68	0/4731
1	B	0.68	0/3462	0.75	1/4686 (0.0%)
All	All	0.65	0/6953	0.72	1/9417 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	429	LEU	CB-CG-CD2	-5.19	102.17	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3414	3307	3296	39	0
1	B	3383	3328	3321	31	0
2	A	63	58	56	0	0
2	B	63	58	57	3	0
3	A	4	6	6	0	0
3	B	4	6	6	0	0
4	B	7	10	10	0	0
5	A	6	8	8	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	151	0	0	8	0
6	B	185	0	0	2	0
All	All	7280	6781	6760	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:54:ASP:N	1:B:54:ASP:OD1	2.15	0.80
1:A:109:TYR:O	1:A:110:VAL:HG23	1.87	0.72
1:B:188:LEU:HD23	1:B:192[B]:LEU:HD13	1.75	0.68
1:A:226:TRP:CZ2	1:A:439:ILE:HD12	2.30	0.66
1:B:221:THR:HG21	1:B:461:VAL:HG21	1.81	0.63
1:A:221:THR:HG21	1:A:461:VAL:HG21	1.85	0.59
2:B:501:MYA:H8A	6:B:757:HOH:O	2.03	0.57
1:A:430:GLN:NE2	6:A:602:HOH:O	2.23	0.57
1:B:377:VAL:HG23	1:B:377:VAL:O	2.08	0.54
1:A:234:LYS:HD3	1:A:418:ASP:OD1	2.09	0.53
1:B:232:VAL:HG21	1:B:253:ILE:HA	1.91	0.53
1:A:327:GLN:NE2	1:A:328:ASP:OD1	2.42	0.52
1:B:229:ILE:HG23	1:B:235:LEU:HD12	1.92	0.51
1:A:88[A]:CYS:HG	1:A:101:THR:HG1	1.56	0.51
1:A:230:ILE:O	6:A:601[B]:HOH:O	2.19	0.50
1:A:308:HIS:ND1	6:A:607:HOH:O	2.34	0.50
1:A:256:HIS:CD2	6:A:656:HOH:O	2.65	0.50
1:A:127:ARG:HH11	1:A:127:ARG:HG2	1.76	0.50
1:A:38:SER:N	6:A:611:HOH:O	2.45	0.50
1:A:287:MET:HG2	6:A:640:HOH:O	2.11	0.50
1:B:222:ILE:CD1	1:B:445:ARG:HG3	2.42	0.49
1:B:345:PHE:HB3	1:B:354:THR:HG21	1.95	0.49
1:B:42:HIS:ND1	2:B:501:MYA:O7A	2.46	0.48
1:A:169[A]:MET:HE1	1:A:207:GLN:HB2	1.94	0.48
1:A:393:PHE:HE2	1:A:426:MET:HE3	1.78	0.48
1:A:258:ILE:HG23	1:A:259:PRO:HD2	1.95	0.48
1:B:76:ILE:HD11	1:B:85:TRP:CD1	2.49	0.47
1:A:61:SER:O	1:A:454:PRO:HD3	2.13	0.47
1:B:314:LYS:HG2	1:B:317:TYR:HE2	1.79	0.46
1:B:109:TYR:CD1	2:B:501:MYA:H6A	2.50	0.46
1:A:262:LYS:O	1:A:406:ASN:ND2	2.49	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:350:LYS:NZ	6:A:609[A]:HOH:O	2.42	0.45
1:A:255:TYR:O	1:A:436:LYS:HE3	2.16	0.45
1:A:393:PHE:CE2	1:A:426:MET:HE3	2.52	0.45
1:B:157:ALA:HA	1:B:173:VAL:O	2.18	0.44
1:B:385:THR:HG23	1:B:385:THR:O	2.18	0.44
1:B:84:SER:HB3	1:B:146:ASN:HD21	1.83	0.44
1:B:143:VAL:HG21	1:B:192[B]:LEU:HD11	2.00	0.44
1:A:231[B]:ASN:O	1:A:235:LEU:HG	2.17	0.44
1:A:232:VAL:HB	1:A:253:ILE:HG12	1.99	0.44
1:B:138:ASN:O	1:B:199:ARG:HD3	2.18	0.43
1:A:124:GLU:H	1:A:124:GLU:CD	2.22	0.43
1:A:209:ILE:HD13	1:A:463[B]:ILE:HD13	2.01	0.43
1:A:221:THR:HG22	1:A:446:TYR:HB2	1.99	0.43
1:B:402:LYS:HE2	1:B:430:GLN:O	2.19	0.43
1:B:226:TRP:CD1	1:B:432:ILE:HD12	2.54	0.43
1:B:107:ASP:HB2	1:B:108:HIS:CD2	2.53	0.42
1:A:225:TYR:CE1	1:A:423:LEU:HG	2.54	0.42
1:A:367:LEU:C	1:A:367:LEU:HD23	2.40	0.42
1:B:212[B]:CYS:SG	1:B:216:ILE:HD11	2.59	0.42
1:A:231[A]:ASN:O	1:A:235:LEU:HG	2.18	0.42
1:A:290:LYS:HD2	1:A:297:MET:HG2	2.01	0.42
1:A:157:ALA:HA	1:A:173:VAL:O	2.19	0.42
1:A:290:LYS:HD2	1:A:297:MET:CG	2.50	0.42
1:B:377:VAL:O	1:B:377:VAL:CG2	2.67	0.42
1:A:107:ASP:HB2	1:A:108:HIS:CD2	2.55	0.42
1:B:429:LEU:HA	1:B:432:ILE:HG12	2.01	0.42
1:B:222:ILE:CD1	1:B:445:ARG:CG	2.98	0.42
1:B:454:PRO:O	1:B:456:ILE:HG23	2.20	0.41
1:B:247:MET:HB2	1:B:251:SER:OG	2.20	0.41
1:B:355:TYR:HB2	1:B:368:PHE:CZ	2.55	0.41
1:A:148:THR:O	1:A:149:LYS:HB2	2.20	0.41
1:B:323:ILE:H	1:B:323:ILE:HG13	1.72	0.41
1:A:145:VAL:O	1:A:149:LYS:N	2.49	0.41
1:A:231[B]:ASN:ND2	6:A:620:HOH:O	2.53	0.41
1:A:429:LEU:HD23	1:A:429:LEU:HA	1.97	0.41
1:A:229:ILE:HG23	1:A:235:LEU:HD12	2.02	0.41
1:B:405:PHE:CD1	1:B:431:VAL:HG21	2.55	0.41
1:B:429:LEU:HD12	1:B:432:ILE:CG1	2.51	0.41
1:A:226:TRP:CH2	1:A:439:ILE:HD12	2.56	0.40
1:B:161:LYS:NZ	6:B:618:HOH:O	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	419/432 (97%)	400 (96%)	18 (4%)	1 (0%)	47	55
1	B	410/432 (95%)	387 (94%)	23 (6%)	0	100	100
All	All	829/864 (96%)	787 (95%)	41 (5%)	1 (0%)	51	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	76	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	370/403 (92%)	360 (97%)	10 (3%)	44	57
1	B	373/403 (93%)	361 (97%)	12 (3%)	39	50
All	All	743/806 (92%)	721 (97%)	22 (3%)	41	53

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	54	ASP
1	B	70	ASP
1	B	93	ILE
1	B	98	PHE
1	B	136	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	147	GLU
1	B	188	LEU
1	B	215	ASN
1	B	237	GLU
1	B	321	ASP
1	B	342	LYS
1	B	457	SER
1	A	38	SER
1	A	81	ASP
1	A	98	PHE
1	A	168	ILE
1	A	252	LEU
1	A	261	ASP
1	A	309	SER
1	A	327	GLN
1	A	361	ASN
1	A	433	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	GOL	A	503[A]	-	5,5,5	0.97	0	5,5,5	1.08	0
3	EDO	A	502	-	3,3,3	0.55	0	2,2,2	0.24	0
4	PEG	B	503	-	6,6,6	0.35	0	5,5,5	0.30	0
2	MYA	A	501	-	50,65,65	2.88	16 (32%)	65,91,91	1.63	16 (24%)
3	EDO	B	502	-	3,3,3	0.51	0	2,2,2	0.30	0
2	MYA	B	501	-	50,65,65	2.98	15 (30%)	65,91,91	1.61	14 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	503[A]	-	-	4/4/4/4	-
3	EDO	A	502	-	-	1/1/1/1	-
4	PEG	B	503	-	-	2/4/4/4	-
2	MYA	A	501	-	-	0/44/80/80	0/3/3/3
3	EDO	B	502	-	-	1/1/1/1	-
2	MYA	B	501	-	-	6/44/80/80	0/3/3/3

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	MYA	C2X-C3X	-11.83	1.26	1.52
2	B	501	MYA	C2X-C3X	-11.53	1.27	1.52
2	A	501	MYA	C3X-C4X	7.17	1.70	1.53
2	B	501	MYA	O4X-C1X	6.66	1.57	1.42
2	B	501	MYA	C3X-C4X	6.63	1.69	1.53
2	B	501	MYA	O4X-C4X	-6.19	1.31	1.45
2	B	501	MYA	C5-N4	6.08	1.47	1.33
2	A	501	MYA	O4X-C4X	-6.08	1.31	1.45
2	B	501	MYA	C9-N8	5.53	1.45	1.33
2	B	501	MYA	C1X-N9A	-5.43	1.31	1.47
2	A	501	MYA	C1X-N9A	-5.23	1.32	1.47
2	A	501	MYA	O4X-C1X	5.15	1.54	1.42
2	A	501	MYA	C5-N4	4.71	1.44	1.33

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	MYA	C6A-N6A	4.49	1.45	1.34
2	A	501	MYA	C9-N8	4.45	1.43	1.33
2	A	501	MYA	C6A-N6A	4.01	1.44	1.34
2	A	501	MYA	C6A-N1A	-3.60	1.33	1.37
2	B	501	MYA	C6A-N1A	-3.59	1.33	1.37
2	B	501	MYA	O2X-C2X	3.05	1.50	1.43
2	A	501	MYA	O2X-C2X	2.94	1.49	1.43
2	A	501	MYA	P2A-O5A	-2.76	1.48	1.56
2	A	501	MYA	O9-C9	-2.43	1.18	1.23
2	A	501	MYA	P2A-O6A	2.42	1.66	1.60
2	A	501	MYA	C2X-C1X	2.42	1.61	1.53
2	B	501	MYA	P2A-O6A	2.35	1.66	1.60
2	B	501	MYA	O9-C9	-2.32	1.18	1.23
2	A	501	MYA	C8A-N9A	-2.27	1.32	1.37
2	A	501	MYA	C5A-N7A	-2.25	1.34	1.39
2	B	501	MYA	O5-C5	-2.11	1.19	1.23
2	B	501	MYA	C5A-N7A	-2.05	1.35	1.39
2	B	501	MYA	C8A-N9A	-2.05	1.33	1.37

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	MYA	C5A-C4A-N3A	-4.77	118.47	127.41
2	B	501	MYA	C2-S1-C2M	4.74	112.59	102.74
2	A	501	MYA	C5A-C4A-N3A	-3.86	120.16	127.41
2	A	501	MYA	C7-C6-C5	3.73	118.56	112.36
2	A	501	MYA	N3A-C2A-N1A	-3.49	122.22	128.82
2	A	501	MYA	C5M-C4M-C3M	-3.41	105.84	112.88
2	B	501	MYA	N9A-C4A-N3A	3.39	134.14	127.09
2	B	501	MYA	N3A-C2A-N1A	-3.36	122.47	128.82
2	B	501	MYA	C2-C3-N4	-2.86	106.42	112.42
2	A	501	MYA	O5-C5-N4	-2.81	117.72	123.01
2	B	501	MYA	O6A-P2A-O3A	2.79	111.35	101.37
2	A	501	MYA	O3X-C3X-C2X	2.66	121.32	111.68
2	B	501	MYA	O4X-C1X-C2X	-2.61	100.95	106.64
2	A	501	MYA	C3X-C2X-C1X	2.58	105.59	99.89
2	B	501	MYA	C2X-C3X-C4X	2.53	107.70	103.22
2	A	501	MYA	N9A-C4A-N3A	2.51	132.31	127.09
2	A	501	MYA	C6-C7-N8	-2.45	106.95	111.90
2	A	501	MYA	O5-C5-C6	2.45	126.49	122.02
2	A	501	MYA	O5X-P1A-O1A	2.43	111.94	106.15
2	B	501	MYA	C8A-N7A-C5A	2.39	108.57	104.24

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	MYA	O3X-C3X-C4X	-2.29	101.80	110.08
2	B	501	MYA	C6-C7-N8	-2.15	107.55	111.90
2	A	501	MYA	C2-C3-N4	-2.11	107.99	112.42
2	B	501	MYA	C4A-C5A-N7A	-2.10	106.55	110.39
2	A	501	MYA	C2A-N1A-C6A	2.08	120.41	113.17
2	B	501	MYA	C2A-N3A-C4A	2.08	120.81	113.53
2	B	501	MYA	C14-C11-C12	2.06	111.60	108.23
2	A	501	MYA	C8A-N7A-C5A	2.05	107.95	104.24
2	B	501	MYA	C3X-C2X-C1X	2.03	104.39	99.89
2	A	501	MYA	O6A-P2A-O3A	2.00	108.54	101.37

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	501	MYA	S1-C2-C3-N4
5	A	503[A]	GOL	C1-C2-C3-O3
4	B	503	PEG	O2-C3-C4-O4
5	A	503[A]	GOL	O1-C1-C2-C3
5	A	503[A]	GOL	O2-C2-C3-O3
2	B	501	MYA	CAM-CBM-CCM-CDM
3	B	502	EDO	O1-C1-C2-O2
5	A	503[A]	GOL	O1-C1-C2-O2
2	B	501	MYA	C12-O6A-P2A-O3A
2	B	501	MYA	C12-O6A-P2A-O4A
2	B	501	MYA	C3M-C4M-C5M-C6M
2	B	501	MYA	C5M-C6M-C7M-C8M
4	B	503	PEG	O1-C1-C2-O2
3	A	502	EDO	O1-C1-C2-O2

There are no ring outliers.


1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	MYA	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	418/432 (96%)	0.46	46 (11%) 5 4	31, 65, 121, 158	0
1	B	411/432 (95%)	0.03	16 (3%) 39 37	27, 51, 92, 126	1 (0%)
All	All	829/864 (95%)	0.25	62 (7%) 14 13	27, 58, 111, 158	1 (0%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	238	ALA	13.1
1	A	379	ASN	7.7
1	A	378	ILE	7.0
1	A	252	LEU	6.6
1	A	377	VAL	6.6
1	A	242	GLY	6.3
1	A	248	THR	6.2
1	A	235	LEU	6.2
1	A	241	ILE	6.2
1	A	240	PHE	5.9
1	A	236	LEU	5.7
1	A	239	GLY	5.1
1	A	326	LEU	4.8
1	A	82	GLY	4.6
1	A	320	LEU	4.4
1	A	244	PRO	4.4
1	A	322	LYS	4.2
1	A	323	ILE	4.2
1	A	147	GLU	4.0
1	A	246	ASN	4.0
1	A	247	MET	4.0
1	B	42	HIS	3.9
1	A	249	MET	3.9
1	A	325	ASP	3.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	46	ASN	3.5
1	B	244	PRO	3.4
1	A	243	ILE	3.4
1	A	316	ALA	3.4
1	B	186	LYS	3.4
1	A	321	ASP	3.2
1	A	301	ILE	3.0
1	A	255	TYR	2.9
1	A	324	GLU	2.9
1	B	81	ASP	2.9
1	A	312	LEU	2.8
1	B	185	SER	2.7
1	A	292	VAL	2.7
1	B	119	PHE	2.6
1	A	145	VAL	2.6
1	A	253	ILE	2.6
1	A	307	ASN	2.5
1	B	187	ARG	2.5
1	B	145	VAL	2.5
1	B	183	LEU	2.5
1	A	115	SER	2.4
1	A	234	LYS	2.4
1	A	429	LEU	2.4
1	A	318	MET	2.4
1	B	384	PRO	2.4
1	A	245	ARG	2.3
1	A	81	ASP	2.3
1	A	293	SER	2.2
1	A	229	ILE	2.2
1	B	246	ASN	2.2
1	A	168	ILE	2.2
1	B	83	PHE	2.2
1	A	317	TYR	2.1
1	B	43	LYS	2.1
1	A	328	ASP	2.1
1	B	247	MET	2.1
1	B	184	ARG	2.0
1	A	308	HIS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MYA	B	501	63/63	0.79	0.25	36,89,125,139	0
3	EDO	A	502	4/4	0.80	0.13	73,88,94,98	0
5	GOL	A	503[A]	6/6	0.81	0.17	56,76,92,94	0
3	EDO	B	502	4/4	0.82	0.10	57,74,90,92	0
4	PEG	B	503	7/7	0.88	0.15	57,72,88,88	0
2	MYA	A	501	63/63	0.97	0.13	37,54,68,81	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.