



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 6, 2024 – 09:03 AM EST

PDB ID : 2FCU  
Title : SyrB2 with alpha-ketoglutarate  
Authors : Blasiak, L.C.; Drennan, C.L.  
Deposited on : 2005-12-12  
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	
1	B	313	

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5368 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called syringomycin biosynthesis enzyme 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	309	Total	C	N	O	S	0	5	0
			2494	1572	427	483	12			
1	B	299	Total	C	N	O	S	0	3	0
			2386	1509	409	456	12			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	cloning artifact	GB 5748808
A	-1	SER	-	cloning artifact	GB 5748808
A	0	HIS	-	cloning artifact	GB 5748808
B	-2	GLY	-	cloning artifact	GB 5748808
B	-1	SER	-	cloning artifact	GB 5748808
B	0	HIS	-	cloning artifact	GB 5748808

- Molecule 2 is ((2R,3S,4S,5S)-3,4-DIHYDROXY-5-(HYDROXYMETHYL)-5-((2R,3S,4S,5S,6R)-3,4,5-TRIHYDROXY-6-METHOXY-TETRAHYDRO-2H-PYRAN-2-YLOXY)-TETRAHYDROFURAN-2-YL)METHYL NONANOATE (three-letter code: DSU) (formula: C<sub>22</sub>H<sub>40</sub>O<sub>12</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	1
			34	22	12		

- Molecule 3 is 2-OXOGLUTARIC ACID (three-letter code: AKG) (formula:  $C_5H_6O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	5	5		
3	B	1	Total	C	O	0	0
			10	5	5		


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	252	Total 252	O 252	0	0
4	B	182	Total 182	O 182	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: syringomycin biosynthesis enzyme 2



- Molecule 1: syringomycin biosynthesis enzyme 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.26Å 90.03Å 128.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.80 – 1.60 45.02 – 1.60	Depositor EDS
% Data completeness (in resolution range)	98.6 (42.80-1.60) 98.6 (45.02-1.60)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.98 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.188 , 0.212 0.193 , 0.215	Depositor DCC
$R_{free}$ test set	5846 reflections (6.63%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.0	Xtrriage
Anisotropy	0.155	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 45.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	5368	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DSU, AKG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/2588	0.73	3/3508 (0.1%)
1	B	0.52	0/2467	0.68	0/3348
All	All	0.55	0/5055	0.71	3/6856 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	296	ARG	NE-CZ-NH1	6.40	123.50	120.30
1	A	73	ASP	CB-CG-OD1	5.83	123.55	118.30
1	A	296	ARG	NE-CZ-NH2	-5.39	117.60	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2494	0	2332	11	0
1	B	2386	0	2211	20	0
2	A	34	0	5	0	0
3	A	10	0	4	0	0
3	B	10	0	4	0	0
4	A	252	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	182	0	0	1	0
All	All	5368	0	4556	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:SER:OG	1:A:31[B]:GLU:HG3	1.73	0.88
1:B:28:SER:OG	1:B:31[B]:GLU:HG3	1.79	0.83
1:A:153:ILE:H	1:A:220:GLN:HE21	1.33	0.75
1:B:28:SER:OG	1:B:31[B]:GLU:CG	2.35	0.74
1:A:220:GLN:HE22	1:B:225:GLN:HE22	1.35	0.71
1:A:159:GLN:HE22	1:A:193:ARG:HH22	1.39	0.70
1:B:159:GLN:HE22	1:B:193:ARG:HH22	1.41	0.68
1:B:97:LEU:HD21	1:B:291:THR:HG21	1.75	0.68
1:B:63:ILE:HD12	1:B:66:TYR:CD2	2.36	0.59
1:B:69:HIS:HE1	4:B:904:HOH:O	1.87	0.56
1:B:258[A]:SER:HB2	1:B:291:THR:HB	1.87	0.55
1:B:8:THR:H	1:B:11:GLN:HE21	1.54	0.54
1:A:299:THR:OG1	1:A:300:HIS:HD2	1.91	0.54
1:A:34:GLU:HG3	1:A:38:ARG:NH2	2.24	0.52
1:B:159:GLN:NE2	1:B:193:ARG:HH22	2.07	0.52
1:A:300:HIS:HE1	4:A:1148:HOH:O	1.93	0.50
1:A:159:GLN:NE2	1:A:193:ARG:HH22	2.06	0.49
1:B:285:GLN:OE1	1:B:291:THR:HG23	2.12	0.49
1:B:285:GLN:OE1	1:B:291:THR:CG2	2.61	0.49
1:B:28:SER:OG	1:B:31[B]:GLU:HG2	2.12	0.49
1:B:191:VAL:HG22	1:B:193:ARG:HG3	1.97	0.47
1:B:8:THR:H	1:B:11:GLN:NE2	2.13	0.46
1:A:57:ILE:N	1:A:57:ILE:HD12	2.32	0.44
1:B:67:ASP:HA	1:B:69:HIS:CD2	2.54	0.42
1:B:129:GLN:HE22	1:B:269:ILE:HG23	1.85	0.42
1:B:54:LEU:O	1:B:55:ASP:HB2	2.19	0.42
1:B:96:VAL:HB	1:B:287:ILE:HD12	2.00	0.42
1:A:310:VAL:O	1:A:310:VAL:HG12	2.18	0.41
1:A:8:THR:H	1:A:11:GLN:NE2	2.19	0.41
1:B:131:ILE:HD12	1:B:264:PRO:HG3	2.02	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/313 (100%)	308 (99%)	4 (1%)	0	100	100
1	B	296/313 (95%)	291 (98%)	5 (2%)	0	100	100
All	All	608/626 (97%)	599 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/265 (100%)	261 (98%)	4 (2%)	65	44
1	B	249/265 (94%)	243 (98%)	6 (2%)	49	24
All	All	514/530 (97%)	504 (98%)	10 (2%)	55	34

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	A	42	ARG
1	A	152	ASN
1	A	179	GLU
1	B	33	LYS
1	B	152	ASN
1	B	191	VAL
1	B	248	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	291	THR
1	B	293	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	152	ASN
1	A	159	GLN
1	A	220	GLN
1	A	300	HIS
1	B	11	GLN
1	B	18	ASN
1	B	69	HIS
1	B	129	GLN
1	B	152	ASN
1	B	159	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	AKG	A	897	-	9,9,9	2.05	1 (11%)	11,11,11	1.44	2 (18%)
3	AKG	B	898	-	9,9,9	1.91	1 (11%)	11,11,11	1.43	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	AKG	A	897	-	-	1/9/9/9	-
3	AKG	B	898	-	-	0/9/9/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	897	AKG	C2-C1	-5.54	1.46	1.53
3	B	898	AKG	C2-C1	-4.81	1.47	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	897	AKG	C3-C2-C1	3.64	122.73	115.97
3	B	898	AKG	C3-C2-C1	3.60	122.65	115.97
3	A	897	AKG	O5-C2-C1	-2.05	116.45	119.43

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	897	AKG	O2-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	309/313 (98%)	0.13	11 (3%) 42 40	8, 14, 29, 36	0
1	B	299/313 (95%)	0.46	36 (12%) 4 3	10, 19, 38, 48	0
All	All	608/626 (97%)	0.29	47 (7%) 13 11	8, 17, 34, 48	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	182	ALA	7.8
1	B	138	PHE	6.5
1	B	272	TYR	6.5
1	A	2	SER	6.2
1	B	181	ASP	5.5
1	B	273	GLY	5.4
1	A	180	PRO	4.7
1	B	180	PRO	4.5
1	B	274	GLY	4.4
1	B	186	VAL	4.3
1	B	124	ALA	4.3
1	B	184	ASN	4.2
1	A	55	ASP	4.1
1	B	61	THR	3.8
1	A	182	ALA	3.8
1	A	124	ALA	3.5
1	B	125	SER	3.4
1	B	177	THR	3.4
1	B	183	ASN	3.4
1	B	55	ASP	3.3
1	A	53	ASP	3.3
1	B	196	PHE	3.1
1	B	185	SER	3.1
1	B	178	TYR	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	187	VAL	3.0
1	A	310	VAL	3.0
1	B	126	GLY	2.9
1	A	185	SER	2.9
1	B	290	GLU	2.9
1	B	208	ASN	2.9
1	B	198	TYR	2.8
1	B	123	ASN	2.7
1	B	271	GLU	2.7
1	B	265	ASP	2.7
1	B	179	GLU	2.6
1	A	125	SER	2.6
1	B	3	LYS	2.5
1	B	175	ARG	2.5
1	A	272	TYR	2.5
1	B	133	PRO	2.4
1	B	53	ASP	2.4
1	B	54	LEU	2.4
1	B	289	ASP	2.4
1	B	293	GLU	2.4
1	A	58	SER	2.2
1	B	267	ASP	2.1
1	B	207	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
-----	------	-------	-----	-------	------	-----	-----------------------------	-------

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	DSU	A	896[N]	34/34	0.81	0.14	17,25,31,32	0
3	AKG	B	898	10/10	0.90	0.12	14,18,22,29	0
3	AKG	A	897	10/10	0.94	0.12	10,13,21,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.