



# Full wwPDB X-ray Structure Validation Report ⓘ

May 18, 2020 – 02:34 pm BST

PDB ID : 2FE8  
Title : SARS coronavirus papain-like protease: structure of a viral deubiquitinating enzyme  
Authors : Ratia, K.; Santarsiero, B.D.; Mesecar, A.D.  
Deposited on : 2005-12-15  
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11



## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7978 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Replicase polypeptide 1ab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	315	2482	1574	412	478	18	0	0	0
1	B	314	2473	1568	411	476	18	0	0	0
1	C	313	2466	1564	410	474	18	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	INITIATING METHIONINE	UNP P59641
B	1	MET	-	INITIATING METHIONINE	UNP P59641
C	1	MET	-	INITIATING METHIONINE	UNP P59641

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		
2	C	1	Total	Zn	0	0
			1	1		

- Molecule 3 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	3	Total	Br	0	0
			3	3		
3	A	3	Total	Br	0	0
			3	3		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	3	Total	Br	0	0
			3	3		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	206	Total	O	0	0
			206	206		
5	B	177	Total	O	0	0
			177	177		
5	C	147	Total	O	0	0
			147	147		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Replicase polyprotein 1ab



- Molecule 1: Replicase polyprotein 1ab



- Molecule 1: Replicase polyprotein 1ab



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	142.87Å 103.06Å 91.73Å 90.00° 111.15° 90.00°	Depositor
Resolution (Å)	20.00 – 1.85 45.24 – 1.84	Depositor EDS
% Data completeness (in resolution range)	97.7 (20.00-1.85) 96.9 (45.24-1.84)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.76 (at 1.84Å)	Xtrriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.201 , 0.229 0.200 , 0.228	Depositor DCC
$R_{free}$ test set	5203 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	25.2	Xtrriage
Anisotropy	0.176	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 45.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.008 for $-1/2^*h+1/2^*k+1, 1/2^*h-1/2^*k+1, 1/2^*h+1/2^*k$ 0.014 for $-1/2^*h-1/2^*k+1, -1/2^*h-1/2^*k-1, 1/2^*h-1/2^*k$	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7978	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.51	0/2537	0.70	0/3441
1	B	0.47	0/2528	0.67	0/3430
1	C	0.46	0/2521	0.66	0/3420
All	All	0.48	0/7586	0.68	0/10291

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	269	TYR	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2482	0	2425	34	0
1	B	2473	0	2414	31	0
1	C	2466	0	2407	29	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	3	0	0	1	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
4	C	5	0	0	0	0
5	A	206	0	0	2	0
5	B	177	0	0	3	0
5	C	147	0	0	4	0
All	All	7978	0	7246	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:VAL:HG13	1:A:223:ILE:HD11	1.53	0.88
1:C:39:GLY:HA2	1:C:85:MET:HE1	1.64	0.79
1:C:39:GLY:HA2	1:C:85:MET:CE	2.14	0.77
1:B:249:PRO:HG3	1:B:302:THR:CG2	2.16	0.75
1:A:229:ARG:HG3	1:A:230:ASP:H	1.55	0.71
1:A:130:ALA:O	1:A:134:GLN:HG3	1.92	0.69
1:A:23:ASP:HB3	1:A:26:MET:HE3	1.77	0.66
1:B:78:GLU:HG3	1:C:78:GLU:HG3	1.78	0.65
1:B:249:PRO:HG3	1:B:302:THR:HG22	1.78	0.65
1:A:45:ILE:HD11	1:A:48:HIS:NE2	2.12	0.64
1:B:307:LYS:HE3	5:B:441:HOH:O	1.96	0.64
1:B:270:GLN:HG2	1:C:208:TYR:CE1	2.33	0.64
1:A:23:ASP:HB3	1:A:26:MET:CE	2.29	0.62
1:A:229:ARG:HG3	1:A:230:ASP:N	2.13	0.62
1:A:223:ILE:HD13	1:A:233:GLN:HB2	1.81	0.62
1:B:268:ASN:O	1:B:272:GLY:N	2.34	0.61
1:C:223:ILE:HD13	1:C:233:GLN:HB2	1.84	0.59
1:B:270:GLN:HG2	1:C:208:TYR:CD1	2.37	0.59
1:C:265:TYR:OH	1:C:272:GLY:HA3	2.01	0.59

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:GLN:HG3	1:A:312:THR:HG22	1.83	0.59
1:A:191:LYS:HB2	1:A:229:ARG:HH12	1.68	0.58
1:C:191:LYS:HE3	1:C:229:ARG:HD2	1.85	0.57
1:A:204:GLU:HG3	5:A:486:HOH:O	2.05	0.56
1:C:265:TYR:CZ	1:C:272:GLY:HA3	2.41	0.56
1:C:58:VAL:HG22	5:C:386:HOH:O	2.05	0.55
1:C:191:LYS:HE3	1:C:229:ARG:CD	2.36	0.55
1:B:66:ARG:HD3	5:B:376:HOH:O	2.07	0.55
1:B:249:PRO:HG3	1:B:302:THR:HG21	1.89	0.55
1:A:315:ILE:OXT	1:A:315:ILE:HD12	2.07	0.54
1:A:203:VAL:HG12	1:A:207:MET:HE2	1.89	0.54
1:A:23:ASP:OD2	1:A:26:MET:HE2	2.08	0.53
1:A:192:HIS:HB2	1:A:229:ARG:HH22	1.74	0.53
1:A:45:ILE:HG12	1:A:46:LYS:O	2.08	0.53
1:A:134:GLN:HA	3:A:318:BR:BR	2.63	0.52
1:A:42:VAL:HB	1:A:45:ILE:CG2	2.40	0.52
1:B:23:ASP:HB3	1:B:26:MET:HE3	1.92	0.51
1:B:204:GLU:HG3	5:B:432:HOH:O	2.11	0.51
1:B:82:GLY:HA3	1:C:76:LEU:CD1	2.41	0.51
1:C:157:ASN:HB3	5:C:443:HOH:O	2.10	0.51
1:C:249:PRO:HG3	1:C:302:THR:CG2	2.41	0.51
1:B:283:LEU:HD12	1:B:296:GLU:HA	1.93	0.51
1:A:188:VAL:CG1	1:A:223:ILE:HD11	2.34	0.50
1:B:49:VAL:HG13	1:B:52:GLU:OE2	2.12	0.49
1:B:4:LYS:O	1:B:24:MET:HG2	2.12	0.49
1:C:39:GLY:HA2	1:C:85:MET:HE3	1.94	0.49
1:C:187:ASN:OD1	1:C:197:THR:HG22	2.12	0.49
1:B:268:ASN:OD1	1:B:269:TYR:HD1	1.96	0.49
1:C:189:VAL:HG22	1:C:195:GLN:HB3	1.95	0.49
1:C:223:ILE:CD1	1:C:233:GLN:HB2	2.42	0.48
1:B:152:ILE:HG12	1:B:173:LEU:HD11	1.95	0.48
1:A:23:ASP:CB	1:A:26:MET:CE	2.92	0.48
1:B:266:THR:O	1:B:272:GLY:HA2	2.13	0.48
1:B:6:ILE:HG13	1:B:22:VAL:HG22	1.96	0.47
1:B:268:ASN:OD1	1:B:269:TYR:N	2.48	0.47
1:B:4:LYS:C	1:B:24:MET:HG2	2.34	0.47
1:A:23:ASP:CB	1:A:26:MET:HE2	2.44	0.47
1:B:78:GLU:HG3	1:C:78:GLU:CG	2.45	0.47
1:B:171:THR:O	1:B:175:GLN:HG2	2.14	0.47
1:C:249:PRO:HG3	1:C:302:THR:HG21	1.96	0.46
1:A:42:VAL:O	1:A:45:ILE:HG22	2.16	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:TYR:CE2	1:A:141:ARG:HD2	2.50	0.46
1:C:66:ARG:HD2	5:C:437:HOH:O	2.17	0.46
1:C:191:LYS:HE3	1:C:229:ARG:CG	2.47	0.44
1:A:171:THR:O	1:A:175:GLN:HG3	2.18	0.43
1:B:148:PHE:CE2	1:B:152:ILE:HD11	2.52	0.43
1:C:227:CYS:SG	1:C:229:ARG:HB2	2.57	0.43
1:C:32:PHE:HB3	1:C:58:VAL:HG21	2.00	0.43
1:A:267:GLY:HA2	1:B:224:PRO:HG3	2.00	0.43
1:C:196:LYS:N	1:C:196:LYS:HD2	2.34	0.43
1:A:103:THR:OG1	1:A:143:GLY:HA2	2.18	0.43
1:A:214:TYR:OH	1:A:238:GLN:HG2	2.20	0.41
1:A:229:ARG:HH11	1:A:229:ARG:HG2	1.85	0.41
1:C:172:HIS:HB2	5:C:428:HOH:O	2.20	0.41
1:A:128:PHE:O	1:A:134:GLN:HG2	2.21	0.41
1:A:197:THR:HG21	1:A:315:ILE:HD11	2.03	0.41
1:A:45:ILE:HG12	1:A:46:LYS:N	2.35	0.41
1:A:127:LYS:HE3	1:A:178:ASN:HD22	1.86	0.40
1:A:283:LEU:HB2	1:A:294:MET:O	2.22	0.40
1:B:121:LEU:HD11	1:B:173:LEU:HD13	2.03	0.40
1:B:23:ASP:HB3	1:B:26:MET:CE	2.51	0.40
1:A:307:LYS:HE3	5:A:473:HOH:O	2.20	0.40
1:C:165:ASP:HB3	1:C:168:GLU:HB3	2.03	0.40
1:B:137:TYR:CE2	1:B:141:ARG:HD2	2.56	0.40
1:B:77:ASP:OD1	1:B:83:ARG:NH2	2.53	0.40
1:C:260:LEU:HG	1:C:307:LYS:HG3	2.03	0.40
1:B:90:HIS:HB2	1:B:160:VAL:HG21	2.04	0.40
1:B:270:GLN:O	1:C:200:LEU:HD11	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/315 (99%)	307 (98%)	6 (2%)	0	100	100
1	B	312/315 (99%)	304 (97%)	7 (2%)	1 (0%)	41	26
1	C	311/315 (99%)	301 (97%)	10 (3%)	0	100	100
All	All	936/945 (99%)	912 (97%)	23 (2%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	269	TYR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/272 (100%)	267 (98%)	5 (2%)	59	45
1	B	271/272 (100%)	265 (98%)	6 (2%)	52	36
1	C	270/272 (99%)	264 (98%)	6 (2%)	52	36
All	All	813/816 (100%)	796 (98%)	17 (2%)	53	38

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	ILE
1	A	81	LEU
1	A	229	ARG
1	A	263	ASN
1	A	315	ILE
1	B	66	ARG
1	B	251	GLU
1	B	263	ASN
1	B	264	GLU
1	B	270	GLN
1	B	285	ARG
1	C	173	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	196	LYS
1	C	255	GLN
1	C	263	ASN
1	C	269	TYR
1	C	313	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 12 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	SO4	C	320	-	4,4,4	0.27	0	6,6,6	0.18	0
4	SO4	B	320	-	4,4,4	0.24	0	6,6,6	0.19	0
4	SO4	A	320	-	4,4,4	0.27	0	6,6,6	0.18	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	315/315 (100%)	-0.31	9 (2%) 51 50	15, 25, 40, 69	0
1	B	314/315 (99%)	-0.20	7 (2%) 62 61	15, 27, 41, 73	0
1	C	313/315 (99%)	-0.14	10 (3%) 47 45	18, 28, 53, 71	0
All	All	942/945 (99%)	-0.22	26 (2%) 53 52	15, 26, 46, 73	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	271	CYS	8.7
1	A	226	VAL	7.1
1	C	191	LYS	5.0
1	B	269	TYR	4.5
1	B	270	GLN	4.0
1	A	228	GLY	3.9
1	C	1	MET	3.9
1	B	314	THR	3.6
1	C	3	VAL	3.2
1	A	230	ASP	3.1
1	B	78	GLU	3.0
1	B	272	GLY	2.7
1	A	227	CYS	2.7
1	C	226	VAL	2.6
1	A	138	TYR	2.6
1	C	269	TYR	2.6
1	C	313	THR	2.6
1	A	225	CYS	2.5
1	C	229	ARG	2.5
1	A	229	ARG	2.4
1	C	2	GLU	2.4
1	A	70	PHE	2.3
1	C	192	HIS	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	267	GLY	2.3
1	A	191	LYS	2.3
1	C	228	GLY	2.3

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	A	316	1/1	0.92	0.06	58,58,58,58	0
2	ZN	C	316	1/1	0.95	0.04	58,58,58,58	0
4	SO4	B	320	5/5	0.97	0.10	43,44,45,47	0
3	BR	A	317	1/1	0.98	0.03	36,36,36,36	0
3	BR	C	317	1/1	0.98	0.04	39,39,39,39	0
3	BR	C	319	1/1	0.98	0.09	51,51,51,51	0
3	BR	B	319	1/1	0.99	0.08	54,54,54,54	0
4	SO4	A	320	5/5	0.99	0.09	36,36,38,39	0
2	ZN	B	316	1/1	0.99	0.07	27,27,27,27	0
3	BR	A	318	1/1	0.99	0.07	54,54,54,54	0
3	BR	B	317	1/1	0.99	0.03	40,40,40,40	0
3	BR	A	319	1/1	0.99	0.12	44,44,44,44	0
4	SO4	C	320	5/5	0.99	0.07	38,39,40,40	0
3	BR	B	318	1/1	0.99	0.03	41,41,41,41	0
3	BR	C	318	1/1	1.00	0.04	38,38,38,38	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.